


This article was downloaded by: [Australian National University]

On: 30 September 2013, At: 16:06

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Molecular Physics: An International Journal at the Interface Between Chemistry and Physics

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tmph20>

Distribution of $r_{12} \cdot p_{12}$ in quantum systems

Yves A. Bernard^a, Pierre-François Loos^a & Peter M.W. Gill^a

^a Research School of Chemistry, Australian National University, Canberra, ACT, 0200, Australia

Accepted author version posted online: 13 Jun 2013. Published online: 08 Jul 2013.

To cite this article: Yves A. Bernard, Pierre-François Loos & Peter M.W. Gill (2013) Distribution of $r_{12} \cdot p_{12}$ in quantum systems, *Molecular Physics: An International Journal at the Interface Between Chemistry and Physics*, 111:16-17, 2414-2426, DOI: [10.1080/00268976.2013.811302](https://doi.org/10.1080/00268976.2013.811302)

To link to this article: <http://dx.doi.org/10.1080/00268976.2013.811302>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

INVITED ARTICLE

Distribution of $\mathbf{r}_{12} \cdot \mathbf{p}_{12}$ in quantum systems

Yves A. Bernard, Pierre-François Loos and Peter M.W. Gill*

Research School of Chemistry, Australian National University, Canberra, ACT 0200, Australia

(Received 31 January 2013; final version received 28 May 2013)

We introduce the two-particle probability density $X(x)$ of $x = \mathbf{r}_{12} \cdot \mathbf{p}_{12} = (\mathbf{r}_1 - \mathbf{r}_2) \cdot (\mathbf{p}_1 - \mathbf{p}_2)$. The fundamental equations involved in the derivation of this new intracule $X(x)$, which we call the Posmom intracule, are derived and we show how to derive $X(x)$ from the many-particle wave-function. We contrast it with the Dot intracule [Y.A. Bernard, D.L. Crittenden, and P.M.W. Gill, Phys. Chem. Chem. Phys. **10**, 3447 (2008)] which can be derived from the Wigner distribution and show the relationships between the Posmom intracule and the one-particle Posmom density [Y.A. Bernard, D.L. Crittenden, and P.M.W. Gill, J. Phys. Chem. A **114**, 11984 (2010)]. To illustrate the information provided by the Posmom intracule, we apply this new formalism to various two-electron systems: the three-dimensional parabolic quantum dot, the helium-like ions and the ground and excited states of the helium atom.

Keywords: two-particle density distribution; Wigner distribution; intracule; Posmom

PACS numbers: 31.15.X-; 31.15.ve; 31.15.vj; 81.07.Ta

1. Introduction

Intracules are two-particle density distribution obtained from the spinless second-order reduced density matrix [1]

$$\rho_2 \begin{pmatrix} \mathbf{r}_1, \mathbf{r}'_1 \\ \mathbf{r}_2, \mathbf{r}'_2 \end{pmatrix} = \int \Psi^*(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \dots, \mathbf{r}_N) \times \Psi(\mathbf{r}'_1, \mathbf{r}'_2, \mathbf{r}_3, \dots, \mathbf{r}_N) d\mathbf{r}_3 \cdots d\mathbf{r}_N, \quad (1)$$

where $\Psi(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \dots, \mathbf{r}_N)$ is the N -particle position wave function. Intracules are usually normalised to the number of particle pairs $N(N-1)/2$.

The seminal intracule is the Position intracule

$$P(u) = \int \rho_2 \begin{pmatrix} \mathbf{r}, \mathbf{r} \\ \mathbf{r} + \mathbf{u}, \mathbf{r} + \mathbf{u} \end{pmatrix} d\mathbf{r} d\Omega_u, \quad (2)$$

which was introduced long ago by Coulson and Neilson [2] to study correlation effects in the helium atom. In Equation (2), $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_2$, $u = |\mathbf{u}| \equiv r_{12}$ and Ω_u is the angular part of \mathbf{u} . $P(u)$ gives the probability density for finding two particles separated by a distance u and has been widely studied [3–21].

The corresponding Momentum intracule [22,23] is

$$M(v) = \frac{1}{(2\pi)^3} \int \rho_2 \begin{pmatrix} \mathbf{r}, \mathbf{r} + \mathbf{q} \\ \mathbf{r} + \mathbf{u} + \mathbf{q}, \mathbf{r} + \mathbf{u} \end{pmatrix} \times e^{i\mathbf{q} \cdot \mathbf{v}} d\mathbf{r} d\mathbf{q} d\mathbf{u} d\Omega_v, \quad (3)$$

where $\mathbf{v} = \mathbf{p}_1 - \mathbf{p}_2$, $v = |\mathbf{v}| \equiv p_{12}$ and Ω_v is the angular part of \mathbf{v} . $M(v)$ gives the probability density for finding two particles moving with a relative momentum v .

Starting with the Wigner distribution [24,25], one can construct a family of intracules [15,16], which provide two-electron position and/or momentum information. Within this family, the patriarch is the Omega intracule

$$\Omega(u, v, \omega) = \frac{1}{(2\pi)^3} \int \rho_2 \begin{pmatrix} \mathbf{r}, \mathbf{r} + \mathbf{q} \\ \mathbf{r} + \mathbf{u} + \mathbf{q}, \mathbf{r} + \mathbf{u} \end{pmatrix} \times e^{i\mathbf{q} \cdot \mathbf{v}} \delta(\omega - \theta_{uv}) d\mathbf{r} d\mathbf{q} d\Omega_u d\Omega_v, \quad (4)$$

where $\omega \equiv \theta_{uv}$ is the dynamical angle between the vector \mathbf{u} and \mathbf{v} . $\Omega(u, v, \omega)$ can be interpreted as the *joint quasi-probability density* for u, v and ω . The *quasi* prefix emphasises that $\Omega(u, v, \omega)$ is not a rigorous probability density and, indeed, it may take negative values [24]. Based on the observation of Rassolov [26] that both relative position and relative momentum are important to describe the correlation between pairs of electrons, and because the Omega intracule contains information on both quantities, $\Omega(u, v, \omega)$ has been extensively used in intracule functional theory (IFT) [16,17,27–33].

Appropriate integrations [16] reduce the Omega intracule to lower order intracules such as $P(u)$, $M(v)$ and the Angle intracule [27,28]

$$\Upsilon(\omega) = \int_0^\infty \int_0^\infty \Omega(u, v, \omega) du dv, \quad (5)$$

*Corresponding author. Email: peter.gill@anu.edu.au

which provides information on the angle ω between \mathbf{u} and \mathbf{v} . A similar reduction yields the Dot intracule [17,30]

$$D(x) = \int_0^\infty \int_x^\infty \frac{\Omega(u, z/u, \omega)}{u z \sin \omega} dz du. \quad (6)$$

The variable $x = \mathbf{u} \cdot \mathbf{v} = uv \cos \omega$ combines information on the relative position and momentum of the particles, and it is easy to show that it gives the rate of change of u^2 , i.e.

$$x = \frac{1}{2} \frac{d}{dt} u^2. \quad (7)$$

In this way, x sheds light on the motion of the electrons. For example, $x = 0$ implies that the electrons are moving in such a way that their separation is constant. This could arise, for example, if they were in a circular orbit around their centre of mass.

Although $D(x)$ is usually a non-negative function and has proven useful for understanding electronic behaviour [27] and for estimating electron correlation energies in atomic and molecular systems [29,30], its connection to the Omega suggests that it is not a rigorous probability density. However, in the following section, we show how to derive the *exact* probability distribution of x .

2. The Posmom intracule

We define the Posmom intracule $X(x)$ to be the exact probability density for the variable $x = \mathbf{u} \cdot \mathbf{v}$. It is the two-particle version of the Posmom density $S(s)$ where $s = \mathbf{r} \cdot \mathbf{p}$ [34–36] and, as we have argued that s describes particle trajectories, we now propose that x likewise characterises pair trajectories.

The quantum mechanical operator

$$\bar{s} = -i\hbar \left(\frac{3}{2} + \mathbf{r} \cdot \nabla_{\mathbf{r}} \right) \quad (8)$$

is known to be an unbounded self-adjoint operator [37,38] and its two-particle equivalent is

$$\bar{x} = -2i\hbar \left(\frac{3}{2} + \mathbf{u} \cdot \nabla_{\mathbf{u}} \right), \quad (9)$$

where ∇ is the gradient operator. Both \bar{s} and \bar{x} correspond to quantum mechanical observables.

Following the same approach used in [34] to obtain $S(s)$ from \bar{s} , one can show that the Posmom intracule can be expressed as the Fourier transform

$$X(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{X}(k) e^{ikx} dk \quad (10)$$

of the two-particle hyperbolic autocorrelation function

$$\hat{X}(k) = \int \rho_2 \left(\begin{matrix} \mathbf{r} \\ \mathbf{r} + e^{k\hbar} \mathbf{u} \end{matrix}, \begin{matrix} \mathbf{r} + \sinh(k\hbar) \mathbf{u} \\ \mathbf{r} + \cosh(k\hbar) \mathbf{u} \end{matrix} \right) d\mathbf{r} d\mathbf{u}. \quad (11)$$

This expression can be simplified after defining the intracule density matrix

$$\rho_u(\mathbf{u}, \mathbf{u}') = \int \rho_2 \left(\begin{matrix} \mathbf{U} - \mathbf{u}/2, \mathbf{U} - \mathbf{u}'/2 \\ \mathbf{U} + \mathbf{u}/2, \mathbf{U} + \mathbf{u}'/2 \end{matrix} \right) d\mathbf{U}, \quad (12)$$

where $\mathbf{U} = \mathbf{r}_1 + \mathbf{r}_2$ is the extracule vector, and yields

$$\hat{X}(k) = \int \rho_u(e^{+k\hbar} \mathbf{u}, e^{-k\hbar} \mathbf{u}) d\mathbf{u}. \quad (13)$$

In an entirely analogous way, the Dot intracule can be expressed as the Fourier transform

$$D(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{D}(k) e^{ikx} dk \quad (14)$$

of the f -Dot function [30]

$$\hat{D}(k) = \int \rho_2 \left(\begin{matrix} \mathbf{r} \\ \mathbf{r} + \mathbf{u} + k\hbar \mathbf{u} \end{matrix}, \begin{matrix} \mathbf{r} + k\hbar \mathbf{u} \\ \mathbf{r} + \mathbf{u} \end{matrix} \right) d\mathbf{r} d\mathbf{u}, \quad (15)$$

and the latter can be reduced to

$$\hat{D}(k) = \int \rho_u((1 + k\hbar) \mathbf{u}, (1 - k\hbar) \mathbf{u}) d\mathbf{u}. \quad (16)$$

Comparing (13) with (16) and the Taylor expansion of the exponential function

$$e^{\pm k\hbar} = 1 \pm k\hbar + \frac{k^2}{2} \hbar^2 + \dots \quad (17)$$

reveals that the probability density $D(x)$ derived from the Wigner distribution is a first-order approximation to the exact density $X(x)$. Thus, the quasi-intracule is correct to $O(\hbar)$ and becomes exact in the classical limit $\hbar \rightarrow 0$. Remarkably, one can construct the exact density from the approximate density using the mapping

$$\hat{X}(k) = \frac{\hat{D}(\tanh(k\hbar))}{\cosh^3(k\hbar)}. \quad (18)$$

Table 1 gives the one- and two-particle hyperbolic autocorrelation functions, the corresponding first-order (Wigner) approximations, and the relations between them. In Table 1, the one-particle density matrix is given by

$$\rho_1(\mathbf{r}_1, \mathbf{r}'_1) = \frac{2}{N-1} \int \rho_2 \left(\begin{matrix} \mathbf{r}_1, \mathbf{r}'_1 \\ \mathbf{r}_2, \mathbf{r}_2 \end{matrix} \right) d\mathbf{r}_2. \quad (19)$$

Table 1. One- and two-particle hyperbolic autocorrelation functions.

	Density	Intracule
Dot	$\hat{S}_W(k) = \int \rho_1((1 + k\hbar/2)\mathbf{r}, (1 - k\hbar/2)\mathbf{r}) d\mathbf{r}$	$\hat{D}(k) = \int \rho_u((1 + k\hbar)\mathbf{u}, (1 - k\hbar)\mathbf{u}) d\mathbf{u}$
Posmom	$\hat{S}(k) = \int \rho_1(e^{+k\hbar/2}\mathbf{r}, e^{-k\hbar/2}\mathbf{r}) d\mathbf{r}$	$\hat{X}(k) = \int \rho_u(e^{+k\hbar}\mathbf{u}, e^{-k\hbar}\mathbf{u}) d\mathbf{u}$
Relation	$\hat{S}(k) = \frac{\hat{S}_W(2 \tanh(k\hbar/2))}{\cosh^3(k\hbar/2)}$ $= \hat{S}_W(k) + O(\hbar^2)$	$\hat{X}(k) = \frac{\hat{D}(\tanh(k\hbar))}{\cosh^3(k\hbar)}$ $= \hat{D}(k) + O(\hbar^2)$

If the wave function is expanded in one-electron functions, $\phi_a(\mathbf{r})$, the reduced two-particle density matrix becomes

$$\rho_2\left(\begin{matrix} \mathbf{r}_1, \mathbf{r}'_1 \\ \mathbf{r}_2, \mathbf{r}'_2 \end{matrix}\right) = \sum_{abcd} P_{abcd} \phi_a(\mathbf{r}_1) \phi_b(\mathbf{r}'_1) \phi_c(\mathbf{r}_2) \phi_d(\mathbf{r}'_2), \quad (20)$$

where P_{abcd} is a two-particle density matrix element. In this case, (11) is given by

$$\hat{X}(k) = \sum_{abcd} P_{abcd} [abcd]_{\hat{X}}, \quad (21)$$

where we have introduced the two-particle hyperbolic autocorrelation integral $[abcd]_{\hat{X}}$. For example, if the basis functions are s -type Gaussians, we obtain (in atomic units)

$$[ssss]_{\hat{X}} = \int e^{-\alpha|\mathbf{r}-\mathbf{A}|^2} e^{-\beta|\mathbf{r}+\sinh(k)\mathbf{u}-\mathbf{B}|^2} e^{-\gamma|\mathbf{r}+\exp(k)\mathbf{u}-\mathbf{C}|^2} e^{-\delta|\mathbf{r}+\cosh(k)\mathbf{u}-\mathbf{D}|^2} d\mathbf{r} d\mathbf{u} = \frac{\pi^3}{J^{3/2}} \exp\left[\frac{1}{\xi} \left(\frac{|\mathbf{H}|^2}{J} - F\right)\right], \quad (22)$$

where

$$\xi = \alpha + \beta + \gamma + \delta, \quad (23a)$$

$$J = \xi[\beta \sinh^2(k) + \gamma \exp^2(k) + \delta \cosh^2(k)] - [\beta \sinh(k) + \gamma \exp(k) + \delta \cosh(k)]^2, \quad (23b)$$

$$F = \alpha\beta|\mathbf{A}-\mathbf{B}|^2 + \alpha\gamma|\mathbf{A}-\mathbf{C}|^2 + \alpha\delta|\mathbf{A}-\mathbf{D}|^2 + \beta\gamma|\mathbf{B}-\mathbf{C}|^2 + \beta\delta|\mathbf{B}-\mathbf{D}|^2 + \gamma\delta|\mathbf{C}-\mathbf{D}|^2, \quad (23c)$$

and

$$\mathbf{H} = \sinh(k)\mathbf{G}_B + \exp(k)\mathbf{G}_C + \cosh(k)\mathbf{G}_D, \quad (24a)$$

$$\mathbf{G}_B = \alpha\beta(\mathbf{B}-\mathbf{A}) + \beta\gamma(\mathbf{B}-\mathbf{C}) + \beta\delta(\mathbf{B}-\mathbf{D}), \quad (24b)$$

$$\mathbf{G}_C = \alpha\gamma(\mathbf{C}-\mathbf{A}) + \beta\gamma(\mathbf{C}-\mathbf{B}) + \gamma\delta(\mathbf{C}-\mathbf{D}), \quad (24c)$$

$$\mathbf{G}_D = \alpha\delta(\mathbf{D}-\mathbf{A}) + \beta\delta(\mathbf{D}-\mathbf{B}) + \gamma\delta(\mathbf{D}-\mathbf{C}). \quad (24d)$$

The scalars and vectors above are independent of the choice of origin and $\hat{X}(k)$ and $X(x)$ are therefore likewise indepen-

dent. This contrasts with the annoying origin-dependence [35] of the one-electron posmom density $S(s)$.

Integrals of higher angular momentum can be generated by differentiating $[ssss]_{\hat{X}}$ with respect to the Cartesian coordinates of the basis function centres, as first suggested by Boys [39], or, more efficiently, using recurrence relations [31]. We have written a programme to compute $X(x)$ within an *spd* Gaussian basis set and implemented this in a development version of the Q-CHEM 3.2 quantum chemistry package [40].

Equations (21)–(24) are easily modified to generate $P(u)$, $M(v)$, $\Upsilon(\omega)$ and $\hat{D}(k)$. In particular, if the functions $\sinh(k)$, $\exp(k)$ and $\cosh(k)$ are replaced by their first-order approximations (k , $1+k$ and 1 , respectively) in the expressions for J and \mathbf{H} , one obtains the f -Dot integrals $[ssss]_{\hat{D}}$ [30].

In the special case of concentric s -type Gaussians, the intracule integrals become

$$[ssss]_P = \frac{4\pi^{5/2}}{\xi^{3/2}} u^2 \exp\left(-\frac{\mu}{\xi} u^2\right), \quad (25a)$$

$$[ssss]_M = \frac{4\pi^{5/2}}{\chi^{3/2}} v^2 \exp\left(-\frac{\nu}{\chi} v^2\right), \quad (25b)$$

$$[ssss]_{\Upsilon} = \frac{\pi^3 (\lambda - 2\eta \cos^2 \omega)}{2\zeta^{3/2} (\lambda + \eta \cos^2 \omega)^{5/2}} \sin \omega, \quad (25c)$$

$$[ssss]_{\hat{D}} = \frac{\pi^3}{K^{3/2}}, \quad (25d)$$

$$[ssss]_{\hat{X}} = \frac{\pi^3}{J^{3/2}}, \quad (25e)$$

where

$$\mu = (\alpha + \beta)(\gamma + \delta), \quad (26a)$$

$$\nu = (\alpha + \gamma)(\beta + \delta), \quad (26b)$$

$$\zeta = (\alpha + \delta)(\beta + \gamma), \quad (26c)$$




Figure 1. Physical interpretation of the variables u , v , ω and x in the weak, medium and strong correlation regimes.

$$\chi = 4(\alpha\beta\gamma + \alpha\beta\delta + \alpha\gamma\delta + \beta\gamma\delta), \quad (26d)$$

$$\lambda = \left(\frac{1}{\alpha + \delta} + \frac{1}{\beta + \gamma} \right) \left(\frac{\alpha\delta}{\alpha + \delta} + \frac{\beta\gamma}{\beta + \gamma} \right), \quad (26e)$$

$$\eta = \left(\frac{\alpha}{\alpha + \delta} - \frac{\beta}{\beta + \gamma} \right)^2, \quad (26f)$$

$$K = \xi[\beta k^2 + \gamma(1+k)^2 + \delta] - [\beta k + \gamma(1+k) + \delta]^2. \quad (26g)$$

In the calculations described below, we have computed $X(x)$ and $D(x)$ numerically using Equations (10) and (14). $\hat{D}(k)$, $\hat{X}(k)$, $D(x)$ and $X(x)$ are all even functions and we will therefore focus only on $x \geq 0$ and $k \geq 0$.

Physical interpretations of the variables u , v , ω and x are summarised in Figure 1. The three limiting configurations $\omega = 0, \pi/2$ and π (which correspond to $x = uv, 0$ and $-uv$) are depicted for the weak (u and v large), medium (where one of u and v is large and the other is small) and strong correlation (u and v small) regimes. A faithful description of electron correlation requires information about the relative position \mathbf{u} and momentum \mathbf{v} , but also on the mutual orientation ω of these two vectors, which gives insight into the nature of the electrons' mutual orbit. The Dot and Posmom intracules provide information about the distribution of values of $x = uv \cos \omega$, and thus about the type of correlation regime (weak, medium or strong). However, as noted

above, being a first-order approximation of $X(x)$, the information gathered in $D(x)$ is slightly biased. The effects of this approximation will be investigated below.

In Section 3, the Posmom intracule is investigated alongside $D(x)$, $\Upsilon(\omega)$, $P(u)$ and $M(v)$ for the two electrons in a parabolic quantum dot. In Section 4, we turn our attention to the electrons in a helium atom or helium-like ion. We also compare the Posmom intracules for ground and excited states and study the effect of the dimensionality of the space \mathcal{D} . Atomic units are used throughout.

3. Parabolic quantum dots

In our study of the Posmom intracule in parabolic quantum dots [41], we consider three different treatments of the Coulomb interaction between the two electrons. First, the non-interacting case, in which it is simply ignored; second, the Hartree–Fock (HF) case [42] in which it is approximated in a mean-field sense; third, the exact treatment which is possible for certain values of the harmonic confinement force constant [43,44].

3.1. Hamiltonian and wave functions

The Hamiltonian is

$$\mathcal{H} = -\frac{1}{2} (\nabla_1^2 + \nabla_2^2) + V(r_1) + V(r_2) + \frac{1}{r_{12}}, \quad (27)$$

where

$$V(r) = \frac{r^2}{2\kappa^2} \quad (28)$$

is the external harmonic potential and $1/\kappa^2$ is the force constant.

The 1S ground state of the non-interacting system has the wave function

$$\Psi_0(\mathbf{r}_1, \mathbf{r}_2) = \psi_0(\mathbf{r}_1)\psi_0(\mathbf{r}_2), \quad (29)$$

$$\psi_0(\mathbf{r}) = (\pi\kappa)^{-3/4} \exp\left(-\frac{r^2}{2\kappa}\right), \quad (30)$$

and the energy

$$E_0 = \frac{3}{\kappa}. \quad (31)$$

The more accurate HF wave function

$$\Psi_{\text{HF}}(\mathbf{r}_1, \mathbf{r}_2) = \psi_{\text{HF}}(\mathbf{r}_1)\psi_{\text{HF}}(\mathbf{r}_2) \quad (32)$$

is not known in closed form, but can be efficiently treated numerically by expanding $\psi_{\text{HF}}(\mathbf{r})$ in a Gaussian basis

$$\psi_{\text{HF}}(\mathbf{r}) = \sum_{j=1}^{N_G} c_j \exp(-\alpha_j r^2). \quad (33)$$

The HF energy can be directly minimised with respect to the coefficients c_j and exponents α_j using a numerical solver [45], thus avoiding the self-consistent field procedure usually needed for this kind of calculation [46,47].

The exact wave function and energy can be found in closed form [44] for certain values of κ . For example, for $\kappa = 2$

$$\Psi_2(\mathbf{r}_1, \mathbf{r}_2) = \left(1 + \frac{r_{12}}{2}\right) \psi_0(\mathbf{r}_1)\psi_0(\mathbf{r}_2), \quad (34a)$$

$$E_2 = 2, \quad (34b)$$

and, for $\kappa = 10$,

$$\Psi_{10}(\mathbf{r}_1, \mathbf{r}_2) = \left(1 + \frac{r_{12}}{2} + \frac{r_{12}^2}{20}\right) \psi_0(\mathbf{r}_1)\psi_0(\mathbf{r}_2), \quad (35a)$$

$$E_{10} = 1/2. \quad (35b)$$

Table 2 shows the convergence of E_{HF} with N_G for $\kappa = 2$ and $\kappa = 10$. The correlation energy

$$E_c = E_{\text{exact}} - E_{\text{HF}} \quad (36)$$

for $\kappa = 2$ ($E_c = 38.438\,871\,755\, mE_h$) agrees with earlier work [14,46]. For $\kappa = 10$, we find $E_c = 29.041\,525\,56\, mE_h$.

Table 2. Energies of parabolic quantum dots for different treatments of the interelectronic interaction.

	N_G	$\kappa = 2$	$\kappa = 10$
E_0	—	1.5	0.3
E_{HF}	1	2.04	0.53
	2	2.038 439	0.529 04
	3	2.038 438 9	0.529 041 5
	4	2.038 438 871 8	0.529 041 525 6
	5	2.038 438 871 755	0.529 041 525 56
	O'Neill and Gill ^a	2.038 438 87	—
	Ragot ^b	2.038 438 871 76	—
E_{exact}	—	2.	0.5

^aReference [14]: 7 basis functions.

^bReference [46]: 11 basis functions.

3.2. Position Intracule

The non-interacting Position intracule is

$$P_0(u) = \sqrt{\frac{2}{\pi\kappa^3}} u^2 \exp\left(-\frac{u^2}{2\kappa}\right), \quad (37)$$

and the HF intracule $P_{\text{HF},\kappa}(u)$ is found from (25a) and (33). For $\kappa = 2$ and $\kappa = 10$, the exact intracules are given by

$$P_2(u) = \frac{(1 + u/2)^2}{8 + 5\sqrt{\pi}} u^2 \exp\left(-\frac{u^2}{4}\right), \quad (38)$$

$$P_{10}(u) = \frac{(1 + u/2 + u^2/20)^2}{5/2(240 + 61\sqrt{5\pi})} u^2 \exp\left(-\frac{u^2}{20}\right). \quad (39)$$

Equation (38) has been reported previously [14].

3.3. Momentum intracule

Using the same notation as above, the non-interacting Momentum intracule is

$$M_0(v) = \frac{2}{\pi} \kappa^{3/2} v^2 \exp\left(-\frac{\kappa v^2}{2}\right), \quad (40)$$

$M_{\text{HF},2}(v)$ and $M_{\text{HF},10}(v)$ are obtained from (25b) and (33), and the exact Momentum intracules are

$$M_2(v) = \frac{8v^2}{8 + 5\sqrt{\pi}} \left[\sqrt{\frac{2}{\pi}} + e^{-\frac{v^2}{2}} + \left(\frac{1}{iv} + iv\right) \times \operatorname{erf}\left(\frac{iv}{\sqrt{2}}\right) e^{-\frac{v^2}{2}} \right]^2, \quad (41)$$

$$M_{10}(v) = \frac{80\sqrt{5}v^2}{48\sqrt{5} + 61\sqrt{\pi}} \left[\sqrt{\frac{10}{\pi}} + (4 - 5v^2) e^{-\frac{5v^2}{2}} + \left(\frac{1}{iv} + 5iv\right) \operatorname{erf}\left(\sqrt{\frac{5}{2}}iv\right) e^{-\frac{5v^2}{2}} \right]^2, \quad (42)$$

where $\text{erf}(z)$ is the error function [48]. Equation (41) has been reported previously [14].

3.4. Angle intracule

The Angle intracule of two non-interacting particles is entirely determined by the Jacobian factor and is [28]

$$\Upsilon_0(\omega) = \frac{1}{2} \sin \omega. \tag{43}$$

$\Upsilon_{\text{HF},2}(\omega)$ and $\Upsilon_{\text{HF},10}(\omega)$ are obtained from (25c) and (33). $\Upsilon_2(\omega)$ and $\Upsilon_{10}(\omega)$ have been obtained by numerical integration of (4) and (5). Equation (4) can be reduced to a two-dimensional integral, and the resulting four-dimensional numerical integration in Equation (5) was performed carefully to ensure accuracy of $O(10^{-3})$ for each value of ω .

3.5. Dot and Posmom intracules

$\hat{D}_{\text{HF},2}(k)$ and $\hat{D}_{\text{HF},10}(k)$ [Equations (33) and (25d)], as well as $\hat{X}_{\text{HF},2}(k)$ and $\hat{X}_{\text{HF},10}(k)$ [Equations (33) and (25e)] have been obtained numerically. Table 3 gathers the non-interacting and exact ($\kappa = 2$ and 10) Dot and Posmom intracules in Fourier and real space. The similarity between the Dot and Posmom expressions is striking.

3.6. Holes

The correlation hole was originally defined [2] as the difference between the exact and HF Position intracule

$$\Delta P(u) = P(u) - P_{\text{HF}}(u), \tag{44}$$

but this can be extended to any intracule I

$$\Delta I = I - I_{\text{HF}}. \tag{45}$$

One can also define the HF hole as the difference between the HF and non-interacting intracules

$$\Delta I_{\text{HF}} = I_{\text{HF}} - I_0. \tag{46}$$

Figure 2 shows all of the intracules for $\kappa = 2$ and Figure 3 shows the holes created as the Coulomb interaction is introduced.

One can see from $P(u)$ and $M(v)$ in Figure 2(a) or 2(b) that the electrons are found at larger separations and move with lower relative momenta in the HF approximation than in the non-interacting case. However, the non-interacting and HF intracules, $\Upsilon(\omega)$, $D(x)$ and $X(x)$, are almost identical. The fact that $\Upsilon(\omega)$, $D(x)$ and $X(x)$ are all invariant under a uniform scaling leads us to conclude that the introduction of the Coulomb operator at the mean-field level leads to an almost exact dilation of the system.

Table 3. Dot and Posmom intracules for the non-interacting and the exact treatments in Fourier and real spaces for two parabolic quantum dots. Note that Γ is the gamma function [48] and $\tilde{K}_n = x^n K_n$, where K_n represents the n th modified Bessel function of the second kind [48].

		Dot	Posmom
Fourier space	Non-int.	$\frac{1}{(1+k^2)^{3/2}}$	$\frac{1}{\cosh^{3/2}(2k)}$
	$\kappa = 2$	$\frac{1}{8+5\sqrt{\pi}} \left[-\frac{\sqrt{\pi}}{(1+k^2)^{3/2}} + \frac{8}{(1+k^2)^2} + \frac{6\sqrt{\pi}}{(1+k^2)^{5/2}} \right]$	$\frac{1}{8+5\sqrt{\pi}} \left[\frac{2\sqrt{\pi}}{\cosh^{3/2}(2k)} + \frac{8 \cosh(k)}{\cosh^2(2k)} + \frac{3\sqrt{\pi}}{\cosh^{5/2}(2k)} \right]$
	$\kappa = 10$	$\frac{1}{240+61\sqrt{5\pi}} \left[\frac{\sqrt{5\pi}}{(1+k^2)^{3/2}} - \frac{80}{(1+k^2)^2} + \frac{320}{(1+k^2)^3} + \frac{60\sqrt{5\pi}}{(1+k^2)^{7/2}} \right]$	$\frac{1}{240+61\sqrt{5\pi}} \left[\frac{16\sqrt{5\pi}}{\cosh^{3/2}(2k)} + \frac{80 \cosh(k)}{\cosh^2(2k)} + \frac{30\sqrt{5\pi}}{\cosh^{5/2}(2k)} + \frac{160 \cosh(k)}{\cosh^3(2k)} + \frac{15\sqrt{5\pi}}{\cosh^{7/2}(2k)} \right]$
Real space	Non-int.	$\frac{1}{\pi} x K_1(x)$	$\frac{1}{\sqrt{2\pi^3}} \left \Gamma\left(\frac{3+i\lambda}{4}\right) \right ^2$
	$\kappa = 2$	$\frac{1}{8\sqrt{\pi+5\pi}} [-\tilde{K}_1(x) + 2\sqrt{2}\tilde{K}_{3/2}(x) + 2\tilde{K}_2(x)]$	$\frac{\sqrt{2}}{(8+5\sqrt{\pi})\pi} \left[\left \Gamma\left(\frac{3+i\lambda}{4}\right) \right ^2 + 2 \left \Gamma\left(\frac{5+i\lambda}{4}\right) \right ^2 + \pi \frac{x \sinh(\frac{x\pi}{4}) + \cosh(\frac{x\pi}{4})}{\cosh(\frac{x\pi}{4})} \right]$
	$\kappa = 10$	$\frac{1}{240\sqrt{\pi+61\sqrt{5\pi}}} \left\{ 20\sqrt{2} [-\tilde{K}_{3/2}(x) + \tilde{K}_{5/2}(x)] + \sqrt{5} [\tilde{K}_1(x) + 4\tilde{K}_3(x)] \right\}$	$\frac{\sqrt{2}}{\sqrt{2\pi} (240+61\sqrt{5\pi})} \left[(x^2 + 25) \left \Gamma\left(\frac{3+i\lambda}{4}\right) \right ^2 + 40 \left \Gamma\left(\frac{5+i\lambda}{4}\right) \right ^2 + \frac{10x}{\sqrt{5}} \frac{\cosh(\frac{x\pi}{4}) + 4x \sinh(\frac{x\pi}{4})}{\cosh(\frac{x\pi}{4})} \right]$




Figure 2. Intracules for a parabolic quantum dot with $\kappa = 2$: non-interacting (—), HF (---) and exact (⋯). (a) Position. (b) Momentum. (c) Angle. (d) Dot. (e) Posmom.

Figure 3 reveals that the HF holes and correlation holes of $P(u)$ and $M(v)$ are surprisingly similar in size and shape. It also shows that the introduction of correlation decreases the probabilities of $\omega \approx \pi/2$ and $x \approx 0$, indicat-

ing that the correlated electrons spend less time circularly orbiting their centre of mass. This conclusion is supported by both the non-rigorous $\Upsilon(\omega)$ and $D(x)$ intracules and the rigorous $X(x)$ intracule.




Figure 3. Intracule holes for the parabolic quantum dots with $\kappa = 2$ and $\kappa = 10$: HF hole (—) and correlation hole (- - -). (a) Position $\kappa = 2$. (b) Position $\kappa = 10$. (c) Momentum $\kappa = 2$. (d) Momentum $\kappa = 10$. (e) Angle $\kappa = 2$. (f) Angle $\kappa = 10$. (g) Dot $\kappa = 2$. (h) Dot $\kappa = 10$. (i) Posmom $\kappa = 2$. (j) Posmom $\kappa = 10$.

Table 4. HF, radial and exact energies of various helium-like ions.

Atom	H ⁻	He	Li ⁺	B ³⁺	Ne ⁸⁺
Z	1	2	3	5	10
HF	-0.487 93	-2.861 67	-7.236 41	-21.986 2	-93.861 1
Radial	-0.514 5	-2.879 0	-7.252 5	-22.001 5	-93.875 9
Exact	-0.527 75	-2.903 72	-7.279 91	-22.031 0	-93.906 8
E_c	-0.03982	-0.04205	-0.04350	-0.0448	-0.0457
% E_c^{rad}	66.7	41.3	37.0	34.2	32.3

However, the differences between $D(x)$ and $X(x)$ are significant. $D(x)$ comes from the Wigner distribution and, as Equations (13) and (16) show, it is the $O(\hbar)$ approximation to $X(x)$. Its Fourier transform $\hat{D}(k)$ decays as k^{-3} for large k and this creates a discontinuity in the second derivative $D''(x)$ at $x = 0$ [30,49]. In contrast, $X(x)$ is smooth at $x = 0$. One of the consequences of this misbehaviour at $x = 0$ is that, for the $\kappa = 2$ quantum dot, the Dot intracule's prediction $D(0) = 0.219$ overestimates the exact value $X(0) = 0.168$ by 30%.

4. Helium-like ions

We now turn our attention to the helium-like ions. The Hamiltonian is obtained by substituting the harmonic potential $v(r)$ in (27) with the Coulombic potential

$$V(r) = -\frac{Z}{r}, \quad (47)$$

where Z is the nuclear charge. As before, the HF wave function and energy can be found by expanding the HF orbital in a Gaussian basis, optimising both the coefficients and exponents. We consider five values of Z , corresponding to the H⁻, He, Li⁺, B³⁺ and Ne⁸⁺ ions and, in this Section, we focus on their Position and Posmom intracules.

4.1. Ground state

The HF orbital of the ¹S ground state was approximated by a Gaussian expansion (33) with $N_G = 11$. The exact wave

function was approximated by the 64-term Hylleraas-type expansion [50]

$$\Psi(\mathbf{r}_1, \mathbf{r}_2) = \sum_{nlm}^3 c_{nlm} (r_1 + r_2)^n (r_1 - r_2)^{2l} r_{12}^m e^{-\alpha(r_1+r_2)}. \quad (48)$$

We also considered the 64-term radially-correlated wave function [51,52]

$$\Psi_{\text{rad}}(\mathbf{r}_1, \mathbf{r}_2) = \sum_{nl}^7 c_{nl} (r_1 + r_2)^n (r_1 - r_2)^{2l} e^{-\alpha(r_1+r_2)}. \quad (49)$$

Table 4 gathers the HF, exact and radially correlated energies obtained from (33), (48) and (49), respectively. Only the correct figures are reported [52–54], as well as the percentage of radial correlation (% E_c^{rad}). As above, the correlation hole is defined as the difference between exact and HF intracules [Equation (45)]. We also define the radial and angular holes [55]

$$\Delta I_{\text{rad}} = I_{\text{rad}} - I_{\text{HF}}, \quad (50)$$

$$\Delta I_{\text{ang}} = I - I_{\text{rad}}. \quad (51)$$

Both $P(u)$ and $X(x)$ were obtained numerically and are shown in Figure 4. Increasing the nuclear charge barely affects $X_{\text{HF}}(x)$ because it produces an almost uniform contraction of the system. However, the effect on the exact $X(x)$




Figure 4. Posmom intracules of helium-like ions: $Z = 1$ (- · · -), $Z = 2$ (- · -), $Z = 3$ (···), $Z = 5$ (- - -) and $Z = 10$ (—). (a) Hartree-Fock. (b) Exact.




Figure 5. Correlation holes of helium-like ions: $Z = 1$ (- · · -), $Z = 2$ (- · -), $Z = 3$ (···), $Z = 5$ (- - -) and $Z = 10$ (—). (a) Posmom. (b) Position.

is much larger and the values of $X(0)$ are 0.1642, 0.1917, 0.1968, 0.2003 and 0.2027 for $Z = 1, 2, 3, 5$ and 10 , respectively. These values reveal that the electrons in H^- spend less time mutually orbiting than those in Ne^{8+} . This is consistent with the conventional view that H^- is a more strongly correlated system than Ne^{8+} .

Figure 5 shows the Posmom and Position holes of the ions. The depth of $\Delta X(x)$ decreases as Z increases but the depth of $\Delta P(u)$ is almost constant as it is squeezed toward the origin. $\Delta P(u)$ exhibits a secondary hole, discussed in detail by Pearson *et al.* [18], but this subtle correlation effect is not visible in $\Delta X(x)$.

Radial correlation provides the majority (67%) of the total correlation energy in H^- but this decreases to 41% in He, and to 32% in Ne^{8+} , as angular correlation effects becomes dominant. This shift is visible in $\Delta P_{rad}(u)$ and $\Delta P_{ang}(u)$, as shown in Figure 6, but $\Delta X_{rad}(x)$ is always larger than $\Delta X_{ang}(x)$ and becomes almost identical for Ne^{8+} .

If we compare the insets in Figure 5(b) and Figure 6(c), we see that the *radial* secondary hole is several times deeper than the *total* secondary hole. This implies that the radial secondary hole, which has been noted previously by Katriel *et al.* [55], is almost entirely cancelled by an angular secondary hole.




Figure 6. Radial and angular holes in He-like ions: $Z = 1$ (- · · -), $Z = 2$ (- · -), $Z = 3$ (···), $Z = 5$ (- - -) and $Z = 10$ (—). (a) Posmom radial correlation hole. (b) Posmom angular correlation hole. (c) Position radial correlation hole. (d) Position angular correlation hole.




Figure 7. Posmom intracule for states of the He atom: $1s^2\ ^1S$ (—), $1s2s\ ^1S$ (---), $1s2s\ ^3S$ (···), $2s^2\ ^1S$ (- · -), $2p^2\ ^1S$ (- · · -).

4.2. Excited states

We have calculated the HF Posmom intracule for several excited states of the He atom using a Gaussian basis of 36 s -type functions with exponents $2^{-15}, 2^{-14}, \dots, 2^{20}$ and 31 p -type functions with exponents $2^{-10}, 2^{-14}, \dots, 2^{20}$. The maximum overlap method (MOM) has been employed for finding excited-state solutions to the HF self-consistent field equations [56].

The intracules in the ground state ($1s^2\ ^1S$) and four excited states ($1s2s\ ^1S$, $1s2s\ ^3S$, $2s^2\ ^1S$ and $2p^2\ ^1S$) are shown in Figure 7. Table 5 lists the values of $X(0)$ for these and other excited states.

When the electron pair occupies a more diffuse orbital, $X(x)$ becomes broader and $X(0)$ drops from 0.206 in the $1s^2$ state to 0.165 in the $2s^2$ state, and to 0.127 for the $2p^2$ state. The decrease is even more marked when the electrons occupy orbitals in different shells, such as in the $1s2s\ ^1S$ state where $X(0) = 0.087$. However, if the two orbitals have the same principal quantum number, such as in the $2s2p\ ^1P$ state, the decrease is smaller.

The Dot intracule has been calculated for the first excited state $1s2s\ ^3S$ [30] and shows a small dip in $D(x)$ around $x = 0$, which we have previously attributed to the Fermi hole. We now believe that that explanation was incorrect and that the dip is a failure of $D(x)$ to capture the behaviour of $X(x)$.

Table 5. Origin Posmom intracule, $X(0)$, for various singlet and triplet excited states of the helium atom ($i, j = x, y, z$).

Configuration	Singlet	Triplet
$1s^2$	0.2060	—
$1s2s$	0.0866	0.09520
$1s2p$	0.0928	0.09525
$2s^2$	0.1647	—
$2s2p$	0.1455	0.1153
$2p_i^2$	0.1267	—
$2p_i2p_j$	0.1567	0.1538

4.3. D -dimensional helium atom

Following the pioneering work of Loeser and Herschbach on the effect of dimensionality on the HF [57,58] and exact energies [59] of He-like ions, several other two-electron systems have recently been studied in D dimensions [47,60–64].

The generalisation of the Posmom intracule for a D -dimensional space is straightforward. Equations (10), (11) and (13) are unchanged and Equation (25e) becomes

$$[s s s s]_{\hat{X}} = \frac{\pi^D}{J^{D/2}}. \quad (52)$$

We used a large, even-tempered Gaussian basis, optimising the coefficients c_j to minimise the HF energy [58,62]

$$E_{\text{HF}} = 2 \int \psi_{\text{HF}}(r) \left[-\frac{\nabla^2}{2} + V(r) \right] \psi_{\text{HF}}(r) dr + \iint \psi_{\text{HF}}^2(r_1) \left\langle \frac{1}{r_{12}} \right\rangle_{\mathcal{D}} \psi_{\text{HF}}^2(r_2) dr_1 dr_2, \quad (53)$$

where

$$\left\langle \frac{1}{r_{12}} \right\rangle_{\mathcal{D}} = \frac{F\left(\frac{3-D}{2}, \frac{1}{2}, \frac{D}{2}, \frac{\min(r_1, r_2)^2}{\max(r_1, r_2)^2}\right)}{\max(r_1, r_2)}, \quad (54)$$

$$\nabla^2 = \frac{d^2}{dr^2} + \frac{D-1}{r} \frac{d}{dr}, \quad (55)$$

$$d\mathbf{r} = \frac{2\pi^{D/2}}{\Gamma(D/2)} r^{D-1} dr, \quad (56)$$

and F is the Gauss hypergeometric function [48]. Our energies for $D = 2, 3, 4$ and 5 agree within a microhartree with the benchmark values of Herschbach and co-workers [58].

Figure 8 shows how $X(x)$ changes with D . The observation that the intracule broadens as D increases is consistent with the conclusion of Herrick and Stillinger [65] that the electrons in D -helium can avoid each other more easily when D is large. Furthermore, they have shown that the binding energy of the ground state in $D = 5$ corresponds




Figure 8. HF Posmom intracules of the D -dimensional He atom: $D = 2$ (—), $D = 3$ (---), $D = 4$ (···), $D = 5$ (- · -).

exactly to the binding energy of the $2p_i2p_j^3P$ state in $\mathcal{D} = 3$. This feature is due to interdimensional degeneracies, first noticed by van Vleck [66], and observed for various systems [61,65,67–70]. We observe likewise that $X(x)$ for the $1s^2$ state in $\mathcal{D} = 5$ is identical to $X(x)$ for the $2p_i2p_j^3P$ state in $\mathcal{D} = 3$.

5 Conclusion

We have introduced a new two-particle density distribution, the Posmom intracule, which condenses information about both the relative position and relative momentum of the particles. We have shown how to construct this distribution from the many-particle wave function and we have shown that the Dot intracule $D(x)$ is a first-order approximation of the Posmom intracule $X(x)$. We have applied our new formalism to two-electron quantum dots and the helium-like ions. A comparison between various intracules (Position, Momentum, Angle, Dot and Posmom) has been carried out, showing the interrelated information conveyed by these two-particle probability distributions.

The Posmom intracule, unlike the Dot intracule, is the rigorous quantum mechanical probability density for $x = \mathbf{u} \cdot \mathbf{v}$. This variable yields information about the rate at which two electrons approach and therefore vanishes when, for example, the electrons are in a circular orbit. Such detailed knowledge about the relative motion of pairs of electrons is the necessary ingredient for IFT and we therefore anticipate that $X(x)$ will play an important role in the future development of that approach to the electronic structure problem.

Acknowledgements

We thank Joshua Hollett for helpful comments. YAB thanks the ANU Research School of Chemistry for a PhD scholarship. PMWG thanks the Australian Research Council (Grants DP0984806, DP1094170, and DP120104740) for funding. PFL thanks the Australian Research Council for a Discovery Early Career Researcher Award (Grant DE130101441). PFL and PMWG thank the NCI National Facility for a generous grant of supercomputer time.

References

- [1] E.R. Davidson, *Reduced Density Matrices in Quantum Chemistry* (Academic, New York, 1976).
- [2] C.A. Coulson and A.H. Neilson, Proc. Phys. Soc. (London) **78**, 831 (1961).
- [3] R.F. Curl, Jr, and C.A. Coulson, Proc. Phys. Soc. (London) **85**, 647 (1965).
- [4] R. Benesch, J. Phys. B **4**, 1403 (1971).
- [5] A.J. Thakkar and V.H. Smith, Jr, Chem. Phys. Lett. **42**, 476 (1976).
- [6] A.J. Thakkar and V.H. Smith, Jr, J. Chem. Phys. **67**, 1191 (1977).
- [7] J.P. Dahl and M. Springborg, J. Chem. Phys. **88**, 4535 (1988).
- [8] J.P. Perdew and Y. Wang, Phys. Rev. B **46**, 12947 (1992).
- [9] T. Koga, Y. Kasai, and A.J. Thakkar, Int. J. Quantum Chem. **46**, 689 (1993).

- [10] J. Cioslowski and G. Liu, J. Chem. Phys. **109**, 8225 (1998).
- [11] A.M. Lee and P.M.W. Gill, Chem. Phys. Lett. **313**, 271 (1999).
- [12] X. Fradera, C. Sarasola, J.M. Ugalde, and R.J. Boyd, Chem. Phys. Lett. **304**, 393 (1999).
- [13] P.M.W. Gill, A.M. Lee, N. Nair, and R.D. Adamson, J. Mol. Struct. Theochem. **506**, 303 (2000).
- [14] D.P. O’Neill and P.M.W. Gill, Phys. Rev. A **68**, 022505 (2003).
- [15] P.M.W. Gill, D.P. O’Neill, and N.A. Besley, Theor. Chem. Acc. **109**, 241 (2003).
- [16] P.M.W. Gill, D.L. Crittenden, D.P. O’Neill, and N.A. Besley, Phys. Chem. Chem. Phys. **8**, 15 (2006).
- [17] J.K. Pearson, D.L. Crittenden, and P.M.W. Gill, J. Chem. Phys. **130**, 164110 (2009).
- [18] J.K. Pearson, P.M.W. Gill, J.M. Ugalde, and R.J. Boyd, Mol. Phys. **107**, 1089 (2009).
- [19] P-F. Loos, Phys. Rev. A **81**, 032510 (2010).
- [20] P-F. Loos and P.M.W. Gill, Phys. Rev. A **81**, 052510 (2010).
- [21] A.J. Proud, M.P. Walker, and J.K. Pearson, Int. J. Quantum Chem. **113**, 76 (2013).
- [22] K.E. Banyard and C.E. Reed, J. Phys. B **11**, 2957 (1978).
- [23] N.A. Besley, A.M. Lee, and P.M.W. Gill, Mol. Phys. **100**, 1763 (2002).
- [24] E. Wigner, Phys. Rev. **40**, 749 (1932).
- [25] M. Hillery, R.F. O’Connell, M.O. Scully, and E.P. Wigner, Phys. Rep. **106**, 121 (1984).
- [26] V.A. Rassolov, J. Chem. Phys. **110**, 3672 (1999).
- [27] D.L. Crittenden and P.M.W. Gill, J. Chem. Phys. **127**, 014101 (2007).
- [28] E.E. Dumont, D.L. Crittenden, and P.M.W. Gill, Phys. Chem. Chem. Phys. **9**, 5340 (2007).
- [29] D.L. Crittenden, E.E. Dumont, and P.M.W. Gill, J. Chem. Phys. **127**, 141103 (2007).
- [30] Y.A. Bernard, D.L. Crittenden, and P.M.W. Gill, Phys. Chem. Chem. Phys. **10**, 3447 (2008).
- [31] J.W. Hollett and P.M.W. Gill, Phys. Chem. Chem. Phys. **13**, 2972 (2011).
- [32] P.M.W. Gill, Annu. Rep. Prog. Chem., Sect. C **107**, 229 (2011).
- [33] D.L. Crittenden and P.M.W. Gill, *Solving the Schrödinger Equation: Has Everything Been Tried?* (IC Press, London, 2011), Chap. Intracule functional theory.
- [34] Y.A. Bernard and P.M.W. Gill, New J. Phys. **11**, 083015 (2009).
- [35] Y.A. Bernard and P.M.W. Gill, J. Phys. Chem. Lett. **1**, 1254 (2010).
- [36] Y.A. Bernard, D.L. Crittenden, and P.M.W. Gill, J. Phys. Chem. A **114**, 11984 (2010).
- [37] J. Wess, Il Nuovo Cimento, **18**, 1086 (1960).
- [38] W.O. Amrein, *Hilbert Space Methods in Quantum Mechanics* (EPFL Press, Laussane, Switzerland, 2009).
- [39] S.F. Boys, Proc. Roy. Soc. (London) A200, 542 (1950).
- [40] Y. Shao, L. Fusti-Molnar, Y. Jung, J. Kussmann, C. Ochsenfeld, S.T. Brown, A.T.B. Gilbert, L.V. Slipchenko, S.V. Levchenko, D.P. O’Neill, R.A. DiStasio, Jr, R.C. Lochan, T. Wang, G.J.O. Beran, N.A. Besley, J.M. Herbert, C.Y. Lin, T. van Voorhis, S.H. Chien, A. Sodt, R.P. Steele, V.A. Rassolov, P.E. Maslen, P.P. Korambath, R.D. Adamson, B. Austin, J. Baker, E.F.C. Byrd, H. Dachsel, R.J. Doerksen, A. Dreuw, B.D. Dunietz, A.D. Dutoi, T.R. Furlani, S.R. Gwaltney, A. Heyden, S. Hirata, C.P. Hsu, G. Kedziora, R.Z. Khaliullin, P. Klunzinger, A.M. Lee, M.S. Lee, W.Z. Liang, I. Lotan, N. Nair, B. Peters, E.I. Proynov, P.A. Pieniazek, Y.M. Rhee, J. Ritchie, E. Rosta, C.D. Sherrill, A.C. Simmonett, J.E. Subotnik, H.L. Woodcock, III, W. Zhang, A.T. Bell, A.K.

- Chakraborty, D.M. Chipman, F.J. Keil, A. Warshel, W.J. Hehre, H.F. Schaefer, III, J. Kong, A.I. Krylov, P.M.W. Gill, and M. Head-Gordon, *Phys. Chem. Chem. Phys.* **8**, 3172 (2006).
- [41] N.R. Kestner and O. Sinanoglu, *Phys. Rev.* **128**, 2687 (1962).
- [42] A. Szabo and N.S. Ostlund, *Modern Quantum Chemistry : Introduction to Advanced Structure Theory* (Dover publications Inc., Mineola, New York, 1989).
- [43] S. Kais, D.R. Herschbach, and R.D. Levine, *J. Chem. Phys.* **91**, 7791 (1989).
- [44] M. Taut, *Phys. Rev. A* **48**, 3561 (1993).
- [45] Wolfram Research, Inc., *Mathematica, version 8.0* (2011).
- [46] S. Ragot, *J. Chem. Phys.* **128**, 164104 (2008).
- [47] P.-F. Loos and P.M.W. Gill, *J. Chem. Phys.* **132**, 234111 (2010).
- [48] F.W.J. Olver, D.W. Lozier, R.F. Boisvert, and C.W. Clark, editors, *NIST Handbook of Mathematical Functions* (Cambridge University Press, New York, 2010).
- [49] J.P. Boyd, *Chebyshev and Fourier Spectral Methods*, 2nd ed. (Dover, New York, 2000).
- [50] E.A. Hylleraas, *Z. Phys.* **54**, 347 (1929).
- [51] E. Clementi, W. Kraemer, and C. Salez, *J. Chem. Phys.* **53**, 125 (1970).
- [52] T. Koga, *Z. Phys. D* **37**, 301 (1996).
- [53] T. Koga, M. Omura, H. Teruya, and A.J. Thakkar, *J. Phys. B* **28**, 3113 (1995).
- [54] H. Nakashima and H. Nakatsuji, *J. Chem. Phys.* **127**, 224104 (2007).
- [55] J. Katriel, S. Roy, and M. Springborg, *J. Chem. Phys.* **123**, 104104 (2005).
- [56] A.T.B. Gilbert, N.A. Besley, and P.M.W. Gill, *J. Phys. Chem. A* **112**, 13164 (2008).
- [57] D.R. Herschbach, *J. Chem. Phys.* **84**, 838 (1986).
- [58] J.G. Loeser and D.R. Herschbach, *J. Chem. Phys.* **84**, 3893 (1986).
- [59] J.G. Loeser and D.R. Herschbach, *J. Chem. Phys.* **84**, 3882 (1986).
- [60] P.-F. Loos and P.M.W. Gill, *Phys. Rev. Lett.* **103**, 123008 (2009).
- [61] P.-F. Loos and P.M.W. Gill, *Mol. Phys.* **108**, 2527 (2010).
- [62] P.-F. Loos and P.M.W. Gill, *J. Chem. Phys.* **131**, 241101 (2009).
- [63] P.-F. Loos and P.M.W. Gill, *Phys. Rev. Lett.* **105**, 113001 (2010).
- [64] P.-F. Loos and P.M.W. Gill, *Chem. Phys. Lett.* **500**, 1 (2010).
- [65] D.R. Herrick and F.H. Stillinger, *Phys. Rev. A* **11**, 42 (1975).
- [66] J.H. van Vleck, *Central Fields in Two vis-à-vis Three Dimensions: An Historical Divertissement* (London: Butterworths, 1973), p. 26
- [67] D.R. Herrick, *J. Math. Phys.* **16**, 281 (1975).
- [68] D.J. Doren and D.R. Herschbach, *J. Chem. Phys.* **85**, 4557 (1986).
- [69] D.Z. Goodson, D.K. Watson, J.G. Loeser, and D.R. Herschbach, *Phys. Rev. A* **44**, 97 (1991).
- [70] M. Dunn and D.K. Watson, *Phys. Rev. A* **59**, 1109 (1999).