

# A Mountaineering Strategy to Excited States: Highly Accurate Energies and Benchmarks for Medium Sized Molecules

Pierre-François Loos,\* Filippo Lipparini,\* Martial Boggio-Pasqua, Anthony Scemama, and Denis Jacquemin\*

Cite This: *J. Chem. Theory Comput.* 2020, 16, 1711–1741

Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

**ABSTRACT:** Following our previous work focusing on compounds containing up to 3 non-hydrogen atoms [*J. Chem. Theory Comput.* 2018, 14, 4360–4379], we present here highly accurate vertical transition energies obtained for 27 molecules encompassing 4, 5, and 6 non-hydrogen atoms: acetone, acrolein, benzene, butadiene, cyanoacetylene, cyanoformaldehyde, cyanogen, cyclopentadiene, cyclopropanone, cyclopropanethione, diacetylene, furan, glyoxal, imidazole, isobutene, methylenecyclopropane, propynal, pyrazine, pyridazine, pyridine, pyrimidine, pyrrole, tetrazine, thioacetone, thiophene, thiopropynal, and triazine. To obtain these energies, we use equation-of-motion/linear-response coupled cluster theory up to the highest technically possible excitation order for these systems (CC3, EOM-CCSDT, and EOM-CCSDTQ) and selected configuration interaction (SCI) calculations (with tens of millions of determinants in the reference space), as well as the multiconfigurational  $n$ -electron valence state perturbation theory (NEVPT2) method. All these approaches are applied in combination with diffuse-containing atomic basis sets. For all transitions, we report at least CC3/*aug-cc-pVQZ* vertical excitation energies as well as CC3/*aug-cc-pVTZ* oscillator strengths for each dipole-allowed transition. We show that CC3 almost systematically delivers transition energies in agreement with higher-level methods with a typical deviation of  $\pm 0.04$  eV, except for transitions with a dominant double excitation character where the error is much larger. The present contribution gathers a large, diverse, and accurate set of more than 200 highly accurate transition energies for states of various natures (valence, Rydberg, singlet, triplet,  $n \rightarrow \pi^*$ ,  $\pi \rightarrow \pi^*$ , ...). We use this series of theoretical best estimates to benchmark a series of popular methods for excited state calculations: CIS(D), ADC(2), CC2, STEOM-CCSD, EOM-CCSD, CCSDR(3), CCSDT-3, CC3, and NEVPT2. The results of these benchmarks are compared to the available literature data.



## 1. INTRODUCTION

Accurately describing transition energies between the electronic ground state (GS) and excited states (ESs) remains an important challenge in quantum chemistry. When dealing with large compounds in complex environments, one is typically limited to the use of time-dependent density-functional theory (TD-DFT),<sup>1–3</sup> a successful yet far from flawless approach. In particular, to perform TD-DFT calculations, one must choose an “appropriate” exchange-correlation functional, which is difficult yet primordial as the impact of the exchange-correlation functional is exacerbated within TD-DFT as compared to DFT.<sup>4</sup> Such a selection can, of course, rely on the intrinsic features of the various exchange-correlation functional families; e.g., it is well-known that range-separated hybrids provide a more physically sound description of long-range charge-transfer transitions than semilocal exchange-correlation functionals in the adiabatic TD-DFT framework.<sup>5,6</sup> However, to obtain a quantitative assessment of the accuracy that can be expected from TD-DFT calculations, benchmarks cannot be avoided. This is why so many assessments of TD-DFT performance for various ES properties are available.<sup>4</sup>

While several of these benchmarks rely on experimental data as reference (typically band shapes<sup>7–14</sup> or 0-0 energies<sup>8,15–22</sup>), theoretical best estimates (TBEs) based on state-of-the-art computational methods<sup>23–28</sup> are advantageous as they allow comparisons on a perfectly equal footing (same geometry, vertical transitions, no environmental effects, etc.). In such a case, the challenge is in fact to obtain accurate TBEs, as top-notch theoretical models generally come with a dreadful scaling with system size and, in addition, typically require large atomic basis sets to deliver transition energies close to the complete basis set (CBS) limit.<sup>29</sup>

More than 20 years ago, Serrano-Andrés, Roos, and collaborators compiled an impressive series of reference transition energies for typical conjugated organic molecules (butadiene, furan, pyrrole, tetrazine, etc.).<sup>30–38</sup> To this end, they relied on experimental GS geometries and the complete-active-

Received: December 5, 2019

Published: January 27, 2020

space second-order perturbation theory (CASPT2) approach with the largest active spaces and basis sets one could dream of at the time. These CASPT2 values were later used to assess the performance of TD-DFT combined with various exchange-correlation functionals<sup>39,40</sup> and remained for a long time the best theoretical references available on the market. However, beyond comparisons with experiments, which are always challenging when computing vertical transition energies,<sup>14</sup> there was no approach available at that time to ascertain the accuracy of these transition energies. Nowadays, it is of common knowledge that CASPT2 has the tendency of underestimating vertical excitation energies in organic molecules when the IPEA shift is not included. It is also known that the use of a standard value of 0.25 au for this IPEA shift may lead to overestimations of the transition energies, making the use of this shift questionable.<sup>41</sup>

A decade ago, Thiel and co-workers defined TBEs for 104 singlet and 63 triplet valence ESs in 28 small and medium conjugated CNOH organic molecules.<sup>23,25,26</sup> These TBEs were computed on MP2/6-31G(d) structures with several levels of theory, notably CASPT2 and various coupled cluster (CC) variants (CC2, CCSD, and CC3). Interestingly, while the default theoretical protocol used by Thiel and co-workers to define their first series of TBEs was CASPT2,<sup>23</sup> the vast majority of their most recent TBEs (the so-called “TBE-2” in ref 26) were determined at the CC3 level of theory with the *aug-cc-pVTZ* (aVTZ) basis set, often using a basis set extrapolation technique. More specifically, CC3/TZVP values were corrected for basis set incompleteness errors by the difference between, e.g., the CC2/aVTZ and CC2/TZVP results.<sup>25,26</sup> Many works have exploited Thiel’s TBEs for assessing lower-order methods,<sup>24,42–68</sup> highlighting further their value for the electronic structure community. In contrast, the number of extensions/improvements of this original set remains quite limited. For example, Kánnár and Szalay computed, in 2014, CCSDT/TZVP reference energies for 17 singlet states of six molecules.<sup>69</sup> Three years later, the same authors reported 46 CCSDT/aVTZ transition energies in small compounds containing two or three non-hydrogen atoms (ethylene, acetylene, formaldehyde, formalimine, and formamide).<sup>70</sup>

Following the same philosophy, two years ago, we reported a set of 106 transition energies for which it was technically possible to reach the full configuration interaction (FCI) limit by performing high-order CC (up to CCSDTQP) and selected CI (SCI) calculations on CC3/aVTZ GS structures.<sup>28</sup> We exploited these TBEs to benchmark many ES methods.<sup>28</sup> Among our conclusions, we found that CCSDTQ yields near-FCI quality excitation energies, whereas we could not detect any significant differences between CC3 and CCSDT transition energies, both being very accurate with mean absolute errors (MAEs) as small as 0.03 eV compared to FCI.

Although these conclusions agree well with earlier studies,<sup>69–71</sup> they obviously only hold for single excitations, i.e., transitions with % $T_1$ , the single-excitation character in CC3 calculations, in the 80–100% range. Therefore, we also recently proposed a set of 20 TBEs for transitions exhibiting a significant double-excitation character (i.e., with % $T_1$  typically below 80%).<sup>72</sup> Unsurprisingly, our results clearly evidenced that the error in the CC methods is intimately related to the % $T_1$  value. For example, for the ES with a significant yet not dominant double excitation character [such as the infamous  $A_g$  ES of butadiene (% $T_1$  = 75%)] CC methods including triples deliver rather accurate estimates (MAE of 0.11 eV with CC3 and 0.06 eV with CCSDT), surprisingly outperforming second-order

multireference schemes such as CASPT2 or the generally robust *n*-electron valence state perturbation theory (NEVPT2). In contrast, for ESs with a dominant double excitation character, e.g., the low-lying (*n*, *n*) → ( $\pi^*$ ,  $\pi^*$ ) excitation in nitrosomethane (% $T_1$  = 2%), single-reference methods (not including quadruples) have been found to be unsuitable with MAEs of 0.86 and 0.42 eV for CC3 and CCSDT, respectively. In this case, multiconfigurational methods are in practice required to obtain accurate results.<sup>72</sup>

A clear limit of our 2018 work<sup>28</sup> was the size of the compounds put together in our set. These were limited to 1–3 non-hydrogen atoms, hence introducing a potential “chemical” bias. Therefore, we have decided, in the present contribution, to consider larger molecules with organic compounds encompassing 4, 5, and 6 non-hydrogen atoms. For such systems, performing CCSDTQ calculations with large one-electron basis sets is elusive. Moreover, the convergence of the SCI energy with respect to the number of determinants is obviously slower for these larger compounds; hence, extrapolating to the FCI limit with an error of ~0.01 eV is rarely achievable in practice. Consequently, the “brute-force” determination of FCI/CBS estimates, as in our earlier work,<sup>28</sup> is definitely out of reach here. Anticipating this problem, we have recently investigated bootstrap CBS extrapolation techniques.<sup>28,72</sup> In particular, we have demonstrated that, following an ONIOM-like scheme,<sup>73</sup> one can very accurately estimate such a limit by correcting high-level values obtained in a small basis by the difference between CC3 results obtained in a larger basis and those in the same small basis.<sup>28</sup> We globally follow such a strategy here. In addition, we also perform NEVPT2 calculations in an effort to check the consistency of our estimates. This is particularly critical for ES with intermediate % $T_1$  values. Using this protocol, we define a set of more than 200 *aug-cc-pVTZ* reference transition energies, most being within ±0.03 eV of the FCI limit. These reference energies are obtained on CC3/aVTZ geometries, and additional basis set corrections (up to quadruple- $\zeta$  at least) are also provided for CC3. Together with the results obtained in our two earlier works,<sup>28,72</sup> the present TBE will hopefully contribute to climbing a rung higher on the ES accuracy ladder.


## 2. COMPUTATIONAL DETAILS

Unless otherwise stated, all transition energies are computed in the frozen-core approximation (with a large core for the sulfur atoms). Pople’s 6-31+G(d) and Dunning’s *aug-cc-pVXZ* (*X* = D, T, Q, and 5) atomic basis sets are systematically employed in our excited-state calculations. In the following, we employ the aVXZ shorthand notations for these diffuse-containing basis sets. We note that an alternative family of more compact diffuse basis sets (such as *jun-cc-pVTZ*) have been proposed by Truhlar and co-workers.<sup>74</sup> Such variants could be better suited to reach CBS-quality transition energies at a smaller computational cost. As we intend to provide benchmark values here, we nevertheless stick to the original Dunning’s bases, which are directly available in almost any quantum chemistry codes. Various statistical quantities are reported in the remainder of this paper: the mean signed error (MSE), mean absolute error (MAE), root-mean-square error (RMSE), the standard deviation of the errors (SDE), and the maximum positive [Max(+)] and negative [Max(–)] errors. Here, we globally follow the same procedure as in ref 28, so that we only briefly outline the various theoretical methods that we have employed in the subsections below.


**2.1. Geometries.** The molecules considered herein are displayed in Scheme 1. Consistently with our previous work,<sup>28</sup>

Scheme 1. Representation of the Considered Compounds


## 4 (non-H) atoms



## 5 (non-H) atoms



## 6 (non-H) atoms



we systematically use CC3/aVTZ GS geometries obtained without applying the frozen-core approximation. The Cartesian coordinates (in bohr) of each compound can be found in the [Supporting Information](#) (SI). Several structures have been extracted from previous contributions,<sup>75–77</sup> whereas the missing structures were optimized using DALTON<sup>78</sup> and/or CFOUR,<sup>79</sup> applying default parameters in both cases.

**2.2. Selected Configuration Interaction.** Because SCI methods are less widespread than the other methods mentioned in the [Introduction](#), we shall detail further their main features. All the SCI calculations have been performed in the frozen-core approximation with the latest version of QUANTUM PACKAGE<sup>80</sup> using the Configuration Interaction using a Perturbative Selection made Iteratively (CIPSI) algorithm to select the most important determinants in the FCI space. Instead of generating all possible excited determinants like a conventional CI calculation, the iterative CIPSI algorithm performs a sparse exploration of the FCI space via a selection of the most relevant determinants using a second-order perturbative criterion. At each iteration, the variational (or reference) space is enlarged with new determinants. CIPSI can be seen as a deterministic version of the FCIQMC algorithm developed by Alavi and co-workers.<sup>81</sup> We refer the interested reader to ref 80, where our implementation of the CIPSI algorithm is detailed.

Excited-state calculations are performed within a state-averaged formalism which means that the CIPSI algorithm selects determinants simultaneously for the GS and ESs. Therefore, all electronic states share the same set of determinants with different CI coefficients. Our implementation of the CIPSI algorithm for ES is detailed in ref 82. For each system, a preliminary SCI calculation is performed using

Hartree–Fock orbitals in order to generate SCI wave functions with at least 5 000 000 determinants. State-averaged natural orbitals are then computed based on this wave function, and a new, larger SCI calculation is performed with this new set of orbitals. This has the advantage to produce a smoother and faster convergence of the SCI energy toward the FCI limit. For the largest systems, an additional iteration is sometimes required in order to obtain better quality natural orbitals and hence well-converged calculations.

The total SCI energy is defined as the sum of the (zeroth-order) variational energy (computed via diagonalization of the CI matrix in the reference space) and a second-order perturbative correction which takes into account the external determinants, i.e., the determinants which do not belong to the variational space but are linked to the reference space via a nonzero matrix element. The magnitude of this second-order correction,  $E^{(2)}$ , provides a qualitative idea of the “distance” to the FCI limit. For maximum efficiency, the total SCI energy is linearly extrapolated to  $E^{(2)} = 0$  (which effectively corresponds to the FCI limit) using the two largest SCI wave functions. These extrapolated total energies (simply labeled as FCI in the remainder of the paper) are then used to compute vertical excitation energies. Although it is not possible to provide a theoretically sound error bar, we estimate the extrapolation error by the difference in excitation energy between the largest SCI wave function and its corresponding extrapolated value. We believe that it provides a very safe estimate of the extrapolation error. Additional information about the SCI wave functions and excitation energies as well as their extrapolated values can be found in the [SI](#).

Table 1. Vertical Transition Energies (in eV) of Cyanoacetylene, Cyanogen, and Diacetylene<sup>a</sup>

| state   | Cyanoacetylene |       |        |             |      |       |        |             |      |       |        |      | lit. <sup>b</sup> | exp. <sup>c</sup> |        |       |                   |      |      |      |
|---|----------------|-------|--------|-------------|------|-------|--------|-------------|------|-------|--------|------|-------------------|-------------------|--------|-------|-------------------|------|------|------|
|   | 6-31+G(d)      |       |        |             |      |       | aVTZ   |             |      |       |        |      |                   |                   | aVQZ   | aVSZ  |                   |      |      |      |
|   | CC3            | CCSDT | CCSDTQ | FCI         | CC3  | CCSDT | CCSDTQ | FCI         | CC3  | CCSDT | CCSDTQ | FCI  |                   |                   | CC3    | CCSDT | NEVPT2            | CC3  | CC3  |      |
| <sup>1</sup> Σ <sub>g</sub> <sup>-</sup>                  | 6.02           | 6.04  | 6.02   | 6.02 ± 0.01 | 5.92 | 5.92  | 5.91   | 5.84 ± 0.09 | 5.80 | 5.81  | 5.78   | 5.78 | 5.79              | 5.79              | 5.79   | 5.79  | 5.79              | 5.46 | 4.77 |      |
| <sup>1</sup> Δ <sub>g</sub>                               | 6.29           | 6.31  | 6.29   | 6.28 ± 0.01 | 6.17 | 6.19  | 6.17   | 6.14 ± 0.05 | 6.08 | 6.09  | 6.10   | 6.10 | 6.06              | 6.06              | 6.06   | 6.06  | 6.06              | 5.81 | 5.48 |      |
| <sup>3</sup> Σ <sub>g</sub> <sup>+</sup>                  | 4.44           | 4.45  | 4.45   | 4.45 ± 0.03 | 4.43 | 4.43  | 4.43   | 4.41 ± 0.06 | 4.45 | 4.44  | 4.45   | 4.45 | 4.46              | 4.46              | 4.46   | 4.47  | 4.47              |      |      |      |
| <sup>3</sup> Δ <sub>g</sub>                               | 5.35           | 5.34  | 5.34   | 5.32 ± 0.03 | 5.28 | 5.27  | 5.27   | 5.20 ± 0.08 | 5.22 | 5.21  | 5.19   | 5.19 | 5.22              | 5.22              | 5.22   | 5.22  | 5.22              |      |      |      |
| <sup>1</sup> A <sup>+</sup> [F] <sup>d</sup>              | 3.70           | 3.72  | 3.70   | 3.67 ± 0.03 | 3.60 | 3.62  | 3.60   | 3.59 ± 0.02 | 3.54 | 3.56  | 3.50   | 3.50 | 3.54              | 3.54              | 3.54   | 3.54  |                   |      |      |      |
|   | Cyanogen       |       |        |             |      |       |        |             |      |       |        |      | aVTZ              | aVQZ              | aVSZ   | lit.  | exp. <sup>e</sup> |      |      |      |
|   | 6-31+G(d)      |       |        |             |      |       | aVDZ   |             |      |       |        |      | CC3               | CCSDT             | NEVPT2 | CC3   | CC3               | CC3  |      |      |
| <sup>1</sup> Σ <sub>g</sub> <sup>-</sup>                  | 6.62           | 6.63  | 6.62   | 6.58 ± 0.03 | 6.52 | 6.52  | 6.51   | 6.44 ± 0.08 | 6.39 | 6.40  | 6.40   | 6.40 | 6.38              | 6.38              | 6.32   | 6.38  | 6.38              | 6.38 | 5.63 | 5.99 |
| <sup>1</sup> Δ <sub>g</sub>                               | 6.88           | 6.89  | 6.88   | 6.87 ± 0.02 | 6.77 | 6.78  | 6.77   | 6.74 ± 0.04 | 6.66 | 6.67  | 6.67   | 6.66 | 6.64              | 6.64              | 6.66   | 6.64  | 6.64              | 6.64 | 5.99 | 4.13 |
| <sup>3</sup> Σ <sub>g</sub> <sup>+</sup>                  | 4.92           | 4.92  | 4.94   | 4.91 ± 0.06 | 4.89 | 4.89  | 4.89   | 4.87 ± 0.07 | 4.90 | 4.89  | 4.89   | 4.88 | 4.91              | 4.91              | 4.88   | 4.91  | 4.91              | 4.91 | 4.91 | 4.13 |
| <sup>1</sup> Σ <sub>g</sub> <sup>-</sup> [F] <sup>d</sup> | 5.27           | 5.28  | 5.26   | 5.31 ± 0.05 | 5.19 | 5.20  | 5.18   | 5.26 ± 0.09 | 5.06 | 5.07  | 5.07   | 4.97 | 5.05              | 5.05              | 4.97   | 5.05  | 5.05              | 5.05 | 5.05 | 5.05 |
|   | Diacetylene    |       |        |             |      |       |        |             |      |       |        |      | aVTZ              | aVQZ              | aVSZ   | lit.  | exp. <sup>f</sup> |      |      |      |
|   | 6-31+G(d)      |       |        |             |      |       | aVDZ   |             |      |       |        |      | CC3               | CCSDT             | NEVPT2 | CC3   | CC3               | CC3  |      |      |
| <sup>1</sup> Σ <sub>g</sub> <sup>-</sup>                  | 5.57           | 5.58  | 5.56   | 5.52 ± 0.06 | 5.44 | 5.45  | 5.43   | 5.47 ± 0.02 | 5.34 | 5.35  | 5.33   | 5.33 | 5.33              | 5.33              | 5.33   | 5.33  | 5.33              | 5.33 | 4.81 | 4.81 |
| <sup>1</sup> Δ <sub>g</sub>                               | 5.83           | 5.85  | 5.83   | 5.84 ± 0.01 | 5.69 | 5.70  | 5.69   | 5.69 ± 0.02 | 5.61 | 5.62  | 5.61   | 5.61 | 5.60              | 5.60              | 5.61   | 5.60  | 5.60              | 5.60 | 5.06 | 5.06 |
| <sup>3</sup> Σ <sub>g</sub> <sup>+</sup>                  | 4.07           | 4.08  | 4.09   | 4.04 ± 0.07 | 4.06 | 4.06  | 4.06   | 4.07 ± 0.04 | 4.08 | 4.08  | 4.08   | 4.08 | 4.10              | 4.10              | 4.08   | 4.10  | 4.11              | 4.11 | 2.7  | 2.7  |
| <sup>3</sup> Δ <sub>g</sub>                               | 4.93           | 4.93  | 4.92   | 4.94 ± 0.01 | 4.86 | 4.85  | 4.85   | 4.85 ± 0.02 | 4.80 | 4.79  | 4.78   | 4.78 | 4.80              | 4.80              | 4.78   | 4.80  | 4.80              | 4.80 | 3.21 | 3.21 |

<sup>a</sup>All states have a valence  $\pi \rightarrow \pi^*$  character. <sup>b</sup>CASPT2 results from refs 106, 107 and 108 (vacuum UV experiments). <sup>c</sup>Vertical fluorescence energy of the lowest excited state. <sup>d</sup>Experimental 0-0 energies from refs 109 (<sup>3</sup>Σ<sub>g</sub><sup>+</sup>), 110 (<sup>1</sup>Σ<sub>g</sub><sup>-</sup>), and 111 (<sup>1</sup>Δ<sub>g</sub>), all analyzing vacuum electronic spectra. <sup>e</sup>Experimental 0-0 energies from ref 112 (singlet ES, vacuum UV experiment) and ref 113 (triplet ES, EELS). In the latter contribution, the 2.7 eV value for the <sup>3</sup>Σ<sub>g</sub><sup>+</sup> state is the onset, whereas an estimate of the vertical energy (4.2 ± 0.2 eV) is given for the <sup>3</sup>Δ<sub>g</sub> state.

Table 2. Vertical Transition Energies (in eV) for Cyclopropanone, Cyclopropanethione, and Methylene cyclopropene

| state                                | Cyclopropanone |       |        |             |      |       |      |       |        | lit.             |                   |
|--------------------------------------|----------------|-------|--------|-------------|------|-------|------|-------|--------|------------------|-------------------|
|                                      | 6-31+G(d)      |       |        |             | aVDZ |       | aVTZ |       |        | th. <sup>a</sup> | exp. <sup>b</sup> |
|                                      | CC3            | CCSDT | CCSDTQ | FCI         | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 |                  |                   |
| <sup>1</sup> B <sub>1</sub> (n → π*) | 4.32           | 4.34  | 4.36   | 4.38 ± 0.02 | 4.22 | 4.23  | 4.21 | 4.24  | 4.04   | 4.25             | 4.13              |
| <sup>1</sup> A <sub>2</sub> (n → π*) | 5.68           | 5.65  | 5.65   | 5.64 ± 0.06 | 5.59 | 5.56  | 5.57 | 5.55  | 5.85   | 5.59             | 5.5               |
| <sup>1</sup> B <sub>2</sub> (n → 3s) | 6.39           | 6.38  | 6.41   |             | 6.21 | 6.19  | 6.32 | 6.31  | 6.51   | 6.90             | 6.22              |
| <sup>1</sup> B <sub>2</sub> (π → π*) | 6.70           | 6.67  | 6.68   |             | 6.56 | 6.54  | 6.54 | 6.53  | 6.82   | 5.96             | 6.1               |
| <sup>1</sup> B <sub>2</sub> (n → 3p) | 6.92           | 6.91  | 6.94   |             | 6.88 | 6.86  | 6.96 | 6.95  | 7.07   | 7.24             | 6.88              |
| <sup>1</sup> A <sub>1</sub> (n → 3p) | 7.00           | 7.00  | 7.03   |             | 6.88 | 6.87  | 7.00 | 6.99  | 7.28   | 7.28             |                   |
| <sup>1</sup> A <sub>1</sub> (π → π*) | 8.51           | 8.49  | 8.51   |             | 8.32 | 8.29  | 8.28 | 8.26  | 8.19   | 7.80             | ~8.1              |
| <sup>3</sup> B <sub>1</sub> (n → π*) | 4.02           | 4.03  |        | 4.00 ± 0.07 | 3.90 | 3.92  | 3.91 | 3.93  | 3.51   | 4.05             |                   |
| <sup>3</sup> B <sub>2</sub> (π → π*) | 4.92           | 4.92  |        | 4.95 ± 0.00 | 4.90 | 4.89  | 4.89 | 4.88  | 5.10   | 4.81             |                   |
| <sup>3</sup> A <sub>2</sub> (n → π*) | 5.48           | 5.44  |        |             | 5.38 | 5.35  | 5.37 | 5.35  | 5.60   | 5.56             |                   |
| <sup>3</sup> A <sub>1</sub> (π → π*) | 6.89           | 6.88  |        |             | 6.79 | 6.78  | 6.83 | 6.79  | 7.16   | 6.98             |                   |

| state                                | Cyclopropanethione |       |        |             |      |       |      |       |        | lit.             |  |
|--------------------------------------|--------------------|-------|--------|-------------|------|-------|------|-------|--------|------------------|--|
|                                      | 6-31+G(d)          |       |        |             | aVDZ |       | aVTZ |       |        | th. <sup>a</sup> |  |
|                                      | CC3                | CCSDT | CCSDTQ | FCI         | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 |                  |  |
| <sup>1</sup> A <sub>2</sub> (n → π*) | 3.46               | 3.44  | 3.44   | 3.45 ± 0.01 | 3.47 | 3.45  | 3.43 | 3.41  | 3.52   | 3.23             |  |
| <sup>1</sup> B <sub>1</sub> (n → π*) | 3.45               | 3.44  | 3.45   | 3.44 ± 0.05 | 3.42 | 3.42  | 3.43 | 3.44  | 3.50   | 3.47             |  |
| <sup>1</sup> B <sub>2</sub> (π → π*) | 4.67               | 4.64  | 4.62   | 4.59 ± 0.09 | 4.66 | 4.64  | 4.64 | 4.62  | 4.77   | 4.34             |  |
| <sup>1</sup> B <sub>2</sub> (n → 3s) | 5.26               | 5.24  | 5.27   |             | 5.23 | 5.21  | 5.34 | 5.31  | 5.35   | 4.98             |  |
| <sup>1</sup> A <sub>1</sub> (π → π*) | 5.53               | 5.52  | 5.51   |             | 5.52 | 5.50  | 5.49 | 5.47  | 5.54   | 5.52             |  |
| <sup>1</sup> B <sub>2</sub> (n → 3p) | 5.83               | 5.81  | 5.83   |             | 5.86 | 5.84  | 5.93 | 5.90  | 5.99   | 5.88             |  |
| <sup>3</sup> A <sub>2</sub> (n → π*) | 3.33               | 3.31  |        | 3.29 ± 0.03 | 3.34 | 3.32  | 3.30 |       | 3.38   | 3.20             |  |
| <sup>3</sup> B <sub>1</sub> (n → π*) | 3.34               | 3.33  |        |             | 3.30 | 3.30  | 3.31 | 3.32  | 3.40   | 3.30             |  |
| <sup>3</sup> B <sub>2</sub> (π → π*) | 4.01               | 4.00  |        | 4.03 ± 0.03 | 4.03 | 4.02  | 4.02 |       | 4.17   | 3.86             |  |
| <sup>3</sup> A <sub>1</sub> (π → π*) | 4.06               | 4.04  |        |             | 4.09 | 4.07  | 4.03 |       | 4.13   | 3.99             |  |

| state   | Methylene cyclopropene |       |        |             |      |                   |      |       |        |                  |                  |                   |
|---|------------------------|-------|--------|-------------|------|-------------------|------|-------|--------|------------------|------------------|-------------------|
|   | 6-31+G(d)              |       |        |             | aVDZ |                   | aVTZ |       |        | lit.             |                  |                   |
|   | CC3                    | CCSDT | CCSDTQ | FCI         | CC3  | CCSDT             | CC3  | CCSDT | NEVPT2 | th. <sup>c</sup> | th. <sup>d</sup> | exp. <sup>e</sup> |
| <sup>1</sup> B <sub>2</sub> (π → π*)              | 4.38                   | 4.37  | 4.34   | 4.32 ± 0.03 | 4.32 | 4.31              | 4.31 | 4.31  | 4.37   | 4.13             | 4.36             | 4.01              |
| <sup>1</sup> B <sub>1</sub> (π → 3s)              | 5.65                   | 5.66  | 5.66   |             | 5.35 | 5.35              | 5.44 | 5.44  | 5.49   | 5.32             | 5.44             | 5.12              |
| <sup>1</sup> A <sub>2</sub> (π → 3p)              | 5.97                   | 5.98  | 5.98   | 5.92 ± 0.10 | 5.86 | 5.88              | 5.95 | 5.96  | 6.00   | 5.83             |                  |                   |
| <sup>1</sup> A <sub>1</sub> (π → π*) <sup>f</sup> | 6.17                   | 6.18  | 6.17   | 6.20 ± 0.01 | 6.15 | 6.15              | 6.13 | 6.13  | 6.36   |                  | 6.13             | 6.02              |
| <sup>3</sup> B <sub>2</sub> (π → π*)              | 3.50                   | 3.50  |        | 3.44 ± 0.06 | 3.49 | 3.49 <sup>g</sup> | 3.50 | 3.49  | 3.66   | 3.24             |                  |                   |
| <sup>3</sup> A <sub>1</sub> (π → π*)              | 4.74                   | 4.74  |        | 4.67 ± 0.10 | 4.74 | 4.74 <sup>g</sup> | 4.74 |       | 4.87   | 4.52             |                  |                   |

<sup>a</sup>CASPT2 results from ref 124. <sup>b</sup>Electron impact experiment from ref 126. Note that the 5.5 eV peak was assigned differently in the original paper, and we follow here the analysis of Serrano-Andrés,<sup>124</sup> whereas the 6.1 eV assignment was “supposed” in the original paper; experimental  $\lambda_{\max}$  values have been measured at 3.62 and 6.52 eV for the <sup>1</sup>B<sub>1</sub>(n → π\*) and <sup>1</sup>B<sub>2</sub>(π → π\*) transitions, respectively.<sup>127</sup> <sup>c</sup>CASPT2 results from refs 35 and 37. <sup>d</sup>CC3 results from ref 27. <sup>e</sup> $\lambda_{\max}$  in pentane at -78 °C from ref 128. <sup>f</sup>Significant state mixing with the <sup>1</sup>A<sub>1</sub>(π → 3p) transition, making unambiguous attribution difficult. <sup>g</sup>As can be seen in the SI, our FCI/aVDZ estimates are 3.45 ± 0.04 and 4.79 ± 0.02 eV for the two lowest triplet states of methylene cyclopropene hinting that the CC3 and CCSDT results might be slightly too low for the second transition.

**2.3. NEVPT2.** The NEVPT2 calculations have been performed with MOLPRO<sup>83</sup> within the partially contracted scheme (PC-NEVPT2), which is theoretically superior to its strongly contracted version due to the larger number of perturbors and greater flexibility.<sup>84–86</sup> These NEVPT2 calculations are performed on top of a state-averaged complete-active-space self-consistent field calculation always including at least the ground state with the excited state of interest. Active spaces carefully chosen and tailored for the desired transitions have been selected. The definition of the active space considered for each system as well as the number of states in the state-averaged calculation are provided in the SI.

**2.4. Other Wave Function Calculations.** For the other levels of theory, we apply a variety of programs, namely, CFOUR,<sup>79</sup> DALTON,<sup>78</sup> GAUSSIAN,<sup>87</sup> ORCA,<sup>88</sup> MRCC,<sup>89,90</sup>

and Q-CHEM.<sup>91</sup> CFOUR is used for CC3,<sup>92,93</sup> CCSDT-3,<sup>94,95</sup> CCSDT,<sup>96</sup> and CCSDTQ;<sup>97</sup> DALTON for CC2,<sup>98,99</sup> CCSD,<sup>100</sup> CCSDR(3),<sup>101</sup> and CC3;<sup>92,93</sup> GAUSSIAN for CIS(D);<sup>102,103</sup> ORCA for the similarity-transformed (ST) equation-of-motion (EOM) CCSD (STEOM-CCSD);<sup>66,104</sup> MRCC for CCSDT<sup>96</sup> and CCSDTQ,<sup>97</sup> and Q-CHEM for ADC(2).<sup>105</sup> Default program settings were applied. We note that for STEOM-CCSD we only report states that are characterized by an active character percentage of 98% or larger. In all the softwares mentioned above, point group symmetry was systematically employed to reduce the computational effort. It should be noted that we do not perform “GS” CC calculations in a specific symmetry to deduce ES energies. All the CC results reported below correspond to excited-state calculations within the EOM or linear-response (LR) formalisms, both delivering strictly

identical results for transition energies. These formalisms are also applied to get the triplet ES energies directly from the closed-shell singlet GS. In other words, all our calculations systematically consider a restricted closed-shell ground state. Finally, the reported CC3 oscillator strengths have been determined within the LR formalism.

### 3. MAIN RESULTS

In the following, we present results obtained for molecules containing four, five, and six (non-hydrogen) atoms. In all cases, we test several atomic basis sets and push the CC excitation order as high as technically possible. Given that the SCI energy converges rather slowly for these systems, we provide an estimated error bar for these extrapolated FCI values (*vide supra*). In most cases, these extrapolated FCI reference data are used as a “safety net” to demonstrate the overall consistency of the various approaches rather than as definitive reference values (see the next section). As a further consistency check, we also report NEVPT2/aVTZ excitation energies for all states. We underline that, except when specifically discussed, all ESs present a dominant single-excitation character (see also next section) so that we do not expect serious CC breakdowns. This is especially true for triplet ESs that are known to be characterized by very large  $\%T_1$  values in the vast majority of the cases.<sup>23</sup> Consequently, we concentrate most of our computational effort on the obtention of accurate transition energies for singlet states. To assign the different ESs, we use literature data, as well as the usual criteria, i.e., relative energies, spatial and spin symmetries, compositions from the underlying molecular orbitals, and oscillator strengths. This allows clear-cut assignments for the vast majority of the cases. There are however some state/method combinations for which strong mixing between ESs of the same symmetry makes unambiguous assignments almost impossible.

#### 3.1. Molecules with Four Non-Hydrogen Atoms.

**3.1.1. Cyanoacetylene, Cyanogen, and Diacetylene.** The ESs of these three closely related linear molecules containing two triple bonds have been quite rarely theoretically investigated,<sup>106,114–117</sup> though (rather old) experimental measurements of their 0-0 energies are available for several ESs.<sup>107–113,118</sup> Our main results are collected in Tables 1 and S1. We consider only low-lying valence  $\pi \rightarrow \pi^*$  transitions, which are all characterized by a strongly dominant single excitation nature ( $\%T_1 > 90\%$ , *vide infra*). For cyanoacetylene, the FCI/6-31+G(d) estimates come with small error bars, and one notices an excellent agreement between these values and their CCSDTQ counterparts, a statement holding for the Dunning double- $\zeta$  basis set results for which the FCI uncertainties are however larger. Using the CCSDTQ values as references, it appears that the previously obtained CASPT2 estimates<sup>106</sup> are, as expected, too low and that the CC3 transition energies are slightly more accurate than their CCSDT counterparts, although all CC estimates of Table 1 come, for a given basis set, in a very tight energetic window. There is also a very neat agreement between the CC/aVTZ and NEVPT2/aVTZ. All these facts provide strong evidence that the CC estimates can be fully trusted for these three linear systems. The basis set effects are quite significant for the valence ES of cyanoacetylene with successive drops of the transition energies by approximately 0.10 eV, when going from 6–31+G(d) to aVDZ and from aVDZ to aVTZ. The lowest triplet state appears less basis set sensitive, though. As expected, extending further the basis set size (to quadruple- and quintuple- $\zeta$ ) leaves the results pretty much

unchanged. The same observation holds when adding a second set of diffuse functions or when correlating the core electrons (see the SI). Obviously, both cyanogen and diacetylene yield very similar trends, with limited methodological effects and quite large basis set effects, except for the  $^1\Sigma_g^+ \rightarrow ^3\Sigma_u^+$  transitions. We note that all CC3 and CCSDT values are, at worst, within  $\pm 0.02$  eV of the FCI window, i.e., all methods presented in Table 1 provide very consistent estimates. For all the states reported in Table 1, the average absolute deviation between NEVPT2/aVTZ and CC3/aVTZ (CCSDT/aVTZ) is as small as 0.02 (0.03) eV, the lowest absorption and emission energies of cyanogen being the only two cases showing significant deviations. As a final note, all our vertical absorption (emission) energies are significantly larger (smaller) than the experimentally measured 0-0 energies, as they should be. We refer the interested reader to previous works,<sup>114,117</sup> for comparisons between theoretical (CASPT2 and CC3) and experimental 0-0 energies for these three compounds.

**3.1.2. Cyclopropanone, Cyclopropanethione, and Methylene-cyclopropane.** These three related compounds present a three-membered  $sp^2$  carbon cycle conjugated to an external  $\pi$  bond. While the ESs of methylenecyclopropane have regularly been investigated with theoretical tools in the past,<sup>27,35,37,75,119–123</sup> the only investigations of vertical transitions we could find for the two other derivatives are a detailed CASPT2 study of Serrano-Andrés and co-workers in 2002<sup>124</sup> and a more recent work reporting the three lowest-lying singlet states of cyclopropanone at the CASPT2/6-31G level.<sup>125</sup>

Our results are listed in Tables 2 and S2. As above, considering the 6-31+G(d) basis set, we notice very small differences between CC3, CCSDT, and CCSDTQ, the latter method giving transition energies systematically falling within the FCI extrapolation uncertainty, except in one case (the lowest totally symmetric state of methylenecyclopropane for which the CCSDTQ value is “off” by 0.02 eV only). Depending on the state, it is either CC3 or CCSDT that is closest to CCSDTQ. In fact, considering the CCSDTQ/6-31+G(d) data listed in Table 2 as reference, the MAEs of CC3 and CCSDT are 0.019 and 0.016 eV, respectively, hinting that the improvement brought by the latter, more expensive method is limited for these three compounds. For the lowest  $B_2$  state of methylenecyclopropane, one of the most challenging cases ( $\%T_1 = 85\%$ ), it is clear from the FCI value that only CCSDTQ is energetically close, the CC3 and CCSDT results being slightly too large by  $\sim 0.05$  eV. It seems reasonable to believe that the same observation can be made for the corresponding state of cyclopropanethione, although in this case the FCI error bar is too large, which prevents any definitive conclusion. Interestingly, at the CC3 level of theory, the rather small 6-31+G(d) basis set provides data within 0.10 eV of the CBS limit for 80% of the transitions. There are, of course, exceptions to this rule, e.g., the strongly dipole-allowed  $^1A_1(\pi \rightarrow \pi^*)$  ES of cyclopropanone and the  $^1B_1(\pi \rightarrow 3s)$  ES of methylenecyclopropane which are significantly over blueshifted with the Pople basis set (Table S2). For cyclopropanone, our CCSDT/aVTZ estimates do agree reasonably well with the CASPT2 data of Serrano-Andrés, except for the  $^1B_2(\pi \rightarrow \pi^*)$  state that we locate significantly higher in energy and the three Rydberg states that our CC calculations predict at significantly lower energies. The present NEVPT2 results are globally in better agreement with the CC values, though non-negligible deviations pertain. Even if comparisons with experiment should be made very cautiously, we note that, for the Rydberg states, the present CC data are

Table 3. Vertical Transition Energies (in eV) of Acrolein, Butadiene, and Glyoxal

| state                    | Acrolein  |       |                          |      |       |      |       |        |      | lit.             |                  |                   |
|--------------------------|-----------|-------|--------------------------|------|-------|------|-------|--------|------|------------------|------------------|-------------------|
|                          | 6-31+G(d) |       |                          |      | aVDZ  |      | aVTZ  |        |      | th. <sup>a</sup> | th. <sup>b</sup> | exp. <sup>c</sup> |
|                          | CC3       | CCSDT | FCI                      | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 |      |                  |                  |                   |
| <sup>1</sup> A''(n → π*) | 3.83      | 3.80  | 3.85 ± 0.01              | 3.77 | 3.74  | 3.74 | 3.73  | 3.76   | 3.63 | 3.83             | 3.71             |                   |
| <sup>1</sup> A'(π → π*)  | 6.83      | 6.86  | 6.59 ± 0.05 <sup>f</sup> | 6.67 | 6.70  | 6.65 | 6.69  | 6.67   | 6.10 | 6.92             | 6.41             |                   |
| <sup>1</sup> A''(n → π*) | 6.94      | 6.89  |                          | 6.75 | 6.72  | 6.75 |       | 7.16   | 6.26 | 7.40             |                  |                   |
| <sup>1</sup> A'(n → 3s)  | 7.22      | 7.23  |                          | 6.99 | 7.00  | 7.07 |       | 7.05   | 6.97 | 7.19             | 7.08             |                   |
| <sup>3</sup> A''(n → π*) | 3.55      | 3.53  | 3.60 ± 0.01              | 3.47 | 3.45  | 3.46 |       | 3.46   | 3.39 | 3.61             |                  |                   |
| <sup>3</sup> A'(π → π*)  | 3.94      | 3.95  | 3.98 ± 0.03              | 3.95 | 3.95  | 3.94 |       | 3.95   | 3.81 | 3.87             |                  |                   |
| <sup>3</sup> A'(π → π*)  | 6.25      | 6.23  |                          | 6.22 | 6.21  | 6.19 |       | 6.23   |      | 6.21             |                  |                   |
| <sup>3</sup> A''(n → π*) | 6.81      | 6.74  |                          | 6.60 |       | 6.61 |       | 6.83   |      | 7.36             |                  |                   |

| state                                | Butadiene |       |        |                          |      |       |      |       |        | lit.             |                  |                   |
|--------------------------------------|-----------|-------|--------|--------------------------|------|-------|------|-------|--------|------------------|------------------|-------------------|
|                                      | 6-31+G(d) |       |        |                          | aVDZ |       | aVTZ |       |        | th. <sup>b</sup> | th. <sup>d</sup> | exp. <sup>e</sup> |
|                                      | CC3       | CCSDT | CCSDTQ | FCI                      | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 |                  |                  |                   |
| <sup>1</sup> B <sub>u</sub> (π → π*) | 6.41      | 6.43  | 6.41   | 6.41 ± 0.02              | 6.25 | 6.27  | 6.22 | 6.24  | 6.68   | 6.33             | 6.36             | 5.92              |
| <sup>1</sup> B <sub>g</sub> (π → 3s) | 6.53      | 6.55  | 6.54   |                          | 6.26 | 6.27  | 6.33 | 6.34  | 6.44   | 6.18             | 6.32             | 6.21              |
| <sup>1</sup> A <sub>g</sub> (π → π*) | 6.73      | 6.63  | 6.56   | 6.55 ± 0.04 <sup>f</sup> | 6.68 | 6.59  | 6.67 | 6.60  | 6.70   | 6.56             | 6.60             |                   |
| <sup>1</sup> A <sub>u</sub> (π → 3p) | 6.87      | 6.89  | 6.87   |                          | 6.57 | 6.59  | 6.64 | 6.66  | 6.84   | 6.45             | 6.56             | 6.64              |
| <sup>1</sup> A <sub>u</sub> (π → 3p) | 6.93      | 6.95  | 6.94   | 6.95 ± 0.01              | 6.73 | 6.74  | 6.80 | 6.81  | 7.01   | 6.65             | 6.74             | 6.80              |
| <sup>1</sup> B <sub>u</sub> (π → 3p) | 7.98      | 8.00  | 7.98   |                          | 7.86 | 7.87  | 7.68 |       | 7.45   | 7.08             | 7.02             | 7.07              |
| <sup>3</sup> B <sub>u</sub> (π → π*) | 3.35      | 3.36  |        | 3.37 ± 0.03              | 3.36 | 3.36  | 3.36 |       | 3.40   | 3.20             |                  | 3.22              |
| <sup>3</sup> A <sub>g</sub> (π → π*) | 5.22      | 5.22  |        |                          | 5.21 | 5.21  | 5.20 |       | 5.30   | 5.08             |                  | 4.91              |
| <sup>3</sup> B <sub>g</sub> (π → 3s) | 6.46      | 6.47  |        | 6.40 ± 0.03              | 6.20 | 6.21  | 6.28 |       | 6.38   | 6.14             |                  |                   |

| state                                       | Glyoxal   |       |        |                          |      |       |      |       |        | lit.             |                  |                   |
|---|-----------|-------|--------|--------------------------|------|-------|------|-------|--------|------------------|------------------|-------------------|
|   | 6-31+G(d) |       |        |                          | aVDZ |       | aVTZ |       |        | th. <sup>b</sup> | th. <sup>g</sup> | exp. <sup>h</sup> |
|   | CC3       | CCSDT | CCSDTQ | FCI                      | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 |                  |                  |                   |
| <sup>1</sup> A <sub>u</sub> (n → π*)        | 2.94      | 2.94  | 2.94   | 2.93 ± 0.03              | 2.90 | 2.90  | 2.88 | 2.88  | 2.90   | 3.10             | 2.93             | 2.8               |
| <sup>1</sup> B <sub>g</sub> (n → π*)        | 4.34      | 4.32  | 4.31   | 4.28 ± 0.06              | 4.30 | 4.28  | 4.27 | 4.25  | 4.30   | 4.68             | 4.39             | ~4.4              |
| <sup>1</sup> A <sub>g</sub> (n, n → π*, π*) | 6.74      | 6.24  | 5.67   | 5.60 ± 0.09 <sup>f</sup> | 6.70 | 6.22  | 6.76 | 6.35  | 5.52   | 5.66             |                  |                   |
| <sup>1</sup> B <sub>g</sub> (n → π*)        | 6.81      | 6.83  | 6.79   |                          | 6.59 | 6.61  | 6.58 | 6.61  | 6.64   | 7.54             | 6.63             | 7.45              |
| <sup>1</sup> B <sub>u</sub> (n → 3p)        | 7.72      | 7.74  | 7.76   |                          | 7.55 | 7.56  | 7.67 | 7.69  | 7.84   | 7.83             | 7.61             | ~7.7              |
| <sup>3</sup> A <sub>u</sub> (n → π*)        | 2.55      | 2.55  |        | 2.54 ± 0.04              | 2.49 | 2.49  | 2.49 | 2.49  | 2.49   | 2.63             |                  | 2.5               |
| <sup>3</sup> B <sub>g</sub> (n → π*)        | 3.97      | 3.95  |        |                          | 3.91 | 3.90  | 3.90 | 3.89  | 3.99   | 4.12             |                  | ~3.8              |
| <sup>3</sup> B <sub>u</sub> (π → π*)        | 5.22      | 5.20  |        |                          | 5.20 | 5.19  | 5.17 | 5.15  | 5.17   | 5.35             |                  | ~5.2              |
| <sup>3</sup> A <sub>g</sub> (π → π*)        | 6.35      | 6.35  |        |                          | 6.34 | 6.34  | 6.30 | 6.30  | 6.33   |                  |                  |                   |

<sup>a</sup>CASPT2 results from ref 129. <sup>b</sup>SAC-CI results from ref 130. <sup>c</sup>Vacuum UV spectra from ref 150; for the lowest state, the same 3.71 eV value is reported in ref 151. <sup>d</sup>MR-AQCC results from ref 134, with theoretical best estimates listed for the lowest B<sub>u</sub> and A<sub>g</sub> states. <sup>e</sup>Electron impact experiment from refs 152 and 153 for the singlet states and from ref 154 for the two lowest triplet transitions; note that, for the lowest B<sub>u</sub> state, there is a vibrational structure with peaks at 5.76, 5.92, and 6.05 eV. <sup>f</sup>From ref 72. <sup>g</sup>CC3 results from ref 27. <sup>h</sup>Electron impact experiment from ref 155 except for the second <sup>1</sup>B<sub>g</sub> ES for which the value is from another work (see ref 156); note that for the lowest <sup>1</sup>B<sub>g</sub>(<sup>1</sup>B<sub>u</sub>) ES, a range of 4.2–4.5 (7.4–7.9) eV is given in ref 155.

clearly more consistent with the electron impact measurements<sup>126</sup> than the original CASPT2 values. For cyclopropenethione, we typically obtain transition energies in agreement with or larger than those obtained with CASPT2,<sup>124</sup> though there is no obvious relationship between the valence/Rydberg nature of the ES and the relative CASPT2 error. The average absolute deviation between our NEVPT2 and CC3 results is 0.08 eV only. Finally, in the case of methylenecyclopropene, our values logically agree very well with the recent estimates of Schwabe and Goerigk,<sup>27</sup> obtained at the CC3/aVTZ level of theory on a different geometry. As anticipated, the available CASPT2 values,<sup>35,37</sup> determined without IPEA shift, appear too low as compared to the present NEVPT2 and CCSDT values. For this compound, the available experimental data are based on the wavelength of maximal

absorption determined in the condensed phase.<sup>128</sup> Hence, only a qualitative match is reached between theory and experiment.

**3.1.3. Acrolein, Butadiene, and Glyoxal.** Let us now turn our attention to the excited states of three pseudolinear π-conjugated systems that have been the subject to several investigations in the past, namely, acrolein,<sup>27,119–121,129–133</sup> butadiene,<sup>23,26,27,72,120,130,134–145</sup> and glyoxal.<sup>27,116,120,130,132,146–149</sup> Among these works, it is worth highlighting the detailed theoretical investigation of Saha, Ehara, and Nakatsuji, who reported a huge number of ESs for these three systems using a coherent theoretical protocol based on the symmetry-adapted-cluster configuration interaction (SAC-CI) method.<sup>130</sup> In the following, these three molecules are considered in their most stable *trans* conformation. Our results are listed in Tables 3 and S3.

**Table 4. Vertical Transition Energies (in eV) of Acetone, Cyanoformaldehyde, Isobutene, Propynal, Thioacetone, and Thiopropynal**

| Acetone                              |           |       |             |             |       |       |       |        |                   |                   |                   |                   |
|--------------------------------------|-----------|-------|-------------|-------------|-------|-------|-------|--------|-------------------|-------------------|-------------------|-------------------|
| state                                | 6-31+G(d) |       |             |             | aVDZ  |       | aVTZ  |        |                   | lit.              |                   |                   |
|                                      | CC3       | CCSDT | CCSDTQ      | FCI         | CC3   | CCSDT | CC3   | CCSDT  | NEVPT2            | th. <sup>a</sup>  | th. <sup>b</sup>  | exp. <sup>c</sup> |
| <sup>1</sup> A <sub>2</sub> (n → π*) | 4.55      | 4.52  | 4.53        | 4.60 ± 0.05 | 4.50  | 4.48  | 4.48  | 4.46   | 4.48              | 4.18              | 4.18              | 4.48              |
| <sup>1</sup> B <sub>2</sub> (n → 3s) | 6.65      | 6.64  | 6.68        |             | 6.31  | 6.30  | 6.43  | 6.42   | 6.81              | 6.58              | 6.58              | 6.36              |
| <sup>1</sup> A <sub>2</sub> (n → 3p) | 7.83      | 7.83  | 7.87        |             | 7.37  | 7.36  | 7.45  | 7.43   | 7.65              | 7.34              | 7.34              | 7.36              |
| <sup>1</sup> A <sub>1</sub> (n → 3p) | 7.81      | 7.81  | 7.84        |             | 7.39  | 7.38  | 7.48  | 7.48   | 7.75              | 7.26              | 7.26              | 7.41              |
| <sup>1</sup> B <sub>2</sub> (n → 3p) | 7.87      | 7.87  | 7.91        |             | 7.56  | 7.55  | 7.59  | 7.58   | 7.91              | 7.48              | 7.48              | 7.45              |
| <sup>3</sup> A <sub>2</sub> (n → π*) | 4.21      | 4.19  |             | 4.18 ± 0.04 | 4.16  | 4.14  | 4.15  |        | 4.20              | 3.90              | 3.90              | 4.15              |
| <sup>3</sup> A <sub>1</sub> (π → π*) | 6.32      | 6.30  |             |             | 6.31  | 6.28  | 6.28  |        | 6.28              | 5.98              | 5.98              |                   |
| Cyanoformaldehyde                    |           |       |             |             |       |       |       |        |                   |                   |                   |                   |
| state                                | 6-31+G(d) |       |             | aVDZ        |       | aVTZ  |       |        | lit.              |                   |                   |                   |
|                                      | CC3       | CCSDT | FCI         | CC3         | CCSDT | CC3   | CCSDT | NEVPT2 | exp. <sup>d</sup> |                   |                   |                   |
| <sup>1</sup> A''(n → π*)             | 3.91      | 3.89  | 3.92 ± 0.02 | 3.86        | 3.84  | 3.83  | 3.81  | 3.98   |                   |                   |                   | 3.26              |
| <sup>1</sup> A''(π → π*)             | 6.64      | 6.67  | 6.60 ± 0.07 | 6.51        | 6.54  | 6.42  | 6.46  | 6.44   |                   |                   |                   |                   |
| <sup>3</sup> A''(n → π*)             | 3.53      | 3.51  | 3.48 ± 0.06 | 3.47        | 3.45  | 3.46  |       | 3.58   |                   |                   |                   |                   |
| <sup>3</sup> A'(π → π*)              | 5.07      | 5.07  |             | 5.03        | 5.03  | 5.01  |       | 5.35   |                   |                   |                   |                   |
| Isobutene                            |           |       |             |             |       |       |       |        |                   |                   |                   |                   |
| state                                | 6-31+G(d) |       |             | aVDZ        |       | aVTZ  |       |        | lit.              |                   |                   |                   |
|                                      | CC3       | CCSDT | FCI         | CC3         | CCSDT | CC3   | CCSDT | NEVPT2 | th. <sup>e</sup>  | exp. <sup>f</sup> | exp. <sup>g</sup> |                   |
| <sup>1</sup> B <sub>1</sub> (π → 3s) | 6.77      | 6.77  | 6.78 ± 0.08 | 6.39        | 6.39  | 6.45  | 6.46  | 6.63   | 6.40              | 6.15              | 6.17              |                   |
| <sup>1</sup> A <sub>1</sub> (π → 3p) | 7.16      | 7.17  | 7.16 ± 0.02 | 7.00        | 7.00  | 7.00  | 7.01  | 7.20   | 6.96              |                   | 6.71              |                   |
| <sup>3</sup> A <sub>1</sub> (π → π*) | 4.52      | 4.53  | 4.56 ± 0.02 | 4.54        | 4.54  | 4.53  |       | 4.61   |                   | 4.21              | 4.3               |                   |
| Propynal                             |           |       |             |             |       |       |       |        |                   |                   |                   |                   |
| state                                | 6-31+G(d) |       |             | aVDZ        |       | aVTZ  |       |        | lit.              |                   |                   |                   |
|                                      | CC3       | CCSDT | FCI         | CC3         | CCSDT | CC3   | CCSDT | NEVPT2 | exp. <sup>h</sup> |                   |                   |                   |
| <sup>1</sup> A''(n → π*)             | 3.90      | 3.87  | 3.84 ± 0.06 | 3.85        | 3.82  | 3.82  | 3.80  | 3.95   |                   |                   | 3.24              |                   |
| <sup>1</sup> A''(π → π*)             | 5.69      | 5.73  | 5.64 ± 0.08 | 5.59        | 5.62  | 5.51  | 5.54  | 5.50   |                   |                   |                   |                   |
| <sup>3</sup> A''(n → π*)             | 3.56      | 3.54  | 3.54 ± 0.04 | 3.50        | 3.48  | 3.49  |       | 3.59   |                   |                   | 2.99              |                   |
| <sup>3</sup> A'(π → π*)              | 4.46      | 4.47  | 4.44 ± 0.08 | 4.40        | 4.44  | 4.43  |       | 4.63   |                   |                   |                   |                   |
| Thioacetone                          |           |       |             |             |       |       |       |        |                   |                   |                   |                   |
| state                                | 6-31+G(d) |       |             |             | aVDZ  |       | aVTZ  |        |                   | lit.              |                   |                   |
|                                      | CC3       | CCSDT | CCSDTQ      | FCI         | CC3   | CCSDT | CC3   | CCSDT  | NEVPT2            | exp. <sup>i</sup> |                   |                   |
| <sup>1</sup> A <sub>2</sub> (n → π*) | 2.58      | 2.56  | 2.56        | 2.61 ± 0.05 | 2.59  | 2.57  | 2.55  | 2.53   | 2.55              |                   | 2.33              |                   |
| <sup>1</sup> B <sub>2</sub> (n → 4s) | 5.65      | 5.64  | 5.66        |             | 5.44  | 5.43  | 5.55  | 5.54   | 5.72              |                   | 5.49              |                   |
| <sup>1</sup> A <sub>1</sub> (π → π*) | 6.09      | 6.10  | 6.07        |             | 5.97  | 5.98  | 5.90  | 5.91   | 6.24              |                   | 5.64              |                   |
| <sup>1</sup> B <sub>2</sub> (n → 4p) | 6.59      | 6.59  | 6.59        |             | 6.45  | 6.44  | 6.51  |        | 6.62              |                   | 6.40              |                   |
| <sup>1</sup> A <sub>1</sub> (n → 4p) | 6.95      | 6.95  | 6.96        |             | 6.54  | 6.53  | 6.61  | 6.60   | 6.52              |                   | 6.52              |                   |
| <sup>3</sup> A <sub>2</sub> (n → π*) | 2.36      | 2.34  |             | 2.36 ± 0.00 | 2.36  | 2.35  | 2.34  |        | 2.32              |                   | 2.14              |                   |
| <sup>3</sup> A <sub>1</sub> (π → π*) | 3.45      | 3.45  |             |             | 3.51  | 3.50  | 3.46  |        | 3.48              |                   |                   |                   |
| Thiopropynal                         |           |       |             |             |       |       |       |        |                   |                   |                   |                   |
| state                                | 6-31+G(d) |       |             | aVDZ        |       | aVTZ  |       |        | lit.              |                   |                   |                   |
|                                      | CC3       | CCSDT | FCI         | CC3         | CCSDT | CC3   | CCSDT | NEVPT2 | exp. <sup>j</sup> |                   |                   |                   |
| <sup>1</sup> A''(n → π*)             | 2.09      | 2.06  | 2.08 ± 0.01 | 2.09        | 2.06  | 2.05  | 2.03  | 2.05   |                   |                   | 1.82              |                   |
| <sup>3</sup> A''(n → π*)             | 1.84      | 1.82  |             | 1.83        | 1.82  | 1.81  |       | 1.81   |                   |                   | 1.64              |                   |

<sup>a</sup>CASPT2 results from ref 36. <sup>b</sup>EOM-CCSD results from ref 157. <sup>c</sup>Two lowest singlet states: various experiments summarized in ref 169; three next singlet states: REMPI experiments from ref 172; lowest triplet: trapped electron measurements from ref 166. <sup>d</sup>0-0 energy reported in ref 171. <sup>e</sup>EOM-CCSD results from ref 160. <sup>f</sup>Energy loss experiment from ref 167. <sup>g</sup>VUV experiment from ref 170 (we report the lowest of the π → 3p state for the <sup>1</sup>A<sub>1</sub> state). <sup>h</sup>0-0 energies from refs 165 (singlet) and 163 (triplet). <sup>i</sup>0-0 energies from ref 164. <sup>j</sup>0-0 energies from ref 168.

Acrolein, due to its lower symmetry and high density of ES with mixed characters, is challenging from a theoretical point of view, and CCSDTQ calculations were technically impossible despite all our efforts. For the lowest n → π\* transitions of both spin symmetries, the FCI estimates come with a tiny error bar,

and it is obvious that the CC excitation energies are slightly too low, especially with CCSDT. Nevertheless, at the exception of the second singlet and triplet A'' ESs, the CC3 and CCSDT transition energies are within ±0.03 eV of each other. These A'' ESs are also the only two transitions for which the discrepancies



between CC3 and NEVPT2 exceed 0.20 eV. This hints at a good accuracy for all other transitions. This statement is additionally supported by the fact that the present CC values are nearly systematically bracketed by previous CASPT2 (lower bound)<sup>129</sup> and SAC-CI (upper bound)<sup>130</sup> results, consistently with the typical error sign of these two models. For the two lowest triplet states, the present CC3/aVTZ values are also within  $\pm 0.05$  eV of the recent MRCI estimates (3.50 and 3.89 eV).<sup>131</sup> As can be seen in Table S3, the aVTZ basis set delivers excitation energies very close to the CBS limit: the largest variation when upgrading from aVTZ to aVQZ (+0.04 eV) is obtained for the second  $^1A'$  Rydberg ES. As experimental data are limited to measured UV spectra,<sup>150,151</sup> one has to be ultra cautious in establishing TBES for acrolein (*vide infra*).

The nature and relative energies of the lowest bright  $B_u$  and dark  $A_g$  ESs of butadiene have puzzled theoretical chemists for many years. It is beyond the scope of the present study to provide an exhaustive list of previous calculations and experimental measurements for these two hallmark ESs, and we refer the readers to refs 137 and 141 for a general and broader overview. For the  $B_u$  transition, we believe that the most solid TBE is the 6.21 eV value obtained by Watson and Chan using a computational strategy similar to ours.<sup>137</sup> Our CCSDT/aVTZ value of 6.24 eV is obviously compatible with their reference value, and our TBE/CBS value is actually 6.21 eV as well (*vide infra*). For the  $A_g$  state, we believe that our previous basis set corrected FCI estimate of 6.50 eV<sup>72</sup> remains the most accurate available to date. These two values are slightly lower than the semistochastic heath-bath CI data obtained by Chien et al. with a double- $\zeta$  basis and a slightly different geometry: 6.45 and 6.58 eV for  $B_u$  and  $A_g$ , respectively.<sup>143</sup> For these two thoroughly studied ESs, one can of course find many other estimates, e.g., at the SAC-CI,<sup>130</sup> CC3,<sup>26,27</sup> CASPT2,<sup>26</sup> and NEVPT2<sup>142</sup> levels. Globally, for butadiene, we find an excellent coherence between the CC3, CCSDT, and CCSDTQ estimates, that all fall in a  $\pm 0.02$  eV window. Unsurprisingly, this does not apply to the already mentioned  $^1A_g$  ES that is, respectively, 0.2 and 0.1 eV too high with the two former CC methods, a direct consequence of the large electronic reorganization taking place during this transition. For all the other butadiene ESs listed in Table 3, both CC3 and CCSDT can be trusted. We also note that the NEVPT2 estimates are within 0.1–0.2 eV of the CC values, except for the lowest  $B_u$  ES for which the associated excitation energy is highly dependent on the selected active space (see the SI). Finally, as can be seen in Table S3, aVTZ produces near-CBS excitation energies for most ESs. However, a significant basis set effect exists for the Rydberg  $^1B_u(\pi \rightarrow 3p)$  ES with an energy lowering as large as  $-0.12$  eV when going from aVTZ to aVQZ. For the record, we note that the available electron impact data<sup>152–154</sup> provides the very same ES ordering as our calculations.

Globally, the conclusions obtained for acrolein and butadiene pertain for glyoxal, i.e., highly consistent CC estimates, reasonable agreement between NEVPT2 and CC3 values, and limited basis set effects beyond aVTZ, except for the  $^1B_u(n \rightarrow 3p)$  Rydberg state (see Tables 3 and S3). This Rydberg state also exhibits an unexpectedly large deviation of 0.04 eV between CC3 and CCSDTQ. More interestingly, glyoxal presents a genuine low-lying double ES of  $^1A_g$  symmetry. The corresponding  $(n, n) \rightarrow (\pi^*, \pi^*)$  transition is totally unseen by approaches that cannot model double excitations, e.g., TD-DFT, CCSD, or ADC(2). Compared to the FCI values, the CC3 and CCSDT estimates associated with this transition are too large by  $\sim 1.0$

and  $\sim 0.5$  eV, respectively, whereas both the CCSDTQ and the NEVPT2 approaches are much closer, as already mentioned in our previous work.<sup>72</sup> For the other transitions, the present CC3 estimates are logically consistent with the values of ref 27 obtained with the same approach on a different geometry and remain slightly lower than the SAC-CI estimates of ref 130. Once more, the experimental data<sup>155,156</sup> are unhelpful in view of the targeted accuracy.

**3.1.4. Acetone, Cyanoformaldehyde, Isobutene, Propynal, Thioacetone, and Thiopropynal.** Let us now turn toward six other compounds with four non-hydrogen atoms. There are several earlier studies reporting estimates of the vertical transition energies for both acetone<sup>23,26,27,36,37,39,121,139,157–161</sup> and isobutene.<sup>139,159,160</sup> To the best of our knowledge, for the four other compounds, the previous computational efforts were mainly focused on the 0-0 energies of the lowest-lying state.<sup>116,117,147,148,162</sup> There are also rather few experimental data available for these six derivatives.<sup>163,164,166–172</sup> Our main results are reported in Tables 4 and S4.

For acetone, one should clearly distinguish the valence ES, for which both methodological and basis set effects are small, and the Rydberg transitions that are both very basis set sensitive and upshifted by ca. 0.04 eV with CCSDTQ as compared to CC3 and CCSDT. For this compound, the 1996 CASPT2 transition energies of Merchán and co-workers listed on the right panel of Table 4 are clearly too low, especially for the three valence ESs.<sup>36</sup> As expected, this error can be partially ascribed to the computational setup, as the Urban group obtained CASPT2 excitation energies of 4.40, 4.09, and 6.22 eV for the  $^1A_2$ ,  $^3A_2$ , and  $^3A_1$  ESs,<sup>161</sup> in much better agreement with ours. Their estimates of the three  $n \rightarrow 3p$  transitions, 7.52, 7.57, and 7.53 eV for the  $^1A_2$ ,  $^1A_1$ , and  $^1B_2$  ESs, also systematically fall within 0.10 eV of our current CC values, whereas for these three ESs, the current NEVPT2 values are clearly too large.

In contrast to acetone, both valence and Rydberg ESs of thioacetone are rather insensitive to the excitation order of the CC expansion as illustrated by the maximal discrepancies of  $\pm 0.02$  eV between the CC3/6-31+G(d) and CCSDTQ/6-31+G(d) results. While the lowest  $n \rightarrow \pi^*$  transition of both spin symmetries are rather basis set insensitive, all the other states need quite large one-electron bases to be correctly described (Table S4). As expected, our theoretical vertical transition energies show the same ranking but are systematically larger than the available experimental 0-0 energies.

For the isoelectronic isobutene molecule, we have considered two singlet Rydberg ESs and one triplet valence ES. For these three cases, we note, for each basis, a very nice agreement between CC3 and CCSDT, the CC results being also very close to the FCI estimates obtained with the Pople basis set. The similarity with the CCSD results of Caricato and co-workers<sup>160</sup> is also very satisfying.

For the three remaining compounds, namely, cyanoformaldehyde, propynal, and thiopropynal, we report low-lying valence transitions with a definite single excitation character. The basis set effects are clearly under control (they are only significant for the second  $^1A''$  ES of cyanoformaldehyde), and we could not detect any variation larger than 0.03 eV between the CC3 and the CCSDT values for a given basis, indicating that the CC values are very accurate. This is further confirmed by the FCI data.

**3.1.5. Intermediate Conclusions.** For the 15 molecules with four non-hydrogen atoms considered here, we find extremely consistent transition energies between CC and FCI estimates in

Table 5. Vertical Transition Energies (in eV) of Furan and Pyrrole

| state   | Furan     |       |      |       |      |       |        |                  |                  |                  |                  |                  |                   |                   |
|---|-----------|-------|------|-------|------|-------|--------|------------------|------------------|------------------|------------------|------------------|-------------------|-------------------|
|   | 6-31+G(d) |       | aVDZ |       | aVTZ |       |        | lit.             |                  |                  |                  |                  |                   |                   |
|   | CC3       | CCSDT | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 | th. <sup>a</sup> | th. <sup>b</sup> | th. <sup>c</sup> | th. <sup>d</sup> | th. <sup>e</sup> | exp. <sup>f</sup> | exp. <sup>g</sup> |
| <sup>1</sup> A <sub>2</sub> ( $\pi \rightarrow 3s$ )    | 6.26      | 6.28  | 6.00 | 6.00  | 6.08 | 6.09  | 6.28   | 5.92             | 6.13             | 5.94             | 5.91             | 6.10             | 5.91              |                   |
| <sup>1</sup> B <sub>2</sub> ( $\pi \rightarrow \pi^*$ ) | 6.50      | 6.52  | 6.37 | 6.39  | 6.34 | 6.37  | 6.20   | 6.04             | 6.42             | 6.51             | 6.10             | 6.42             | 6.04              | 6.06              |
| <sup>1</sup> A <sub>1</sub> ( $\pi \rightarrow \pi^*$ ) | 6.71      | 6.67  | 6.62 | 6.58  | 6.58 | 6.56  | 6.77   | 6.16             | 6.71             | 6.89             | 6.44             |                  |                   | 6.44              |
| <sup>1</sup> B <sub>1</sub> ( $\pi \rightarrow 3p$ )    | 6.76      | 6.77  | 6.55 | 6.56  | 6.63 | 6.64  | 6.71   | 6.46             | 6.68             | 6.46             | 6.45             | 6.66             | 6.47              |                   |
| <sup>1</sup> A <sub>2</sub> ( $\pi \rightarrow 3p$ )    | 6.97      | 6.99  | 6.73 | 6.74  | 6.80 | 6.81  | 6.99   | 6.59             | 6.79             | 6.61             | 6.60             | 6.83             | 6.61              |                   |
| <sup>1</sup> B <sub>2</sub> ( $\pi \rightarrow 3p$ )    | 7.53      | 7.54  | 7.39 | 7.40  | 7.23 |       | 7.01   | 6.48             | 6.91             | 6.87             | 6.72             | 7.36             | 6.75              |                   |
| <sup>3</sup> B <sub>2</sub> ( $\pi \rightarrow \pi^*$ ) | 4.28      | 4.28  | 4.25 | 4.23  | 4.22 |       | 4.42   | 3.99             |                  | 4.26             |                  |                  |                   | 4.0               |
| <sup>3</sup> A <sub>1</sub> ( $\pi \rightarrow \pi^*$ ) | 5.56      | 5.54  | 5.51 | 5.49  | 5.48 |       | 5.60   | 5.15             |                  | 5.53             |                  |                  |                   | 5.2               |
| <sup>3</sup> A <sub>2</sub> ( $\pi \rightarrow 3s$ )    | 6.18      | 6.19  | 5.94 | 5.94  | 6.02 |       | 6.08   | 5.86             |                  | 5.89             |                  |                  |                   |                   |
| <sup>3</sup> B <sub>1</sub> ( $\pi \rightarrow 3p$ )    | 6.69      | 6.71  | 6.51 | 6.51  | 6.59 |       | 6.68   | 6.42             |                  | 6.41             |                  |                  |                   |                   |

| state   | Pyrrole   |       |      |       |      |       |        |                  |                  |                  |                  |                  |                   |                   |
|---|-----------|-------|------|-------|------|-------|--------|------------------|------------------|------------------|------------------|------------------|-------------------|-------------------|
|   | 6-31+G(d) |       | aVDZ |       | aVTZ |       |        | lit.             |                  |                  |                  |                  |                   |                   |
|   | CC3       | CCSDT | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 | th. <sup>a</sup> | th. <sup>h</sup> | th. <sup>c</sup> | th. <sup>i</sup> | th. <sup>j</sup> | exp. <sup>k</sup> | exp. <sup>l</sup> |
| <sup>1</sup> A <sub>2</sub> ( $\pi \rightarrow 3s$ )    | 5.25      | 5.25  | 5.15 | 5.14  | 5.24 | 5.24  | 5.51   | 5.08             | 5.45             | 5.10             | 5.20             | 5.27             | 5.22              |                   |
| <sup>1</sup> B <sub>1</sub> ( $\pi \rightarrow 3p$ )    | 5.99      | 5.98  | 5.89 | 5.87  | 5.98 | 6.00  | 6.32   | 5.85             | 6.21             | 5.79             | 5.95             | 6.00             |                   |                   |
| <sup>1</sup> A <sub>2</sub> ( $\pi \rightarrow 3p$ )    | 6.27      | 6.27  | 5.94 | 5.93  | 6.01 |       | 6.44   | 5.83             | 6.14             | 5.81             | 5.94             | 7.03             |                   | 5.87              |
| <sup>1</sup> B <sub>2</sub> ( $\pi \rightarrow \pi^*$ ) | 6.33      | 6.33  | 6.28 | 6.28  | 6.25 | 6.26  | 6.48   | 5.92             | 6.95             | 5.96             | 6.04             | 6.08             |                   | 5.98              |
| <sup>1</sup> A <sub>1</sub> ( $\pi \rightarrow \pi^*$ ) | 6.43      | 6.40  | 6.35 | 6.32  | 6.32 | 6.30  | 6.53   | 5.92             | 6.59             | 6.53             | 6.37             | 6.15             |                   |                   |
| <sup>1</sup> B <sub>2</sub> ( $\pi \rightarrow 3p$ )    | 7.20      | 7.20  | 7.00 | 7.00  | 6.83 |       | 6.62   | 5.78             | 6.26             | 6.61             | 6.57             |                  |                   |                   |
| <sup>3</sup> B <sub>2</sub> ( $\pi \rightarrow \pi^*$ ) | 4.59      | 4.58  | 4.56 | 4.54  | 4.53 |       | 4.74   | 4.27             |                  | 4.53             |                  |                  | 4.21              |                   |
| <sup>3</sup> A <sub>2</sub> ( $\pi \rightarrow 3s$ )    | 5.22      | 5.22  | 5.12 | 5.12  | 5.21 |       | 5.49   | 5.04             |                  | 5.07             |                  |                  | 5.1               |                   |
| <sup>3</sup> A <sub>1</sub> ( $\pi \rightarrow \pi^*$ ) | 5.54      | 5.54  | 5.49 | 5.48  | 5.46 |       | 5.56   | 5.16             |                  | 5.53             |                  |                  |                   |                   |
| <sup>3</sup> B <sub>1</sub> ( $\pi \rightarrow 3p$ )    | 5.91      | 5.90  | 5.82 | 5.81  | 5.92 |       | 6.28   | 5.82             |                  | 5.74             |                  |                  |                   |                   |

<sup>a</sup>CASPT2 results from ref 32. <sup>b</sup>NEVPT2 results from ref 180. <sup>c</sup>MR-CC results from ref 182. <sup>d</sup>ADC(3) results from ref 184. <sup>e</sup>CC3 results from ref 27. <sup>f</sup>Various experiments summarized in ref 178. <sup>g</sup>Electron impact from ref 185: for the <sup>1</sup>A<sub>1</sub> state two values (6.44 and 6.61 eV) are reported, whereas for the two lowest triplet states, two values (3.99 and 5.22 eV) can be found in ref 186. <sup>h</sup>NEVPT2 results from ref 187. <sup>i</sup>Best estimate from ref 188, based on CC calculations. <sup>j</sup>XMS-CASPT2 results from ref 189. <sup>k</sup>Electron impact from refs 185 and 190. <sup>l</sup>Vapor UV spectra from refs 191, 192, and 193.

the vast majority of the cases. Importantly, we confirm our previous conclusions obtained on smaller compounds:<sup>28</sup> (i) CCSDTQ values systematically fall within (or are extremely close to) the FCI error bar, and (ii) both CC3 and CCSDT are also highly trustable when the considered ES does not exhibit a strong double excitation character. Indeed, considering the 54 “single” ES cases for which CCSDTQ estimates could be obtained (only excluding the lowest <sup>1</sup>A<sub>g</sub> ES of butadiene and glyoxal), we determined negligible MSE < 0.01 eV, tiny MAE (0.01 and 0.02 eV), and small maximal deviations (0.05 and 0.04 eV) for CC3 and CCSDT, respectively. This clearly indicates that these two approaches provide chemically accurate estimates (errors below 1 kcal·mol<sup>-1</sup> or 0.043 eV) for most electronic transitions. Interestingly, some of us have shown that CC3 also provides chemically accurate 0-0 energies as compared to experimental values for most valence transitions.<sup>116,117,173</sup> When comparing the NEVPT2 and CC3 (CCSDT) results obtained with aVTZ for the 91 (65) ESs for which comparisons are possible (again excluding only the lowest <sup>1</sup>A<sub>g</sub> states of butadiene and glyoxal), one obtains an MSE of +0.09 (+0.09) eV and an MAE of 0.11 (0.12) eV. This seems to indicate that NEVPT2, as applied here, has a slight tendency to overestimate the transition energies. This contrasts with “standard” CASPT2 which is known to generally underestimate transition energies, as further illustrated and discussed above and below.

**3.2. Five-Membered Rings.** We now consider five-membered rings, and, in particular, five common derivatives

that have been considered in several previous theoretical studies (*vide infra*): cyclopentadiene, furan, imidazole, pyrrole, and thiophene. As the most advanced levels of theory employed in the previous section, namely, CCSDTQ and FCI, become beyond reach for these compounds (except in very rare occasions), one has to rely on the nature of the ES and the consistency between results to deduce TBES.

For furan, *ab initio* calculations have been performed with almost every available wave function method.<sup>23,25–27,32,174–184</sup> However, the present work is, to the best of our knowledge, the first to disclose CCSDT values as well as CC3 energies obtained with a quadruple- $\zeta$  basis set. Our results for 10 low-lying ES states are listed in Tables 5 and S5. All singlet (triplet) transitions are characterized by %T<sub>1</sub> values in the 92–94% (97–99%) range. Consistently, the maximal discrepancy between CC3 and CCSDT is small (0.04 eV). In addition, there is a decent consistency between the present data and the NEVPT2 results of both ref 180 and of the present work, as well as the MR-CC values of ref 182. This holds for almost all transitions except the <sup>1</sup>B<sub>2</sub>( $\pi \rightarrow 3p$ ) excitation that we predict to be significantly higher than in most previous works, even after accounting for the quite large basis set effects (–0.10 eV between the aVTZ and aVQZ estimates, see Table S5). We believe that our estimate is the most accurate to date for this particularly tricky ES. Interestingly, the recent ADC(3) values of ref 184 are consistently smaller by ca. –0.2 eV as compared to CCSDT (see Table 6), in agreement with the error sign we observed in

Table 6. Vertical Transition Energies (in eV) of Cyclopentadiene, Imidazole, and Thiophene

| state   | Cyclopentadiene |       |      |       |      |       |        |                  |                  |                  |                  |                   | exp. <sup>f</sup> | exp. <sup>g</sup> |
|---|-----------------|-------|------|-------|------|-------|--------|------------------|------------------|------------------|------------------|-------------------|-------------------|-------------------|
|   | 6-31+G(d)       |       | aVDZ |       | aVTZ |       |        | lit.             |                  |                  |                  |                   |                   |                   |
|   | CC3             | CCSDT | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 | th. <sup>a</sup> | th. <sup>b</sup> | th. <sup>c</sup> | th. <sup>d</sup> | exp. <sup>e</sup> |                   |                   |
| <sup>1</sup> B <sub>2</sub> ( $\pi \rightarrow \pi^*$ ) | 5.79            | 5.80  | 5.59 | 5.60  | 5.54 | 5.56  | 5.65   | 5.27             | 5.54             | 5.19             | 5.58             | 5.26              |                   | 5.20              |
| <sup>1</sup> A <sub>2</sub> ( $\pi \rightarrow 3s$ )    | 6.08            | 6.08  | 5.70 | 5.70  | 5.77 | 5.78  | 5.92   | 5.65             | 5.58             | 5.62             | 5.79             | 5.68              | 5.63              |                   |
| <sup>1</sup> B <sub>1</sub> ( $\pi \rightarrow 3p$ )    | 6.57            | 6.58  | 6.34 | 6.34  | 6.40 | 6.41  | 6.42   | 6.24             | 6.17             | 6.24             | 6.43             |                   | 6.35              |                   |
| <sup>1</sup> A <sub>2</sub> ( $\pi \rightarrow 3p$ )    | 6.67            | 6.67  | 6.39 | 6.39  | 6.45 | 6.46  | 6.59   | 6.30             | 6.21             | 6.25             | 6.47             |                   |                   | 6.26              |
| <sup>1</sup> B <sub>2</sub> ( $\pi \rightarrow 3p$ )    | 7.06            | 7.07  | 6.55 | 6.55  | 6.56 | 6.56  | 6.60   | 6.25             | 6.22             | 6.27             | 6.58             |                   | 6.31              |                   |
| <sup>1</sup> A <sub>1</sub> ( $\pi \rightarrow \pi^*$ ) | 6.67            | 6.60  | 6.59 | 6.53  | 6.57 | 6.52  | 6.75   | 6.31             | 6.76             | 6.42             | 6.65             |                   |                   | ~6.2              |
| <sup>3</sup> B <sub>2</sub> ( $\pi \rightarrow \pi^*$ ) | 3.33            | 3.33  | 3.32 | 3.31  | 3.32 |       | 3.41   | 3.15             | 3.40             |                  |                  | 3.10              |                   |                   |
| <sup>3</sup> A <sub>1</sub> ( $\pi \rightarrow \pi^*$ ) | 5.16            | 5.15  | 5.14 | 5.13  | 5.12 |       | 5.30   | 4.90             | 5.18             |                  |                  | >4.7              |                   |                   |
| <sup>3</sup> A <sub>2</sub> ( $\pi \rightarrow 3s$ )    | 6.01            | 6.02  | 5.65 | 5.65  | 5.73 |       | 5.73   | 5.63             | 5.56             |                  |                  |                   |                   |                   |
| <sup>3</sup> B <sub>1</sub> ( $\pi \rightarrow 3p$ )    | 6.51            | 6.52  | 6.30 | 6.30  | 6.36 |       | 6.40   | 6.25             | 6.19             |                  |                  |                   |                   |                   |

| state   | Imidazole |       |      |       |      |        |                  |                  |                   |      | exp. <sup>j</sup> |
|---|-----------|-------|------|-------|------|--------|------------------|------------------|-------------------|------|-------------------|
|   | 6-31+G(d) |       | aVDZ |       | aVTZ |        |                  | lit.             |                   |      |                   |
|   | CC3       | CCSDT | CC3  | CCSDT | CC3  | NEVPT2 | th. <sup>h</sup> | th. <sup>i</sup> | exp. <sup>k</sup> |      |                   |
| <sup>1</sup> A''( $\pi \rightarrow 3s$ )                | 5.77      | 5.77  | 5.60 | 5.60  | 5.71 | 5.93   | 5.71             |                  |                   | ~5.2 |                   |
| <sup>1</sup> A'( $\pi \rightarrow \pi^*$ ) <sup>k</sup> | 6.51      | 6.51  | 6.43 | 6.43  | 6.41 | 6.73   | 6.72             | 6.25             |                   | ~6.4 |                   |
| <sup>1</sup> A''( $n \rightarrow \pi^*$ )               | 6.66      | 6.66  | 6.42 | 6.42  | 6.50 | 6.96   | 6.52             | 6.65             |                   |      |                   |
| <sup>1</sup> A'( $\pi \rightarrow 3p$ ) <sup>k</sup>    | 7.04      | 7.02  | 6.93 | 6.89  | 6.87 | 7.00   | 6.49             |                  |                   |      |                   |
| <sup>3</sup> A'( $\pi \rightarrow \pi^*$ )              | 4.83      | 4.81  | 4.78 |       | 4.75 | 4.86   | 4.49             | 4.65             |                   |      |                   |
| <sup>3</sup> A''( $\pi \rightarrow 3s$ )                | 5.72      | 5.72  | 5.57 | 5.56  | 5.67 | 5.91   | 5.68             |                  |                   |      |                   |
| <sup>3</sup> A'( $\pi \rightarrow \pi^*$ )              | 5.88      | 5.88  | 5.78 |       | 5.74 | 5.91   | 5.47             | 5.64             |                   |      |                   |
| <sup>3</sup> A''( $n \rightarrow \pi^*$ )               | 6.48      | 6.46  | 6.37 | 6.35  | 6.33 | 6.48   | 6.07             | 6.25             |                   |      |                   |

| state   | Thiophene |       |      |       |      |       |        |                  |                  |                  |                  |                   | exp. <sup>p</sup> | exp. <sup>q</sup> | exp. <sup>r</sup> |
|---|-----------|-------|------|-------|------|-------|--------|------------------|------------------|------------------|------------------|-------------------|-------------------|-------------------|-------------------|
|   | 6-31+G(d) |       | aVDZ |       | aVTZ |       |        | lit.             |                  |                  |                  |                   |                   |                   |                   |
|   | CC3       | CCSDT | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 | th. <sup>l</sup> | th. <sup>m</sup> | th. <sup>n</sup> | th. <sup>o</sup> | exp. <sup>p</sup> |                   |                   |                   |
| <sup>1</sup> A <sub>1</sub> ( $\pi \rightarrow \pi^*$ )           | 5.79      | 5.77  | 5.70 | 5.68  | 5.65 | 5.64  | 5.84   | 5.33             | 5.41             | 5.70             | 5.64             | 5.16              | 5.13              | 5.16              |                   |
| <sup>1</sup> B <sub>2</sub> ( $\pi \rightarrow \pi^*$ )           | 6.23      | 6.24  | 6.05 | 6.06  | 5.96 | 5.98  | 6.10   | 5.72             | 5.72             | 6.10             | 5.97             | 5.99              | 5.83              |                   |                   |
| <sup>1</sup> A <sub>2</sub> ( $\pi \rightarrow 3s$ )              | 6.26      | 6.26  | 6.07 | 6.06  | 6.14 | 6.14  | 6.20   | 5.93             | 5.70             | 6.05             | 6.23             |                   |                   |                   |                   |
| <sup>1</sup> B <sub>1</sub> ( $\pi \rightarrow 3p$ )              | 6.18      | 6.17  | 6.19 | 6.17  | 6.14 | 6.14  | 6.19   | 6.30             | 5.87             | 6.30             | 6.17             |                   |                   | 6.71              |                   |
| <sup>1</sup> A <sub>2</sub> ( $\pi \rightarrow 3p$ )              | 6.32      |       | 6.33 | 6.31  | 6.25 | 6.21  | 6.40   | 6.35             | 6.03             | 6.28             | 6.33             |                   |                   |                   |                   |
| <sup>1</sup> B <sub>1</sub> ( $\pi \rightarrow 3s$ )              | 6.62      | 6.62  | 6.42 | 6.41  | 6.50 | 6.49  | 6.71   | 6.23             | 6.12             | 6.36             | 6.68             |                   |                   | 6.47              |                   |
| <sup>1</sup> B <sub>2</sub> ( $\pi \rightarrow 3p$ ) <sup>s</sup> | 7.45      | 7.44  | 7.45 | 7.44  | 7.29 | 7.29  | 7.25   | 6.56             | 6.41             | 6.81             | 6.97             |                   |                   | 6.60              |                   |
| <sup>1</sup> A <sub>1</sub> ( $\pi \rightarrow \pi^*$ )           | 7.50      | 7.46  | 7.41 | 7.35  |      | 7.39  | 6.69   | 7.32             | 7.71             | 7.74             | 6.61             |                   |                   |                   |                   |
| <sup>3</sup> B <sub>2</sub> ( $\pi \rightarrow \pi^*$ )           | 3.95      | 3.94  | 3.96 | 3.94  | 3.94 |       | 4.13   | 3.75             | 3.94             |                  | 3.96             |                   | 3.74              |                   |                   |
| <sup>3</sup> A <sub>1</sub> ( $\pi \rightarrow \pi^*$ )           | 4.90      | 4.90  | 4.82 | 4.81  | 4.77 |       | 4.84   | 4.50             | 4.86             |                  | 4.87             |                   | 4.62              |                   |                   |
| <sup>3</sup> B <sub>1</sub> ( $\pi \rightarrow 3p$ )              | 6.00      | 5.98  | 6.01 | 5.99  | 5.95 |       | 5.98   | 5.90             | 5.94             |                  | 6.01             |                   |                   |                   |                   |
| <sup>3</sup> A <sub>2</sub> ( $\pi \rightarrow 3s$ )              | 6.20      | 6.20  | 6.01 | 6.00  | 6.09 |       | 6.14   | 5.88             | 5.75             |                  | 5.83             |                   |                   |                   |                   |

<sup>a</sup>CASPT2 results from ref 32. <sup>b</sup>SAC-CI results from ref 198. <sup>c</sup>MR-MP results from ref 174. <sup>d</sup>CC3 results from ref 27. <sup>e</sup>Electron impact from ref 199. <sup>f</sup>Gas phase absorption from ref 200. <sup>g</sup>Energy loss from ref 201 for the two valence states; two-photon resonant experiment from ref 202 for the <sup>1</sup>A<sub>2</sub> Rydberg ES. <sup>h</sup>CASPT2 results from ref 38. <sup>i</sup>CC3 results from ref 26. <sup>j</sup>Gas-phase experimental estimates from ref 203. <sup>k</sup>The assignments of these two states as valence and Rydberg are based on the oscillator strength, but both have a partial Rydberg character. The CASSCF spatial extent is in fact larger for the lowest transition, and Roos consequently classified both ESs as Rydberg transitions.<sup>38</sup> <sup>l</sup>CASPT2 results from ref 33. <sup>m</sup>SAC-CI results from ref 204. <sup>n</sup>CCSDR(3) results from ref 205. <sup>o</sup>TBE from ref 206, based on EOM-CCSD for the singlet and ADC(2) for triplets. <sup>p</sup>0-0 energies from ref 207. <sup>q</sup>0-0 energies from ref 208 for the singlets and energy loss experiment from ref 209 for the triplets. <sup>r</sup>0-0 energies from ref 206. <sup>s</sup>Non-negligible mixing with a close-lying ( $\pi \rightarrow \pi^*$ ) transition of the same symmetry.

smaller compounds for ADC(3).<sup>28</sup> Again, we note that the experimental data<sup>156,185,186</sup> provide the same state ordering as our calculations.

Like furan, pyrrole has been extensively investigated in the literature using a large panel of theoretical methods.<sup>23,25–27,32,174,178,181–183,187–189,191,194–197</sup> We report six low-lying singlet and four triplet ESs in Tables 5 and S5. All these transitions have very large %*T*<sub>1</sub> values except for the totally symmetric  $\pi \rightarrow \pi^*$  excitation (%*T*<sub>1</sub> = 86%). For each state, we found highly consistent CC3 and CCSDT results, often

significantly larger than older multireference estimates<sup>32,182,196</sup> but in nice agreement with the very recent XMS-CASPT2 results of the González group,<sup>189</sup> and the present NEVPT2 estimates [at the exception of the <sup>1</sup>A<sub>2</sub>( $\pi \rightarrow 3p$ ) transition]. The match obtained with the 20 years old extrapolated CC values of Christiansen and co-workers<sup>188</sup> is quite remarkable. The only exceptions are the two B<sub>2</sub> transitions that were reported as significantly mixed in this venerable work. For the lowest singlet ES, the FCI/6-31+G(d) value is 5.24 ± 0.02 eV confirming the performance of both CC3 and CCSDT for this transition. As can

Table 7. Vertical Transition Energies (in eV) of Benzene

| state  | 6-31+G(d) |       | aVDZ |       | aVTZ |       |        | lit.             |                  |                  |                  |                   |                   |
|--|-----------|-------|------|-------|------|-------|--------|------------------|------------------|------------------|------------------|-------------------|-------------------|
|  | CC3       | CCSDT | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 | th. <sup>a</sup> | th. <sup>b</sup> | th. <sup>c</sup> | th. <sup>d</sup> | exp. <sup>e</sup> | exp. <sup>f</sup> |
| <sup>1</sup> B <sub>2u</sub> ( $\pi \rightarrow \pi^*$ ) | 5.13      | 5.10  | 5.11 | 5.08  | 5.09 | 5.06  | 5.32   | 4.84             | 5.08             | 5.06             | 5.03             |                   | 4.90              |
| <sup>1</sup> B <sub>1u</sub> ( $\pi \rightarrow \pi^*$ ) | 6.68      | 6.69  | 6.50 | 6.50  | 6.44 | 6.45  | 6.43   | 6.30             | 6.54             | 6.22             | 6.23             |                   | 6.20              |
| <sup>1</sup> E <sub>1g</sub> ( $\pi \rightarrow 3s$ )    | 6.75      | 6.76  | 6.46 | 6.46  | 6.52 | 6.52  | 6.75   | 6.38             | 6.51             | 6.42             |                  |                   | 6.33              |
| <sup>1</sup> A <sub>2u</sub> ( $\pi \rightarrow 3p$ )    | 7.24      | 7.25  | 7.02 | 7.02  | 7.08 | 7.08  | 7.40   | 6.86             | 6.97             | 7.06             |                  |                   | 6.93              |
| <sup>1</sup> E <sub>2u</sub> ( $\pi \rightarrow 3p$ )    | 7.34      | 7.35  | 7.09 | 7.09  | 7.15 | 7.15  | 7.45   | 6.91             | 7.03             | 7.12             |                  |                   | 6.95              |
| <sup>3</sup> B <sub>1u</sub> ( $\pi \rightarrow \pi^*$ ) | 4.18      | 4.16  | 4.19 | 4.17  | 4.18 |       | 4.32   | 3.89             | 4.15             | 3.88             | 4.11             | 3.95              |                   |
| <sup>3</sup> E <sub>1u</sub> ( $\pi \rightarrow \pi^*$ ) | 4.95      | 4.94  | 4.89 | 4.88  | 4.86 |       | 4.92   | 4.49             | 4.86             | 4.72             | 4.75             | 4.75              |                   |
| <sup>3</sup> B <sub>2u</sub> ( $\pi \rightarrow \pi^*$ ) | 6.06      | 6.06  | 5.86 | 5.86  | 5.81 |       | 5.51   | 5.49             | 5.88             | 5.54             | 5.67             | 5.60              |                   |

<sup>a</sup>CASPT2 results from ref 34. <sup>b</sup>CC3 results from ref 215. <sup>c</sup>SAC-CI results from ref 218. <sup>d</sup>RASPT2(18,18) results from ref 220. <sup>e</sup>Electron impact from ref 221. <sup>f</sup>Jet-cooled experiment from ref 225 for the two lowest states, and multiphoton experiments from refs 223 and 224 for the Rydberg states.

be seen in Table S5, aVTZ yields basis-set converged transition energies, except, like in furan, for the Rydberg <sup>1</sup>B<sub>2</sub>( $\pi \rightarrow 3p$ ) transition that is significantly red-shifted (−0.09 eV) when pushing to the quadruple- $\zeta$  basis set. As mentioned in Thiel's work,<sup>23</sup> the experimental spectra of pyrrole are quite broad, and the rare available experiments<sup>185,190–193,195</sup> can only be considered as general guidelines.

Although diverse wave function studies have been performed on cyclopentadiene (including CASPT2,<sup>23,26,32</sup> CC,<sup>23,25,27</sup> SAC-CI,<sup>198</sup> and various multireference approaches<sup>174,181</sup>), this compound has received less attention than other members of the five-membered ring family, namely, furan and pyrrole (*vide infra*). This is probably due to the presence of the methylene group that renders computations significantly more expensive. Most transitions listed in Tables 6 and S6 are characterized by %  $T_1$  exceeding 93%, the only exception being the <sup>1</sup>A<sub>1</sub>( $\pi \rightarrow \pi^*$ ) excitation that has a similar nature as the lowest A<sub>g</sub> state of butadiene (% $T_1$  = 79%). Consistently, the CC3 and CCSDT results are nearly identical for all ESs except for the <sup>1</sup>A<sub>1</sub> ES. By comparing the results obtained for this <sup>1</sup>A<sub>1</sub>( $\pi \rightarrow \pi^*$ ) transition to its butadiene counterpart, one can infer that the CCSDT estimate is probably too large by roughly 0.04–0.08 eV and that the NEVPT2 value is unlikely to be accurate enough to establish a definitive TBE. This statement is also in line with the results of ref 72. For the two B<sub>2</sub>( $\pi \rightarrow \pi^*$ ) transitions, we could obtain FCI/6-31+G(d) estimates of 5.78 ± 0.02 eV (singlet) and 3.33 ± 0.05 eV (triplet), the CC3 and CCSDT transition energies falling inside these energetic windows in both cases. As one can see in Tables 6 and S6, the basis set effects are rather moderate for the electronic transitions of cyclopentadiene, with no variation larger than 0.10 eV (0.02 eV) between aVDZ and aVTZ (aVTZ and aVQZ). When comparing to literature data, our values are unsurprisingly consistent with the CC3 values of Schwabe and Goerigk<sup>27</sup> and tend to be significantly larger than earlier CASPT2<sup>26,32</sup> and MR-MP<sup>174</sup> estimates. As expected, a few gas-phase experiments are available as well for this derivative,<sup>199–202</sup> but they hardly represent grounds for comparison.

Due to its lower symmetry, imidazole has been less investigated, the most advanced studies available probably remain the CASPT2 work of Serrano-Andr s and co-workers from 1996,<sup>38</sup> and the basis-set extrapolated CC3 results of Silva-Junior et al. for the valence transitions from 2010.<sup>26</sup> The experimental data in gas-phase are also limited.<sup>203</sup> Our results are displayed in Tables 6 and S6. The CC3 and CCSDT values are quite consistent despite the fact that the % $T_1$  values of the two singlet A' states are slightly smaller than 90%. These two

states have indeed, at least partially, a Rydberg character (see the footnote in Table 6). The agreement between the CC estimates and previous CASPT2,<sup>38</sup> and current NEVPT2 energies is reasonable, the latter being systematically larger than their CC3 counterparts. For the eight transitions considered here, the basis set effects are moderate and aVTZ yield results within 0.03 eV of their aVQZ counterparts (Table S6 in the SI).

Finally, the ESs of thiophene, which is one of the most important building block in organic electronic devices, were the subject of previous theoretical investigations<sup>33,204–206,210,211</sup> that unveiled a series of transitions that were not yet characterized in the available measurements.<sup>186,190,206–210</sup> To the best of our knowledge, the present work is the first to report CC calculations obtained with (iterative) triples and therefore constitutes the most accurate estimates to date. Indeed, all the transitions listed in Tables 6 and S6 are characterized by a largely dominant single excitation character, with % $T_1$  above 90% except for the two <sup>1</sup>A<sub>1</sub> transitions for which % $T_1$  = 88% and 87%. The agreement between CC3 and CCSDT remains nevertheless excellent for these low-lying totally symmetric transitions. Thiophene is also one of these compounds for which the unambiguous characterization of the nature of the ES is difficult, with, e.g., a strong mixing between the second and third singlet ESs of B<sub>2</sub> symmetry. This makes the assignment of the valence ( $\pi \rightarrow \pi^*$ ) or Rydberg ( $\pi \rightarrow 3p$ ) character of this transition particularly tricky at the CC3 level. We note that contradictory assignments can be found in the literature.<sup>33,204,205</sup> As for the previously discussed isostructural systems, the only ES that undergoes significant basis set effects beyond aVTZ is the Rydberg <sup>1</sup>B<sub>2</sub>( $\pi \rightarrow 3p$ ) (−0.09 eV when upgrading to aVQZ, see Table S6) and that the NEVPT2 estimates tend to be slightly larger than the CC3 values. The data of Table 6 are globally in good agreement with the previously reported values with discrepancies that are significant only for the three highest-lying singlet states.

**3.3. Six-Membered Rings.** Let us now turn toward seven six-membered rings which play a key role in chemistry: benzene, pyrazine, pyridazine, pyridine, pyrimidine, tetrazine, and triazine. To the best of our knowledge, the present work is the first to propose CCSDT reference energies as well as CC3/aVQZ values for all these compounds. Of course, these systems have been investigated before, and beyond Thiel's benchmarks,<sup>23,25,26</sup> it is worth pointing out the early investigation of Del Bene and co-workers<sup>212</sup> performed with a CC approach including perturbative corrections for the triples. Following a theoretically consistent protocol, Nooijen<sup>213</sup> also performed STEOM-CCSD calculations to study the ESs of each of these

Table 8. Vertical Transition Energies (in eV) of Pyrazine and Tetrazine

| state  | Pyrazine  |       |      |       |      |       |        |                  |                  |                  |                  |                   |                   |
|--|-----------|-------|------|-------|------|-------|--------|------------------|------------------|------------------|------------------|-------------------|-------------------|
|  | 6-31+G(d) |       | aVDZ |       | aVTZ |       |        | lit.             |                  |                  |                  |                   |                   |
|  | CC3       | CCSDT | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 | th. <sup>a</sup> | th. <sup>b</sup> | th. <sup>c</sup> | th. <sup>d</sup> | exp. <sup>e</sup> | exp. <sup>f</sup> |
| <sup>1</sup> B <sub>3u</sub> (n → π*)              | 4.28      | 4.28  | 4.19 | 4.19  | 4.14 | 4.15  | 4.17   | 3.83             | 4.12             | 4.25             | 4.19             | 3.93              |                   |
| <sup>1</sup> A <sub>u</sub> (n → π*)               | 5.08      | 5.08  | 4.98 | 4.98  | 4.97 | 4.98  | 4.77   | 4.36             | 4.93             | 5.24             | 4.93             |                   |                   |
| <sup>1</sup> B <sub>2u</sub> (π → π*)              | 5.10      | 5.08  | 5.07 | 5.05  | 5.03 | 5.02  | 5.32   | 4.79             | 4.75             | 4.84             | 5.19             | 4.8               | 4.81              |
| <sup>1</sup> B <sub>2g</sub> (n → π*)              | 5.86      | 5.85  | 5.78 | 5.77  | 5.71 | 5.71  | 5.88   | 5.50             | 5.85             | 6.04             | 5.81             | 5.19              |                   |
| <sup>1</sup> A <sub>g</sub> (n → 3s)               | 6.74      | 6.73  | 6.54 | 6.53  | 6.66 | 6.65  | 6.70   |                  | 6.83             | 7.07             | 6.46             |                   |                   |
| <sup>1</sup> B <sub>1g</sub> (n → π*)              | 6.87      | 6.87  | 6.75 | 6.75  | 6.73 | 6.74  | 6.75   | 6.26             | 6.73             |                  | 6.73             |                   | 6.10              |
| <sup>1</sup> B <sub>1u</sub> (π → π*)              | 7.10      | 7.11  | 6.92 | 6.93  | 6.86 | 6.88  | 6.81   | 6.60             | 6.89             | 6.68             | 6.99             | 6.5               | 6.51              |
| <sup>1</sup> B <sub>1g</sub> (π → 3s)              | 7.36      | 7.37  | 7.13 | 7.14  | 7.20 | 7.21  | 7.33   |                  | 7.31             | 7.08             |                  |                   |                   |
| <sup>1</sup> B <sub>2u</sub> (n → 3p)              | 7.39      | 7.39  | 7.14 | 7.13  | 7.25 |       | 7.25   |                  | 7.45             | 7.67             | 7.06             |                   |                   |
| <sup>1</sup> B <sub>1u</sub> (n → 3p)              | 7.56      | 7.55  | 7.38 | 7.37  | 7.45 |       | 7.42   | 7.28             | 7.50             | 7.73             | 7.31             |                   | 7.67              |
| <sup>1</sup> B <sub>1u</sub> (π → π*)              | 8.19      | 8.23  | 7.99 | 8.03  | 7.94 |       | 8.25   | 7.43             | 7.96             | 8.24             | 8.08             |                   |                   |
| <sup>3</sup> B <sub>3u</sub> (n → π*)              | 3.68      | 3.68  | 3.60 | 3.60  | 3.59 |       | 3.56   | 3.16             |                  |                  |                  | 3.33              |                   |
| <sup>3</sup> B <sub>1u</sub> (π → π*)              | 4.39      | 4.36  | 4.40 | 4.36  | 4.39 |       | 4.57   | 4.15             |                  |                  |                  | 4.04              |                   |
| <sup>3</sup> B <sub>2u</sub> (π → π*)              | 4.56      | 4.55  | 4.46 | 4.45  | 4.40 |       | 4.42   | 4.28             |                  |                  |                  | ~4.4              |                   |
| <sup>3</sup> A <sub>u</sub> (n → π*)               | 5.05      | 5.05  | 4.93 | 4.93  | 4.93 |       | 4.75   | 4.19             |                  |                  |                  | 4.2               |                   |
| <sup>3</sup> B <sub>2g</sub> (n → π*)              | 5.18      | 5.17  | 5.11 | 5.11  | 5.08 |       | 5.21   | 4.81             |                  |                  |                  | 4.49              |                   |
| <sup>3</sup> B <sub>1u</sub> (π → π*)              | 5.38      | 5.37  | 5.32 | 5.31  | 5.29 |       | 5.35   | 4.98             |                  |                  |                  |                   |                   |
| state  | Tetrazine |       |      |       |      |       |        |                  |                  |                  |                  |                   |                   |
|  | 6-31+G(d) |       | aVDZ |       | aVTZ |       |        | lit.             |                  |                  |                  |                   |                   |
|  | CC3       | CCSDT | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 | th. <sup>g</sup> | th. <sup>h</sup> | th. <sup>i</sup> | th. <sup>j</sup> | th. <sup>k</sup>  | exp. <sup>l</sup> |
| <sup>1</sup> B <sub>3u</sub> (n → π*)              | 2.53      | 2.54  | 2.49 | 2.50  | 2.46 | 2.47  | 2.35   | 1.96             | 2.22             | 2.01             | 2.29             | 2.46              | 2.35              |
| <sup>1</sup> A <sub>u</sub> (n → π*)               | 3.75      | 3.75  | 3.69 | 3.70  | 3.67 | 3.69  | 3.58   | 3.06             | 3.62             | 3.09             | 3.41             | 3.78              | 3.6               |
| <sup>1</sup> A <sub>g</sub> (double) <sup>m</sup>  | 6.22      | 5.86  | 6.22 | 5.86  | 6.21 | 5.96  | 4.61   | 4.37             | 5.06             | 4.34             | 4.66             |                   |                   |
| <sup>1</sup> B <sub>1g</sub> (n → π*)              | 5.01      | 5.02  | 4.97 | 4.98  | 4.91 | 4.93  | 4.95   | 4.51             | 4.73             | 4.47             | 4.53             | 4.87              |                   |
| <sup>1</sup> B <sub>2u</sub> (π → π*)              | 5.29      | 5.26  | 5.27 | 5.25  | 5.23 | 5.21  | 5.56   | 4.89             | 4.90             |                  | 5.59             | 5.08              | 4.97              |
| <sup>1</sup> B <sub>2g</sub> (n → π*)              | 5.56      | 5.52  | 5.53 | 5.50  | 5.46 | 5.45  | 5.63   | 5.05             | 5.09             | 4.92             | 5.59             | 5.28              |                   |
| <sup>1</sup> A <sub>u</sub> (n → π*)               | 5.61      | 5.61  | 5.59 | 5.59  | 5.52 | 5.53  | 5.62   | 5.28             | 5.23             | 5.32             | 5.95             | 5.39              | 5.5               |
| <sup>1</sup> B <sub>3g</sub> (double) <sup>m</sup> | 7.64      |       | 7.62 |       | 7.62 |       | 6.15   | 5.16             | 6.30             | 5.26             | 6.01             |                   | 5.92              |
| <sup>1</sup> B <sub>2g</sub> (n → π*)              | 6.24      | 6.22  | 6.17 | 6.16  | 6.13 |       | 6.13   | 5.48             | 6.16             | 5.78             | 6.05             | 6.16              |                   |
| <sup>1</sup> B <sub>1g</sub> (n → π*)              | 7.04      | 7.04  | 6.98 | 6.98  | 6.92 |       | 6.76   | 5.99             | 6.73             | 6.20             | 6.92             | 6.80              |                   |
| <sup>3</sup> B <sub>3u</sub> (n → π*)              | 1.87      | 1.88  | 1.86 | 1.86  | 1.85 |       | 1.73   | 1.45             | 1.71             |                  |                  | 1.87              | 1.7               |
| <sup>3</sup> A <sub>u</sub> (n → π*)               | 3.48      | 3.49  | 3.43 | 3.44  | 3.44 |       | 3.36   | 2.81             | 3.47             |                  |                  | 3.49              | 2.90              |
| <sup>3</sup> B <sub>1g</sub> (n → π*)              | 4.25      | 4.25  | 4.23 | 4.23  | 4.20 |       | 4.24   | 3.76             | 3.97             |                  |                  | 4.18              |                   |
| <sup>3</sup> B <sub>1u</sub> (π → π*)              | 4.54      | 4.49  | 4.54 | 4.49  | 4.54 |       | 4.70   | 4.25             | 3.67             |                  |                  | 4.36              |                   |
| <sup>3</sup> B <sub>2u</sub> (π → π*)              | 4.65      | 4.64  | 4.58 | 4.58  | 4.52 |       | 4.58   | 4.29             | 4.35             |                  |                  | 4.39              |                   |
| <sup>3</sup> B <sub>2g</sub> (n → π*)              | 5.11      | 5.11  | 5.09 | 5.08  | 5.05 |       | 5.27   | 4.67             | 4.78             |                  |                  | 4.89              |                   |
| <sup>3</sup> A <sub>u</sub> (n → π*)               | 5.17      | 5.17  | 5.15 | 5.15  | 5.11 |       | 5.13   | 4.85             | 4.89             |                  |                  | 4.96              |                   |
| <sup>3</sup> B <sub>3g</sub> (double) <sup>m</sup> | 7.35      |       | 7.33 |       | 7.35 |       | 5.51   | 5.08             |                  |                  |                  |                   |                   |
| <sup>3</sup> B <sub>1u</sub> (π → π*)              | 5.51      | 5.50  | 5.46 | 5.46  | 5.42 |       | 5.56   | 5.09             | 5.31             |                  |                  | 5.32              |                   |

<sup>a</sup>CASPT2 results from ref 226. <sup>b</sup>STEOM-CCSD results from ref 213. <sup>c</sup>SAC-CI results from ref 218. <sup>d</sup>CC3 results from ref 27. <sup>e</sup>Double resonance dip spectroscopy from ref 235 (B<sub>3u</sub> and B<sub>2g</sub> ESs) and EEL from ref 236 (others). <sup>f</sup>UV max from ref 234. <sup>g</sup>CASPT2 results from ref 229. <sup>h</sup>Ext-STEOM-CCSD results from ref 231. <sup>i</sup>GVPPT2 results from ref 219. <sup>j</sup>NEVPT2 results from ref 232. <sup>k</sup>CC3 results from ref 26. <sup>l</sup>From ref 238, the singlets are from EEL, except for the 4.97 and 5.92 eV values that are from VUV; the triplets are from EEL, and additional (unassigned) triplet peaks are found at 4.21, 4.6, and 5.2 eV. <sup>m</sup>All these three doubly ESs have a (n, n → π\*, π\*) character.

derivatives. However, these two works only considered singlet ESs.

**3.3.1. Benzene, Pyrazine, and Tetrazine.** These three highly symmetric systems allow CCSDT/aVTZ calculations for singlet states to be directly performed without the need of basis set extrapolations. Benzene was studied many times before,<sup>23,25–27,34,66,69,72,120,136,183,212–220</sup> and we report in Tables 7 and S7 estimates obtained for five singlet and three triplet ESs, all characterized by %T<sub>1</sub> exceeding 90% except for the lowest singlet (86%). As one can see, the two CC approaches are again

yielding very consistent transition energies with variations in the 0.00–0.03 eV range. Besides, aVTZ essentially provides basis set converged transition energies (Table S7). The present CC estimates are also very consistent with earlier CC3 results<sup>215</sup> and are compatible with both the very recent RASPT2<sup>220</sup> and our NEVPT2 values. For the states of both spin symmetries, the CC3 and CCSDT transition energies are slightly larger than the available electron impact/multiphoton measurements<sup>221–225</sup> but do provide energetic gaps between ESs very similar to the measured ones.

Table 9. Vertical Transition Energies (in eV) of Pyridazine and Pyridine

| state   | Pyridazine |       |      |       |      |        |                  |                  |                  |                  |                   |                   |
|---|------------|-------|------|-------|------|--------|------------------|------------------|------------------|------------------|-------------------|-------------------|
|   | 6-31+G(d)  |       | aVDZ |       | aVTZ |        | lit.             |                  |                  |                  |                   |                   |
|   | CC3        | CCSDT | CC3  | CCSDT | CC3  | NEVPT2 | th. <sup>a</sup> | th. <sup>b</sup> | th. <sup>c</sup> | th. <sup>d</sup> | exp. <sup>e</sup> | exp. <sup>f</sup> |
| <sup>1</sup> B <sub>1</sub> (n → π*)              | 3.95       | 3.95  | 3.86 | 3.86  | 3.83 | 3.80   | 3.48             | 3.76             | 3.65             | 3.85             |                   | 3.36              |
| <sup>1</sup> A <sub>2</sub> (n → π*)              | 4.49       | 4.48  | 4.39 | 4.39  | 4.37 | 4.40   | 3.66             | 4.46             | 4.28             | 4.44             |                   | 4.02              |
| <sup>1</sup> A <sub>1</sub> (π → π*)              | 5.36       | 5.32  | 5.33 | 5.30  | 5.29 | 5.58   | 4.86             | 4.92             | 4.86             | 5.20             | 5.0               | 5.01              |
| <sup>1</sup> A <sub>2</sub> (n → π*)              | 5.88       | 5.86  | 5.80 | 5.78  | 5.74 | 5.88   | 5.09             | 5.66             | 5.52             | 5.66             |                   | 5.61              |
| <sup>1</sup> B <sub>2</sub> (n → 3s)              | 6.26       | 6.27  | 6.06 | 6.06  | 6.17 | 6.21   |                  | 6.45             |                  |                  |                   |                   |
| <sup>1</sup> B <sub>1</sub> (n → π*)              | 6.51       | 6.51  | 6.41 | 6.41  | 6.37 | 6.64   | 5.80             | 6.41             | 6.20             | 6.33             |                   | 6.00              |
| <sup>1</sup> B <sub>2</sub> (π → π*)              | 6.96       | 6.97  | 6.79 | 6.80  | 6.74 | 7.10   | 6.61             | 6.77             | 6.44             | 6.68             |                   | 6.50              |
| <sup>3</sup> B <sub>1</sub> (n → π*)              | 3.27       | 3.26  | 3.20 | 3.20  | 3.19 | 3.13   |                  |                  |                  |                  |                   | 3.06              |
| <sup>3</sup> A <sub>2</sub> (n → π*)              | 4.19       | 4.19  | 4.11 | 4.11  | 4.11 | 4.14   |                  |                  |                  |                  |                   | 3.55              |
| <sup>3</sup> B <sub>2</sub> (π → π*)              | 4.39       | 4.36  | 4.39 | 4.35  | 4.38 | 4.49   |                  |                  |                  |                  | 4.0               | 4.33              |
| <sup>3</sup> A <sub>1</sub> (π → π*)              | 4.93       | 4.94  | 4.87 | 4.86  | 4.83 | 4.94   |                  |                  |                  |                  | 4.4               | 4.68              |
| state   | Pyridine   |       |      |       |      |        |                  |                  |                  |                  |                   |                   |
|   | 6-31+G(d)  |       | aVDZ |       | aVTZ |        | lit.             |                  |                  |                  |                   |                   |
|   | CC3        | CCSDT | CC3  | CCSDT | CC3  | NEVPT2 | th. <sup>g</sup> | th. <sup>b</sup> | th. <sup>c</sup> | th. <sup>d</sup> | exp. <sup>h</sup> | exp. <sup>i</sup> |
| <sup>1</sup> B <sub>1</sub> (n → π*)              | 5.12       | 5.10  | 5.01 | 5.00  | 4.96 | 5.15   | 4.91             | 4.90             | 4.80             | 4.95             | 5.24              | 4.78              |
| <sup>1</sup> B <sub>2</sub> (π → π*)              | 5.23       | 5.20  | 5.21 | 5.18  | 5.17 | 5.31   | 4.84             | 4.82             | 4.81             | 5.12             | 4.99              | 4.99              |
| <sup>1</sup> A <sub>2</sub> (n → π*)              | 5.55       | 5.54  | 5.41 | 5.41  | 5.40 | 5.29   | 5.17             | 5.31             | 5.24             | 5.41             | 5.43              | 5.40              |
| <sup>1</sup> A <sub>1</sub> (π → π*)              | 6.84       | 6.84  | 6.64 | 6.63  | 6.63 | 6.69   | 6.42             | 6.62             | 6.36             | 6.60             | 6.38              |                   |
| <sup>1</sup> A <sub>1</sub> (n → 3s)              | 6.92       | 6.92  | 6.71 | 6.71  | 6.76 | 6.99   | 6.70             | 6.96             | 6.64             |                  | 6.28              | 6.25              |
| <sup>1</sup> A <sub>2</sub> (π → 3s)              | 6.98       | 6.99  | 6.74 | 6.75  | 6.81 | 6.86   | 6.75             | 6.90             | 6.53             |                  |                   |                   |
| <sup>1</sup> B <sub>2</sub> (π → π*) <sup>j</sup> | 7.50       | 7.52  | 7.40 | 7.42  | 7.38 | 7.83   | 7.48             | 7.29             | 7.14             | 7.33             | 7.22              | 7.20              |
| <sup>1</sup> B <sub>1</sub> (π → 3p)              | 7.54       | 7.55  | 7.32 | 7.32  | 7.38 | 7.45   | 7.25             | 7.42             | 7.10             |                  |                   |                   |
| <sup>1</sup> A <sub>1</sub> (π → π*)              | 7.56       |       | 7.34 | 7.34  | 7.39 | 6.97   | 7.23             | 7.37             | 7.26             | 7.39             | 7.22              | 6.39              |
| <sup>3</sup> A <sub>1</sub> (π → π*)              | 4.33       | 4.31  | 4.34 | 4.31  | 4.33 | 4.60   | 4.05             |                  |                  | 4.28             |                   | 3.86              |
| <sup>3</sup> B <sub>1</sub> (n → π*)              | 4.57       | 4.56  | 4.47 | 4.47  | 4.46 | 4.58   | 4.41             |                  |                  | 4.42             |                   | 4.12              |
| <sup>3</sup> B <sub>2</sub> (π → π*)              | 4.92       | 4.91  | 4.83 | 4.83  | 4.79 | 4.88   | 4.56             |                  |                  | 4.72             |                   | 4.47              |
| <sup>3</sup> A <sub>1</sub> (π → π*)              | 5.14       | 5.13  | 5.08 |       | 5.05 | 5.19   | 4.73             |                  |                  | 4.96             |                   |                   |
| <sup>3</sup> A <sub>2</sub> (n → π*)              | 5.51       | 5.49  | 5.37 | 5.36  | 5.35 | 5.33   | 5.10             |                  |                  | 5.53             |                   | 5.40              |
| <sup>3</sup> B <sub>2</sub> (π → π*)              | 6.46       | 6.45  | 6.30 | 6.29  | 6.25 | 6.29   | 6.02             |                  |                  | 6.22             |                   | 6.09              |

<sup>a</sup>CASPT2 results from ref 30. <sup>b</sup>STEOM-CCSD results from ref 213. <sup>c</sup>EOM-CCSD( $\tilde{T}$ ) from ref 212. <sup>d</sup>CC3-ext. from ref 26. <sup>e</sup>EEL from ref 239. <sup>f</sup>EEL from ref 240. <sup>g</sup>CASPT2 from ref 241. <sup>h</sup>EEL from ref 242. <sup>i</sup>EEL from ref 243. <sup>j</sup>Significant state mixing with a close-lying Rydberg transition making unambiguous attribution difficult. At the CC3/aVDZ level, the Rydberg state is at 7.26 eV and has a small  $f_j$ , so attribution is rather clear. However, at the CC3/aVTZ level, the two B<sub>2</sub> transitions are at 7.35 and 7.38 eV (hence strongly mixed), so that the attribution has been made using the  $f$  values of 0.174 and 0.319, respectively.

There are many available studies of the ESs of pyrazine<sup>23,24,26,27,30,66,69,120,160,183,212,213,218,226,227</sup> and tetrazine,<sup>23–27,66,69,120,160,212,213,228–233</sup> for which the D<sub>2h</sub> symmetry helps distinguish the different ESs. Our results are collected in Tables 8 and S8. In pyrazine, all transitions are characterized by %T<sub>1</sub> > 85% at the exception of the <sup>1</sup>B<sub>1g</sub>(n → π\*) transition (84%). The excitation energies are basically unchanged going from CC3 to CCSDT except possibly for the highest-lying singlet state considered here. Going from triple- to quadruple- $\zeta$  basis, the variations do not exceed 0.04 eV, even for the four Rydberg ESs treated here. This indicates that one can be highly confident in the present estimates except for the highest-lying singlet ES. Again, the previous CASPT2 estimates<sup>23,30,183,226</sup> appear to be globally too low, while the (unconventional) CASPT3 results<sup>227</sup> seem too high. A similar overestimation can be noticed in previous SAC-CI results<sup>218</sup> and our NEVPT2 values, the latter showing a mean absolute deviation of 0.11 eV compared to CC3. In fact, the most satisfying agreement between the current estimates and previous works is reached with Nooijen's STEOM-CCSD values (except for the highest ES),<sup>213</sup> and the recent Schwabe–Goerigk's CC3 estimates.<sup>27</sup>

The available experimental data<sup>234–237</sup> do not include all theoretically predicted transitions but provide a similar energetic ranking for both singlets and triplets.

For tetrazine, we consider valence ESs only, including three transitions exhibiting a genuine double excitation nature (%T<sub>1</sub> < 10%). Of course, for these double excitations, CC3 and CCSDT cannot be considered as reliable. This is illustrated by the large change in excitation energies between these two CC models. The theoretical best estimates are likely obtained with NEVPT2.<sup>72</sup> For all the other transitions, the %T<sub>1</sub> values are in the 80–90% range for singlets and larger than 95% for triplets. Consequently, the CC3 and CCSDT results are very consistent, the sole exception being the lowest <sup>3</sup>B<sub>1u</sub>(π → π\*) transition for which we note a shift of −0.05 eV when upgrading the level of theory to CCSDT. In all other cases, there is a global consistency between our CC values. Moreover, the basis set effects are very small beyond aVTZ with a maximal variation of 0.02 eV when going to aVQZ (Table S8). The present values are almost systematically larger than the previous CASPT2,<sup>229</sup> STEOM-CCSD,<sup>231</sup> and GVVPT2<sup>219</sup> estimates. Our NEVPT2 values are also globally consistent with the CC3 values with a maximal

Table 10. Vertical Transition Energies (in eV) of Pyrimidine and Triazine

| state                                | Pyrimidine |       |      |       |      |        |                  |                  |                  |                  |                   |                   |
|--------------------------------------|------------|-------|------|-------|------|--------|------------------|------------------|------------------|------------------|-------------------|-------------------|
|                                      | 6-31+G(d)  |       | aVDZ |       | aVTZ |        |                  | lit.             |                  |                  |                   |                   |
|                                      | CC3        | CCSDT | CC3  | CCSDT | CC3  | NEVPT2 | th. <sup>a</sup> | th. <sup>b</sup> | th. <sup>c</sup> | th. <sup>d</sup> | exp. <sup>e</sup> | exp. <sup>f</sup> |
| <sup>1</sup> B <sub>1</sub> (n → π*) | 4.58       | 4.57  | 4.48 | 4.48  | 4.44 | 4.55   | 4.26             | 4.40             | 4.32             | 4.24             | 4.2               | 4.18              |
| <sup>1</sup> A <sub>2</sub> (n → π*) | 4.99       | 4.99  | 4.89 | 4.88  | 4.86 | 4.84   | 4.49             | 4.72             | 4.74             | 4.74             |                   | 4.69              |
| <sup>1</sup> B <sub>2</sub> (π → π*) | 5.47       | 5.44  | 5.44 | 5.41  | 5.41 | 5.53   | 5.47             | 5.04             | 5.29             | 5.01             | 5.12              | 5.18              |
| <sup>1</sup> A <sub>2</sub> (n → π*) | 6.07       | 6.06  | 5.98 | 5.97  | 5.93 | 6.02   |                  | 5.94             | 5.98             | 5.84             | 6.05              | 5.67              |
| <sup>1</sup> B <sub>1</sub> (n → π*) | 6.39       |       | 6.29 | 6.29  | 6.26 | 6.40   | 6.03             | 6.18             | 6.35             | 6.11             |                   | 6.02              |
| <sup>1</sup> B <sub>2</sub> (n → 3s) | 6.81       | 6.80  | 6.61 | 6.59  | 6.72 | 6.77   |                  | 6.85             | 6.84             | 6.57             |                   |                   |
| <sup>1</sup> A <sub>1</sub> (π → π*) | 7.08       | 7.09  | 6.93 | 6.94  | 6.87 | 7.11   | 7.10             | 6.87             | 6.86             | 6.57             | 6.7               | 6.69              |
| <sup>3</sup> B <sub>1</sub> (n → π*) | 4.20       | 4.20  | 4.12 | 4.11  | 4.10 | 4.17   | 3.81             |                  | 4.11             |                  |                   | 3.85              |
| <sup>3</sup> A <sub>1</sub> (π → π*) | 4.55       | 4.52  | 4.56 | 4.52  | 4.55 | 4.67   | 4.35             |                  | 4.39             |                  |                   | 4.42              |
| <sup>3</sup> A <sub>2</sub> (n → π*) | 4.77       | 4.76  | 4.67 | 4.67  | 4.66 | 4.72   | 4.24             |                  | 4.71             |                  |                   | 4.18              |
| <sup>3</sup> B <sub>2</sub> (π → π*) | 5.08       | 5.08  | 5.00 | 5.00  | 4.96 | 5.01   | 4.83             |                  | 4.81             |                  |                   | 4.93              |

| state                                 | Triazine  |       |      |       |      |       |        |                  |                  |                  |                  |                   |
|---------------------------------------|-----------|-------|------|-------|------|-------|--------|------------------|------------------|------------------|------------------|-------------------|
|                                       | 6-31+G(d) |       | aVDZ |       | aVTZ |       |        | lit.             |                  |                  |                  |                   |
|                                       | CC3       | CCSDT | CC3  | CCSDT | CC3  | CCSDT | NEVPT2 | th. <sup>g</sup> | th. <sup>b</sup> | th. <sup>d</sup> | th. <sup>h</sup> | exp. <sup>c</sup> |
| <sup>1</sup> A <sub>1</sub> '(n → π*) | 4.85      | 4.84  | 4.76 | 4.74  | 4.73 | 4.72  | 4.61   | 4.11             | 4.58             | 4.49             | 4.70             |                   |
| <sup>1</sup> A <sub>2</sub> '(n → π*) | 4.84      | 4.84  | 4.78 | 4.78  | 4.74 | 4.75  | 4.89   | 4.30             | 4.74             | 4.54             | 4.71             | 4.59              |
| <sup>1</sup> E''(n → π*)              | 4.89      | 4.89  | 4.82 | 4.81  | 4.78 | 4.78  | 4.88   | 4.32             | 4.69             | 4.56             | 4.75             | 3.97              |
| <sup>1</sup> A <sub>2</sub> '(π → π*) | 5.84      | 5.80  | 5.81 | 5.78  | 5.78 | 5.75  | 5.95   | 5.59             | 5.35             | 5.36             | 5.71             | 5.70              |
| <sup>1</sup> A <sub>1</sub> '(π → π*) | 7.45      | 7.45  | 7.31 | 7.31  | 7.24 | 7.24  | 7.30   |                  | 7.21             | 6.90             | 7.18             | 6.86              |
| <sup>1</sup> E'(n → 3s)               | 7.44      | 7.41  | 7.24 | 7.21  | 7.35 | 7.32  | 7.45   |                  | 7.38             | 7.16             |                  |                   |
| <sup>1</sup> E''(n → π*)              | 7.89      | 7.86  | 7.82 | 7.80  | 7.79 | 7.78  | 7.98   |                  |                  | 7.78             | 7.78             |                   |
| <sup>1</sup> E'(π → π*)               | 8.12      | 8.13  | 7.97 |       | 7.92 | 7.94  | 8.34   |                  | 7.82             | 7.72             | 7.84             | 7.76              |
| <sup>3</sup> A <sub>2</sub> '(n → π*) | 4.40      | 4.40  | 4.35 | 4.35  | 4.33 |       | 4.51   | 3.87             |                  |                  |                  |                   |
| <sup>3</sup> E''(n → π*)              | 4.59      | 4.59  | 4.52 | 4.52  | 4.51 |       | 4.61   | 4.04             |                  |                  |                  |                   |
| <sup>3</sup> A <sub>1</sub> '(n → π*) | 4.87      |       | 4.78 | 4.76  | 4.75 |       | 4.71   | 4.15             |                  |                  |                  |                   |
| <sup>3</sup> A <sub>1</sub> '(π → π*) | 4.88      | 4.85  | 4.88 | 4.85  | 4.88 |       | 5.05   |                  |                  |                  |                  |                   |
| <sup>3</sup> E'(π → π*)               | 5.70      | 5.68  | 5.64 |       | 5.61 |       | 5.73   |                  |                  |                  |                  |                   |
| <sup>3</sup> A <sub>2</sub> '(π → π*) | 6.85      | 6.84  | 6.69 | 6.68  | 6.63 |       | 6.36   | 4.76             |                  |                  |                  |                   |

<sup>a</sup>CASPT2 results from ref 251. <sup>b</sup>STEOM-CCSD results from ref 213. <sup>c</sup>SAC-CI results from ref 218. <sup>d</sup>EOM-CCSD( $\tilde{T}$ ) results from ref 212. <sup>e</sup>UV max from ref 234. <sup>f</sup>EEL from ref 253. <sup>g</sup>CASPT2 results from ref 255. <sup>h</sup>CC3-ext. results from ref 26.

discrepancy of 0.22 eV for the ESs with a dominant single excitation character. One finds a global agreement with Thiel's CC3/aVTZ values,<sup>26</sup> although we note variations of approximately 0.20 eV for specific excitations like the B<sub>2g</sub> transitions. This feature might be due to the use of distinct geometries in the two studies. The experimental EEL values from Palmer's work<sup>238</sup> show a reasonable agreement with our estimates.

**3.3.2. Pyridazine, Pyridine, Pyrimidine, and Triazine.** Those four azabenzenes with a C<sub>2v</sub> or D<sub>3h</sub> spatial symmetry are also popular molecules in terms of ES calculations.<sup>23,25–27,30,66,69,104,120,160,183,212,213,239,241,244–247</sup> Our results for pyridazine and pyridine are gathered in Tables 9 and S9. For the former compound, the available wave function results<sup>23,25–27,30,66,69,120,213,239,245</sup> focused on singlet transitions, at the exception of rather old MRCI,<sup>239</sup> and CASPT2 investigations.<sup>245</sup> Again, the %T<sub>1</sub> values are larger than 85% (95%) for the singlet (triplet) transitions, and the only state for which there is a variation larger than 0.03 eV between the CC3/aVDZ and the CCSDT/aVDZ energies is the <sup>3</sup>B<sub>2</sub>(π → π\*) transition. As in the previous six-membered cycles, the basis set effects are rather small, and aVTZ provides values close to the CBS limit for the considered transitions. For the singlet valence ESs, we find again a rather good match with the results of previous STEOM-CCSD<sup>213</sup> and CC<sup>26,212</sup> calculations. Yet again, these values are significantly higher than the CASPT2

estimates reported in refs 30 and 26. For the triplets, the present data represent the most accurate results published to date. Our NEVPT2 values are very close to their CC3 analogues for the lowest-lying singlet and triplet but positively deviate for the higher-lying ES. Interestingly, beyond the popular 20 year old reference measurements,<sup>239,248</sup> there is a very recent experimental EEL analysis for pyridazine<sup>240</sup> that locates almost all the ESs. The transition energies reported in this very recent work are systematically smaller than our CC estimates by approximately –0.20 eV. Nonetheless, this study provides exactly the same ES ranking as our theoretical protocol.

Pyridine, the hallmark organic heterocycle, has been more scrutinized than pyridazine, and many wave function approaches have been applied to estimate its ES energies.<sup>23,25–27,30,66,69,104,160,183,212,213,241,246,247</sup> Besides, two detailed EEL experiments are also available for pyridine.<sup>242,243</sup> The general trends described above for other six-membered cycles do pertain with (i) large %T<sub>1</sub> values and a nice consistency between CC3 and CCSDT estimates for all transitions listed in Table 9; (ii) small basis set effects beyond aVTZ even for the Rydberg transitions; (iii) qualitative agreement with past CC results; (iv) NEVPT2 transitions energies that are, on average, larger than their CC counterparts; and (v) the same ES ranking as in the most recent measurements.<sup>243</sup> Beyond these aspects, it is worth mentioning that the second <sup>1</sup>B<sub>2</sub>(π → π\*) ES is strongly mixed

Table 11. TBE Values (in eV) for All Considered States alongside Their Corresponding Oscillator Strength,  $f$ , and Percentage of Single Excitations,  $\%T_1$ , Obtained at the CC3/aVTZ level<sup>a</sup>

|                                   | state                                       | $f$   | $\%T_1$ | TBE/aVTZ |                       | TBE/CBS |       |
|-----------------------------------|---|-------|---------|----------|-----------------------|---------|-------|
|                                   |   |       |         | value    | protocol <sup>b</sup> | value   | corr. |
| acetone                           | $^1A_2(V; n \rightarrow \pi^*)$             |       | 91.1    | 4.47     | B                     | 4.48    | aVQZ  |
|                                   | $^1B_2(R; n \rightarrow 3s)$                | 0.000 | 90.5    | 6.46     | B                     | 6.51    | aVQZ  |
|                                   | $^1A_2(R; n \rightarrow 3p)$                |       | 90.9    | 7.47     | B                     | 7.44    | aVQZ  |
|                                   | $^1A_1(R; n \rightarrow 3p)$                | 0.004 | 90.6    | 7.51     | B                     | 7.55    | aVQZ  |
|                                   | $^1B_2(R; n \rightarrow 3p)$                | 0.029 | 91.2    | 7.62     | B                     | 7.63    | aVQZ  |
|                                   | $^3A_2(V; n \rightarrow \pi^*)$             |       | 97.8    | 4.13     | D                     | 4.15    | aVQZ  |
|                                   | $^3A_1(V; \pi \rightarrow \pi^*)$           |       | 98.7    | 6.25     | D                     | 6.27    | aVQZ  |
| acrolein                          | $^1A''(V; n \rightarrow \pi^*)$             | 0.000 | 87.6    | 3.78     | G                     | 3.79    | aVQZ  |
|                                   | $^1A'(V; \pi \rightarrow \pi^*)$            | 0.344 | 91.2    | 6.69     | CCSDT/aVTZ            | 6.69    | aVQZ  |
|                                   | $^1A''(V; n \rightarrow \pi^*)$             | 0.000 | 79.4    | 6.72     | D                     | 6.74    | aVQZ  |
|                                   | $^1A'(R; n \rightarrow 3s)$                 | 0.109 | 89.4    | 7.08     | D                     | 7.12    | aVQZ  |
|                                   | $^3A''(V; n \rightarrow \pi^*)$             |       | 97.0    | 3.51     | H                     | 3.50    | aVQZ  |
|                                   | $^3A'(V; \pi \rightarrow \pi^*)$            |       | 98.6    | 3.94     | D                     | 3.95    | aVQZ  |
|                                   | $^3A'(V; \pi \rightarrow \pi^*)$            |       | 98.4    | 6.18     | D                     | 6.19    | aVQZ  |
| benzene                           | $^3A''(V; n \rightarrow \pi^*)$             |       | 92.7    | 6.54     | E                     | 6.55    | aVQZ  |
|                                   | $^1B_{2u}(V; \pi \rightarrow \pi^*)$        |       | 86.3    | 5.06     | CCSDT/aVTZ            | 5.06    | aVQZ  |
|                                   | $^1B_{1u}(V; \pi \rightarrow \pi^*)$        |       | 92.9    | 6.45     | CCSDT/aVTZ            | 6.44    | aVQZ  |
|                                   | $^1E_{1g}(R; \pi \rightarrow 3s)$           |       | 92.8    | 6.52     | CCSDT/aVTZ            | 6.54    | aVQZ  |
|                                   | $^1A_{2u}(R; \pi \rightarrow 3p)$           | 0.066 | 93.4    | 7.08     | CCSDT/aVTZ            | 7.10    | aVQZ  |
|                                   | $^1E_{2u}(R; \pi \rightarrow 3p)$           |       | 92.8    | 7.15     | CCSDT/aVTZ            | 7.16    | aVQZ  |
|                                   | $^3B_{1u}(V; \pi \rightarrow \pi^*)$        |       | 98.6    | 4.16     | D                     | 4.17    | aVQZ  |
| butadiene                         | $^3E_{1u}(V; \pi \rightarrow \pi^*)$        |       | 97.1    | 4.85     | D                     | 4.86    | aVQZ  |
|                                   | $^3B_{2u}(V; \pi \rightarrow \pi^*)$        |       | 98.1    | 5.81     | D                     | 5.81    | aVQZ  |
|                                   | $^1B_u(V; \pi \rightarrow \pi^*)$           | 0.664 | 93.3    | 6.22     | B                     | 6.21    | aVQZ  |
|                                   | $^1B_g(R; \pi \rightarrow 3s)$              |       | 94.1    | 6.33     | B                     | 6.35    | aVQZ  |
|                                   | $^1A_g(V; \pi \rightarrow \pi^*)$           |       | 75.1    | 6.50     | F                     | 6.50    | aVQZ  |
|                                   | $^1A_u(R; \pi \rightarrow 3p)$              | 0.001 | 94.1    | 6.64     | B                     | 6.66    | aVQZ  |
|                                   | $^1A_u(R; \pi \rightarrow 3p)$              | 0.049 | 94.1    | 6.80     | B                     | 6.82    | aVQZ  |
| cyanoacetylene                    | $^1B_u(R; \pi \rightarrow 3p)$              | 0.055 | 93.8    | 7.68     | C                     | 7.54    | aVQZ  |
|                                   | $^3B_u(V; \pi \rightarrow \pi^*)$           |       | 98.4    | 3.36     | D                     | 3.37    | aVQZ  |
|                                   | $^3A_g(V; \pi \rightarrow \pi^*)$           |       | 98.7    | 5.20     | D                     | 5.21    | aVQZ  |
|                                   | $^3B_g(R; \pi \rightarrow 3s)$              |       | 97.9    | 6.29     | D                     | 6.31    | aVQZ  |
|                                   | $^1\Sigma^-(V; \pi \rightarrow \pi^*)$      |       | 94.3    | 5.80     | A                     | 5.79    | aVSZ  |
|                                   | $^1\Delta(V; \pi \rightarrow \pi^*)$        |       | 94.0    | 6.07     | A                     | 6.05    | aVSZ  |
|                                   | $^3\Sigma^+(V; \pi \rightarrow \pi^*)$      |       | 98.5    | 4.44     | CCSDT/aVTZ            | 4.46    | aVSZ  |
| cyanoformaldehyde                 | $^3\Delta(V; \pi \rightarrow \pi^*)$        |       | 98.2    | 5.21     | CCSDT/aVTZ            | 5.21    | aVSZ  |
|                                   | $^1A''[F](V; \pi \rightarrow \pi^*)$        | 0.004 | 93.6    | 3.54     | A                     | 3.54    | aVQZ  |
|                                   | $^1A''(V; n \rightarrow \pi^*)$             | 0.001 | 89.8    | 3.81     | CCSDT/aVTZ            | 3.82    | aVQZ  |
|                                   | $^1A''(V; \pi \rightarrow \pi^*)$           | 0.000 | 91.9    | 6.46     | CCSDT/aVTZ            | 6.45    | aVQZ  |
| cyanogen                          | $^3A''(V; n \rightarrow \pi^*)$             |       | 97.6    | 3.44     | D                     | 3.45    | aVQZ  |
|                                   | $^3A'(V; \pi \rightarrow \pi^*)$            |       | 98.4    | 5.01     | D                     | 5.02    | aVQZ  |
|                                   | $^1\Sigma_u^-(V; \pi \rightarrow \pi^*)$    |       | 94.1    | 6.39     | A                     | 6.38    | aVSZ  |
|                                   | $^1\Delta_u(V; \pi \rightarrow \pi^*)$      |       | 93.4    | 6.66     | A                     | 6.64    | aVSZ  |
| cyclopentadiene                   | $^3\Sigma_u^+(V; \pi \rightarrow \pi^*)$    |       | 98.5    | 4.91     | B                     | 4.93    | aVSZ  |
|                                   | $^1\Sigma_u^-[F](V; \pi \rightarrow \pi^*)$ |       | 93.4    | 5.05     | A                     | 5.03    | aVSZ  |
|                                   | $^1B_2(V; \pi \rightarrow \pi^*)$           | 0.084 | 93.8    | 5.56     | CCSDT/aVTZ            | 5.55    | aVQZ  |
|                                   | $^1A_2(R; \pi \rightarrow 3s)$              |       | 94.0    | 5.78     | CCSDT/aVTZ            | 5.80    | aVQZ  |
|                                   | $^1B_1(R; \pi \rightarrow 3p)$              | 0.037 | 94.2    | 6.41     | CCSDT/aVTZ            | 6.42    | aVQZ  |
|                                   | $^1A_2(R; \pi \rightarrow 3p)$              |       | 93.8    | 6.46     | CCSDT/aVTZ            | 6.47    | aVQZ  |
|                                   | $^1B_2(R; \pi \rightarrow 3p)$              | 0.046 | 94.2    | 6.56     | CCSDT/aVTZ            | 6.55    | aVQZ  |
|                                   | $^1A_1(V; \pi \rightarrow \pi^*)$           | 0.001 | 78.9    | 6.52     | CCSDT/aVTZ            | 6.52    | aVQZ  |
|                                   | $^3B_2(V; \pi \rightarrow \pi^*)$           |       | 98.4    | 3.31     | D                     | 3.31    | aVQZ  |
| $^3A_1(V; \pi \rightarrow \pi^*)$ |   | 98.6  | 5.11    | D        | 5.12                  | aVQZ    |       |
| $^3A_2(R; \pi \rightarrow 3s)$    |   | 97.9  | 5.73    | D        | 5.75                  | aVQZ    |       |
| $^3B_1(R; \pi \rightarrow 3p)$    |   | 97.9  | 6.36    | D        | 6.38                  | aVQZ    |       |



Table 11. continued

|   | state   | <i>f</i>  | % <i>T</i> <sub>1</sub> | TBE/aVTZ |                       | TBE/CBS |       |
|---|---|---|-------------------------|----------|-----------------------|---------|-------|
|   |   |   |                         | value    | protocol <sup>b</sup> | value   | corr. |
| cyclopropenone                                  | <sup>1</sup> B <sub>1</sub> (V; <i>n</i> → $\pi^*$ )                      | 0.000   | 87.7                    | 4.26     | B                     | 4.28    | aVSZ  |
|   | <sup>1</sup> A <sub>2</sub> (V; <i>n</i> → $\pi^*$ )                      |   | 91.0                    | 5.55     | B                     | 5.56    | aVSZ  |
|   | <sup>1</sup> B <sub>2</sub> (R; <i>n</i> → 3 <i>s</i> )                   | 0.003   | 90.8                    | 6.34     | B                     | 6.40    | aVSZ  |
|   | <sup>1</sup> B <sub>2</sub> (V; $\pi$ → $\pi^*$ )                         | 0.047   | 86.5                    | 6.54     | B                     | 6.56    | aVSZ  |
|   | <sup>1</sup> B <sub>2</sub> (R; <i>n</i> → 3 <i>p</i> )                   | 0.018   | 91.1                    | 6.98     | B                     | 7.01    | aVSZ  |
|   | <sup>1</sup> A <sub>1</sub> (R; <i>n</i> → 3 <i>p</i> )                   | 0.003   | 91.2                    | 7.02     | B                     | 7.08    | aVSZ  |
|   | <sup>1</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                         | 0.320   | 90.8                    | 8.28     | B                     | 8.26    | aVSZ  |
|   | <sup>3</sup> B <sub>1</sub> (V; <i>n</i> → $\pi^*$ )                      |   | 96.0                    | 3.93     | CCSDT/aVTZ            | 3.96    | aVSZ  |
|   | <sup>3</sup> B <sub>2</sub> (V; $\pi$ → $\pi^*$ )                         |   | 97.9                    | 4.88     | CCSDT/aVTZ            | 4.91    | aVSZ  |
|   | <sup>3</sup> A <sub>2</sub> (V; <i>n</i> → $\pi^*$ )                      |   | 97.5                    | 5.35     | CCSDT/aVTZ            | 5.37    | aVSZ  |
|   | <sup>3</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                         |   | 98.1                    | 6.79     | CCSDT/aVTZ            | 6.81    | aVSZ  |
| cyclopropenethione                              | <sup>1</sup> A <sub>2</sub> (V; <i>n</i> → $\pi^*$ )                      |   | 89.6                    | 3.41     | B                     | 3.41    | aVSZ  |
|   | <sup>1</sup> B <sub>1</sub> (V; <i>n</i> → $\pi^*$ )                      | 0.000   | 84.8                    | 3.45     | B                     | 3.48    | aVSZ  |
|   | <sup>1</sup> B <sub>2</sub> (V; $\pi$ → $\pi^*$ )                         | 0.007   | 83.0                    | 4.60     | B                     | 4.62    | aVSZ  |
|   | <sup>1</sup> B <sub>2</sub> (R; <i>n</i> → 3 <i>s</i> )                   | 0.048   | 91.8                    | 5.34     | B                     | 5.40    | aVSZ  |
|   | <sup>1</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                         | 0.228   | 89.0                    | 5.46     | B                     | 5.46    | aVSZ  |
|   | <sup>1</sup> B <sub>2</sub> (R; <i>n</i> → 3 <i>p</i> )                   | 0.084   | 91.3                    | 5.92     | B                     | 5.94    | aVSZ  |
|   | <sup>3</sup> A <sub>2</sub> (V; <i>n</i> → $\pi^*$ )                      |   | 97.2                    | 3.28     | D                     | 3.28    | aVSZ  |
|   | <sup>3</sup> B <sub>1</sub> (V; <i>n</i> → $\pi^*$ )                      |   | 94.5                    | 3.32     | CCSDT/aVTZ            | 3.36    | aVSZ  |
|   | <sup>3</sup> B <sub>2</sub> (V; $\pi$ → $\pi^*$ )                         |   | 96.5                    | 4.01     | D                     | 4.04    | aVSZ  |
|   | <sup>3</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                         |   | 98.2                    | 4.01     | D                     | 4.01    | aVSZ  |
|   | diacetylene   | <sup>1</sup> $\Sigma_u^-$ (V; $\pi$ → $\pi^*$ ) |                         | 94.4     | 5.33                  | A       | 5.32  |
| <sup>1</sup> $\Delta_u$ (V; $\pi$ → $\pi^*$ )   |   |   | 94.1                    | 5.61     | A                     | 5.60    | aVSZ  |
| <sup>3</sup> $\Sigma_u^+$ (V; $\pi$ → $\pi^*$ ) |   |   | 98.5                    | 4.10     | C                     | 4.13    | aVSZ  |
| <sup>3</sup> $\Delta_u$ (V; $\pi$ → $\pi^*$ )   |   |   | 98.2                    | 4.78     | B                     | 4.78    | aVSZ  |
| furan   | <sup>1</sup> A <sub>2</sub> (R; $\pi$ → 3 <i>s</i> )                      |   | 93.8                    | 6.09     | CCSDT/aVTZ            | 6.11    | aVQZ  |
|   | <sup>1</sup> B <sub>2</sub> (V; $\pi$ → $\pi^*$ )                         | 0.163   | 93.0                    | 6.37     | CCSDT/aVTZ            | 6.37    | aVQZ  |
|   | <sup>1</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                         | 0.000   | 92.4                    | 6.56     | CCSDT/aVTZ            | 6.56    | aVQZ  |
|   | <sup>1</sup> B <sub>1</sub> (R; $\pi$ → 3 <i>p</i> )                      | 0.038   | 93.9                    | 6.64     | CCSDT/aVTZ            | 6.66    | aVQZ  |
|   | <sup>1</sup> A <sub>2</sub> (R; $\pi$ → 3 <i>p</i> )                      |   | 93.6                    | 6.81     | CCSDT/aVTZ            | 6.83    | aVQZ  |
|   | <sup>1</sup> B <sub>2</sub> (R; $\pi$ → 3 <i>p</i> )                      | 0.008   | 93.5                    | 7.24     | D                     | 7.14    | aVQZ  |
|   | <sup>3</sup> B <sub>2</sub> (V; $\pi$ → $\pi^*$ )                         |   | 98.4                    | 4.20     | D                     | 4.20    | aVQZ  |
|   | <sup>3</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                         |   | 98.1                    | 5.46     | D                     | 5.47    | aVQZ  |
|   | <sup>3</sup> A <sub>2</sub> (R; $\pi$ → 3 <i>s</i> )                      |   | 97.9                    | 6.02     | D                     | 6.05    | aVQZ  |
|   | <sup>3</sup> B <sub>1</sub> (R; $\pi$ → 3 <i>p</i> )                      |   | 97.9                    | 6.59     | D                     | 6.61    | aVQZ  |
| glyoxal   | <sup>1</sup> A <sub>u</sub> (V; <i>n</i> → $\pi^*$ )                      | 0.000   | 91.0                    | 2.88     | B                     | 2.88    | aVSZ  |
|   | <sup>1</sup> B <sub>g</sub> (V; <i>n</i> → $\pi^*$ )                      |   | 88.3                    | 4.24     | B                     | 4.25    | aVSZ  |
|   | <sup>1</sup> A <sub>g</sub> (V; <i>n</i> , <i>n</i> → $\pi^*$ , $\pi^*$ ) |   | 0.5                     | 5.61     | F                     | 5.60    | aVSZ  |
|   | <sup>1</sup> B <sub>g</sub> (V; <i>n</i> → $\pi^*$ )                      |   | 83.9                    | 6.57     | B                     | 6.58    | aVSZ  |
|   | <sup>1</sup> B <sub>u</sub> (R; <i>n</i> → 3 <i>p</i> )                   | 0.095   | 91.7                    | 7.71     | B                     | 7.78    | aVSZ  |
|   | <sup>3</sup> A <sub>u</sub> (V; <i>n</i> → $\pi^*$ )                      |   | 97.6                    | 2.49     | CCSDT/aVTZ            | 2.50    | aVSZ  |
|   | <sup>3</sup> B <sub>g</sub> (V; <i>n</i> → $\pi^*$ )                      |   | 97.4                    | 3.89     | CCSDT/aVTZ            | 3.91    | aVSZ  |
|   | <sup>3</sup> B <sub>u</sub> (V; $\pi$ → $\pi^*$ )                         |   | 98.5                    | 5.15     | CCSDT/aVTZ            | 5.17    | aVSZ  |
|   | <sup>3</sup> A <sub>g</sub> (V; $\pi$ → $\pi^*$ )                         |   | 98.8                    | 6.30     | CCSDT/aVTZ            | 6.31    | aVSZ  |
|   | imidazole   | <sup>1</sup> A''(R; $\pi$ → 3 <i>s</i> )        | 0.001                   | 93.0     | 5.71                  | D       | 5.73  |
| <sup>1</sup> A'(V; $\pi$ → $\pi^*$ )            |   | 0.124   | 89.6                    | 6.41     | D                     | 6.41    | aVQZ  |
| <sup>1</sup> A''(V; <i>n</i> → $\pi^*$ )        |   | 0.028   | 93.6                    | 6.50     | D                     | 6.53    | aVQZ  |
| <sup>1</sup> A'(R; $\pi$ → 3 <i>p</i> )         |   | 0.035   | 88.9                    | 6.83     | D                     | 6.82    | aVQZ  |
| <sup>3</sup> A'(V; $\pi$ → $\pi^*$ )            |   |   | 98.3                    | 4.73     | E                     | 4.74    | aVQZ  |
| <sup>3</sup> A''(R; $\pi$ → 3 <i>s</i> )        |   |   | 97.6                    | 5.66     | D                     | 5.69    | aVQZ  |
| <sup>3</sup> A'(V; $\pi$ → $\pi^*$ )            |   |   | 97.9                    | 5.74     | E                     | 5.75    | aVQZ  |
| <sup>3</sup> A''(V; <i>n</i> → $\pi^*$ )        |   |   | 97.3                    | 6.31     | D                     | 6.31    | aVQZ  |
| isobutene                                       | <sup>1</sup> B <sub>1</sub> (R; $\pi$ → 3 <i>s</i> )                      | 0.006   | 94.1                    | 6.46     | CCSDT/aVTZ            | 6.48    | aVQZ  |
|   | <sup>1</sup> A <sub>1</sub> (R; $\pi$ → 3 <i>p</i> )                      | 0.228   | 94.2                    | 7.01     | CCSDT/aVTZ            | 7.00    | aVQZ  |
|   | <sup>3</sup> A <sub>1</sub> (V; ( $\pi$ → $\pi^*$ ))                      |   | 98.9                    | 4.53     | D                     | 4.54    | aVQZ  |
| methylenecyclopropene                           | <sup>1</sup> B <sub>2</sub> (V; $\pi$ → $\pi^*$ )                         | 0.011   | 85.4                    | 4.28     | B                     | 4.29    | aVSZ  |

Table 11. continued

|  | state                                      | <i>f</i>                                | % <i>T</i> <sub>1</sub> | TBE/aVTZ |                       | TBE/CBS |       |      |
|--|--|---|-------------------------|----------|-----------------------|---------|-------|------|
|  |  |   |                         | value    | protocol <sup>b</sup> | value   | corr. |      |
| propynal                                 | <sup>1</sup> B <sub>1</sub> (R; π → 3s)    | 0.005                                   | 93.6                    | 5.44     | B                     | 5.47    | aVSZ  |      |
|  | <sup>1</sup> A <sub>2</sub> (R; π → 3p)    |   | 93.3                    | 5.96     | B                     | 5.98    | aVSZ  |      |
|  | <sup>1</sup> A <sub>1</sub> (V; π → π*)    | 0.224                                   | 92.8                    | 6.12     | B                     | 6.03    | aVSZ  |      |
|  | <sup>3</sup> B <sub>2</sub> (V; π → π*)    |   | 97.2                    | 3.49     | CCSDT/aVTZ            | 3.50    | aVSZ  |      |
|  | <sup>3</sup> A <sub>1</sub> (V; π → π*)    |   | 98.6                    | 4.74     | D                     | 4.75    | aVSZ  |      |
|  | <sup>1</sup> A''(V; n → π*)                | 0.000                                   | 89.0                    | 3.80     | CCSDT/aVTZ            | 3.81    | aVQZ  |      |
|  | <sup>1</sup> A''(V; π → π*)                | 0.000                                   | 92.9                    | 5.54     | CCSDT/aVTZ            | 5.53    | aVQZ  |      |
|  | <sup>3</sup> A'(V; n → π*)                 |   | 97.4                    | 3.47     | D                     | 3.48    | aVQZ  |      |
| pyrazine                                 | <sup>3</sup> A'(V; π → π*)                 |   | 98.3                    | 4.47     | D                     | 4.48    | aVQZ  |      |
|  | <sup>1</sup> B <sub>3u</sub> (V; n → π*)   | 0.006                                   | 90.1                    | 4.15     | CCSDT/aVTZ            | 4.15    | aVQZ  |      |
|  | <sup>1</sup> A <sub>u</sub> (V; n → π*)    |   | 88.6                    | 4.98     | CCSDT/aVTZ            | 4.99    | aVQZ  |      |
|  | <sup>1</sup> B <sub>2u</sub> (V; π → π*)   | 0.078                                   | 86.9                    | 5.02     | CCSDT/aVTZ            | 5.01    | aVQZ  |      |
|  | <sup>1</sup> B <sub>2g</sub> (V; n → π*)   |   | 85.6                    | 5.71     | CCSDT/aVTZ            | 5.71    | aVQZ  |      |
|  | <sup>1</sup> A <sub>g</sub> (R; n → 3s)    |   | 91.1                    | 6.65     | CCSDT/aVTZ            | 6.69    | aVQZ  |      |
|  | <sup>1</sup> B <sub>1g</sub> (V; n → π*)   |   | 84.2                    | 6.74     | CCSDT/aVTZ            | 6.74    | aVQZ  |      |
|  | <sup>1</sup> B <sub>1u</sub> (V; π → π*)   | 0.063                                   | 92.8                    | 6.88     | CCSDT/aVTZ            | 6.87    | aVQZ  |      |
|  | <sup>1</sup> B <sub>1g</sub> (R; π → 3s)   |   | 93.8                    | 7.21     | CCSDT/aVTZ            | 7.24    | aVQZ  |      |
|  | <sup>1</sup> B <sub>2u</sub> (R; n → 3p)   | 0.037                                   | 90.8                    | 7.24     | D                     | 7.28    | aVQZ  |      |
|  | <sup>1</sup> B <sub>1u</sub> (R; n → 3p)   | 0.128                                   | 91.4                    | 7.44     | D                     | 7.47    | aVQZ  |      |
|  | <sup>1</sup> B <sub>1u</sub> (V; π → π*)   | 0.285                                   | 90.5                    | 7.98     | D                     | 7.97    | aVQZ  |      |
|  | <sup>3</sup> B <sub>3u</sub> (V; n → π*)   |   | 97.3                    | 3.59     | D                     | 3.59    | aVQZ  |      |
|  | <sup>3</sup> B <sub>1u</sub> (V; π → π*)   |   | 98.5                    | 4.35     | D                     | 4.36    | aVQZ  |      |
|  | <sup>3</sup> B <sub>2u</sub> (V; (π → π*)) |   | 97.6                    | 4.39     | D                     | 4.39    | aVQZ  |      |
|  | pyridazine                                 | <sup>3</sup> A <sub>u</sub> (V; n → π*) |                         | 96.1     | 4.93                  | D       | 4.94  | aVQZ |
| <sup>3</sup> B <sub>2g</sub> (V; n → π*) |  |   | 97.0                    | 5.08     | D                     | 5.09    | aVQZ  |      |
| <sup>3</sup> B <sub>1u</sub> (V; π → π*) |  |   | 97.0                    | 5.28     | D                     | 5.28    | aVQZ  |      |
| <sup>1</sup> B <sub>1</sub> (V; n → π*)  |  | 0.005                                   | 89.0                    | 3.83     | D                     | 3.83    | aVQZ  |      |
| <sup>1</sup> A <sub>2</sub> (V; n → π*)  |  |   | 86.9                    | 4.37     | D                     | 4.38    | aVQZ  |      |
| <sup>1</sup> A <sub>1</sub> (V; π → π*)  |  | 0.016                                   | 85.8                    | 5.26     | D                     | 5.26    | aVQZ  |      |
| <sup>1</sup> A <sub>2</sub> (V; n → π*)  |  |   | 86.2                    | 5.72     | D                     | 5.72    | aVQZ  |      |
| <sup>1</sup> B <sub>2</sub> (R; n → 3s)  |  | 0.001                                   | 88.5                    | 6.17     | D                     | 6.21    | aVQZ  |      |
| <sup>1</sup> B <sub>1</sub> (V; n → π*)  |  | 0.004                                   | 87.0                    | 6.37     | D                     | 6.37    | aVQZ  |      |
| <sup>1</sup> B <sub>2</sub> (V; π → π*)  |  | 0.010                                   | 90.6                    | 6.75     | D                     | 6.74    | aVQZ  |      |
| <sup>3</sup> B <sub>1</sub> (V; n → π*)  |  |   | 97.1                    | 3.19     | D                     | 3.20    | aVQZ  |      |
| <sup>3</sup> A <sub>2</sub> (V; n → π*)  |  |   | 96.2                    | 4.11     | D                     | 4.12    | aVQZ  |      |
| <sup>3</sup> B <sub>2</sub> (V; π → π*)  |  |   | 98.5                    | 4.34     | D                     | 4.35    | aVQZ  |      |
| <sup>3</sup> A <sub>1</sub> (V; π → π*)  |  |   | 97.3                    | 4.82     | D                     | 4.81    | aVQZ  |      |
| pyridine                                 |  | <sup>1</sup> B <sub>1</sub> (V; n → π*) | 0.004                   | 88.4     | 4.95                  | D       | 4.95  | aVQZ |
|  |  | <sup>1</sup> B <sub>2</sub> (V; π → π*) | 0.028                   | 86.5     | 5.14                  | D       | 5.14  | aVQZ |
|  | <sup>1</sup> A <sub>2</sub> (V; n → π*)    |   | 87.9                    | 5.40     | D                     | 5.41    | aVQZ  |      |
|  | <sup>1</sup> A <sub>1</sub> (V; π → π*)    | 0.010                                   | 92.1                    | 6.62     | D                     | 6.61    | aVQZ  |      |
|  | <sup>1</sup> A <sub>1</sub> (R; n → 3s)    | 0.011                                   | 89.7                    | 6.76     | D                     | 6.80    | aVQZ  |      |
|  | <sup>1</sup> A <sub>2</sub> (R; π → 3s)    |   | 93.2                    | 6.82     | D                     | 6.84    | aVQZ  |      |
|  | <sup>1</sup> B <sub>2</sub> (V; π → π*)    | 0.319                                   | 90.0                    | 7.40     | D                     | 7.42    | aVQZ  |      |
|  | <sup>1</sup> B <sub>1</sub> (R; π → 3p)    | 0.045                                   | 93.6                    | 7.38     | D                     | 7.40    | aVQZ  |      |
|  | <sup>1</sup> A <sub>1</sub> (V; π → π*)    | 0.291                                   | 90.5                    | 7.39     | D                     | 7.40    | aVQZ  |      |
|  | <sup>3</sup> A <sub>1</sub> (V; π → π*)    |   | 98.5                    | 4.30     | D                     | 4.31    | aVQZ  |      |
|  | <sup>3</sup> B <sub>1</sub> (V; n → π*)    |   | 97.0                    | 4.46     | D                     | 4.47    | aVQZ  |      |
|  | <sup>3</sup> B <sub>2</sub> (V; π → π*)    |   | 97.3                    | 4.79     | D                     | 4.79    | aVQZ  |      |
|  | <sup>3</sup> A <sub>1</sub> (V; π → π*)    |   | 97.1                    | 5.04     | E                     | 5.04    | aVQZ  |      |
|  | <sup>3</sup> A <sub>2</sub> (V; n → π*)    |   | 95.8                    | 5.36     | D                     | 5.38    | aVQZ  |      |
|  | <sup>3</sup> B <sub>2</sub> (V; π → π*)    |   | 97.7                    | 6.24     | D                     | 6.24    | aVQZ  |      |
|  | pyrimidine                                 | <sup>1</sup> B <sub>1</sub> (V; n → π*) | 0.005                   | 88.6     | 4.44                  | D       | 4.45  | aVQZ |
| <sup>1</sup> A <sub>2</sub> (V; n → π*)  |  |   | 88.5                    | 4.85     | D                     | 4.86    | aVQZ  |      |
| <sup>1</sup> B <sub>2</sub> (V; π → π*)  |  | 0.028                                   | 86.3                    | 5.38     | D                     | 5.37    | aVQZ  |      |
| <sup>1</sup> A <sub>2</sub> (V; n → π*)  |  |   | 86.7                    | 5.92     | D                     | 5.92    | aVQZ  |      |

Table 11. continued

|  | state  | <i>f</i>   | % <i>T</i> <sub>1</sub> | TBE/aVTZ |                       | TBE/CBS    |       |      |
|--|--|--|-------------------------|----------|-----------------------|------------|-------|------|
|  |  |  |                         | value    | protocol <sup>b</sup> | value      | corr. |      |
| pyrrole  | <sup>1</sup> B <sub>1</sub> (V; <i>n</i> → $\pi^*$ )                       | 0.005  | 86.7                    | 6.26     | D                     | 6.27       | aVQZ  |      |
|  | <sup>1</sup> B <sub>2</sub> (R; <i>n</i> → 3 <i>s</i> )                    | 0.005  | 90.3                    | 6.70     | D                     | 6.74       | aVQZ  |      |
|  | <sup>1</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                          | 0.036  | 91.5                    | 6.88     | D                     | 6.87       | aVQZ  |      |
|  | <sup>3</sup> B <sub>1</sub> (V; <i>n</i> → $\pi^*$ )                       |  | 96.8                    | 4.09     | D                     | 4.10       | aVQZ  |      |
|  | <sup>3</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                          |  | 98.3                    | 4.51     | D                     | 4.52       | aVQZ  |      |
|  | <sup>3</sup> A <sub>2</sub> (V; <i>n</i> → $\pi^*$ )                       |  | 96.5                    | 4.66     | D                     | 4.67       | aVQZ  |      |
|  | <sup>3</sup> B <sub>2</sub> (V; $\pi$ → $\pi^*$ )                          |  | 97.4                    | 4.96     | D                     | 4.96       | aVQZ  |      |
|  | <sup>1</sup> A <sub>2</sub> (R; $\pi$ → 3 <i>s</i> )                       |  | 92.9                    | 5.24     | CCSDT/aVTZ            | 5.27       | aVQZ  |      |
|  | <sup>1</sup> B <sub>1</sub> (R; $\pi$ → 3 <i>p</i> )                       | 0.015  | 92.4                    | 6.00     | CCSDT/aVTZ            | 6.03       | aVQZ  |      |
|  | <sup>1</sup> A <sub>2</sub> (R; $\pi$ → 3 <i>p</i> )                       |  | 93.0                    | 6.00     | D                     | 6.02       | aVQZ  |      |
|  | <sup>1</sup> B <sub>2</sub> (V; ( $\pi$ → $\pi^*$ ))                       | 0.164  | 92.5                    | 6.26     | CCSDT/aVTZ            | 6.23       | aVQZ  |      |
|  | <sup>1</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                          | 0.001  | 86.3                    | 6.30     | CCSDT/aVTZ            | 6.29       | aVQZ  |      |
|  | <sup>1</sup> B <sub>2</sub> (R; $\pi$ → 3 <i>p</i> )                       | 0.003  | 92.6                    | 6.83     | D                     | 6.74       | aVQZ  |      |
|  | <sup>3</sup> B <sub>2</sub> (V; $\pi$ → $\pi^*$ )                          |  | 98.3                    | 4.51     | D                     | 4.51       | aVQZ  |      |
|  | <sup>3</sup> A <sub>2</sub> (R; $\pi$ → 3 <i>s</i> )                       |  | 97.6                    | 5.21     | D                     | 5.24       | aVQZ  |      |
|  | tetrazine  | <sup>3</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )    |                         | 97.8     | 5.45                  | D          | 5.46  | aVQZ |
| <sup>3</sup> B <sub>1</sub> (R; $\pi$ → 3 <i>p</i> )                       |  |  | 97.4                    | 5.91     | D                     | 5.94       | aVQZ  |      |
| <sup>1</sup> B <sub>3u</sub> (V; <i>n</i> → $\pi^*$ )                      |  | 0.006  | 89.8                    | 2.47     | CCSDT/aVTZ            | 2.46       | aVQZ  |      |
| <sup>1</sup> A <sub>u</sub> (V; <i>n</i> → $\pi^*$ )                       |  |  | 87.9                    | 3.69     | CCSDT/aVTZ            | 3.70       | aVQZ  |      |
| <sup>1</sup> A <sub>g</sub> (V; <i>n</i> , <i>n</i> → $\pi^*$ , $\pi^*$ )  |  |  | 0.7                     | 4.61     | NEVPT2/aVTZ           | 4.59       | aVQZ  |      |
| <sup>1</sup> B <sub>1g</sub> (V; <i>n</i> → $\pi^*$ )                      |  |  | 83.1                    | 4.93     | CCSDT/aVTZ            | 4.92       | aVQZ  |      |
| <sup>1</sup> B <sub>2u</sub> (V; $\pi$ → $\pi^*$ )                         |  | 0.055  | 85.4                    | 5.21     | CCSDT/aVTZ            | 5.20       | aVQZ  |      |
| <sup>1</sup> B <sub>2g</sub> (V; <i>n</i> → $\pi^*$ )                      |  |  | 81.7                    | 5.45     | CCSDT/aVTZ            | 5.45       | aVQZ  |      |
| <sup>1</sup> A <sub>u</sub> (V; <i>n</i> → $\pi^*$ )                       |  |  | 87.7                    | 5.53     | CCSDT/aVTZ            | 5.53       | aVQZ  |      |
| <sup>1</sup> B <sub>3g</sub> (V; <i>n</i> , <i>n</i> → $\pi^*$ , $\pi^*$ ) |  |  | 0.7                     | 6.15     | NEVPT2/aVTZ           | 6.13       | aVQZ  |      |
| <sup>1</sup> B <sub>2g</sub> (V; <i>n</i> → $\pi^*$ )                      |  |  | 80.2                    | 6.12     | D                     | 6.12       | aVQZ  |      |
| <sup>1</sup> B <sub>1g</sub> (V; <i>n</i> → $\pi^*$ )                      |  |  | 85.1                    | 6.91     | D                     | 6.91       | aVQZ  |      |
| <sup>3</sup> B <sub>3u</sub> (V; <i>n</i> → $\pi^*$ )                      |  |  | 97.1                    | 1.85     | D                     | 1.86       | aVQZ  |      |
| <sup>3</sup> A <sub>u</sub> (V; <i>n</i> → $\pi^*$ )                       |  |  | 96.3                    | 3.45     | D                     | 3.46       | aVQZ  |      |
| <sup>3</sup> B <sub>1g</sub> (V; <i>n</i> → $\pi^*$ )                      |  |  | 97.0                    | 4.20     | D                     | 4.21       | aVQZ  |      |
| thioacetone  |  | <sup>3</sup> B <sub>1u</sub> (V; $\pi$ → $\pi^*$ )   |                         | 98.5     | 4.49                  | D          | 4.49  | aVQZ |
|  | <sup>3</sup> B <sub>2u</sub> (V; $\pi$ → $\pi^*$ )                         |  | 97.5                    | 4.52     | D                     | 4.52       | aVQZ  |      |
|  | <sup>3</sup> B <sub>2g</sub> (V; <i>n</i> → $\pi^*$ )                      |  | 96.4                    | 5.04     | D                     | 5.04       | aVQZ  |      |
|  | <sup>3</sup> A <sub>u</sub> (V; <i>n</i> → $\pi^*$ )                       |  | 96.6                    | 5.11     | D                     | 5.11       | aVQZ  |      |
|  | <sup>3</sup> B <sub>3g</sub> (V; <i>n</i> , <i>n</i> → $\pi^*$ , $\pi^*$ ) |  | 5.7                     | 5.51     | NEVPT2/aVTZ           | 5.50       | aVQZ  |      |
|  | <sup>3</sup> B <sub>1u</sub> (V; $\pi$ → $\pi^*$ )                         |  | 96.6                    | 5.42     | D                     | 5.43       | aVQZ  |      |
|  | <sup>1</sup> A <sub>2</sub> (V; <i>n</i> → $\pi^*$ )                       |  | 88.9                    | 2.53     | B                     | 2.54       | aVQZ  |      |
|  | <sup>1</sup> B <sub>2</sub> (R; <i>n</i> → 4 <i>s</i> )                    | 0.052  | 91.3                    | 5.56     | B                     | 5.61       | aVQZ  |      |
|  | <sup>1</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                          | 0.242  | 90.6                    | 5.88     | B                     | 5.86       | aVQZ  |      |
|  | <sup>1</sup> B <sub>2</sub> (R; <i>n</i> → 4 <i>p</i> )                    | 0.028  | 92.4                    | 6.51     | C                     | 6.52       | aVQZ  |      |
|  | <sup>1</sup> A <sub>1</sub> (R; <i>n</i> → 4 <i>p</i> )                    | 0.023  | 91.6                    | 6.61     | B                     | 6.64       | aVQZ  |      |
|  | <sup>3</sup> A <sub>2</sub> (V; <i>n</i> → $\pi^*$ )                       |  | 97.4                    | 2.33     | D                     | 2.34       | aVQZ  |      |
|  | <sup>3</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                          |  | 98.7                    | 3.45     | D                     | 3.46       | aVQZ  |      |
|  | thiophene  | <sup>1</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )    | 0.070                   | 87.6     | 5.64                  | CCSDT/aVTZ | 5.63  | aVQZ |
|  |  | <sup>1</sup> B <sub>2</sub> (V; $\pi$ → $\pi^*$ )    | 0.079                   | 91.5     | 5.98                  | CCSDT/aVTZ | 5.96  | aVQZ |
|  |  | <sup>1</sup> A <sub>2</sub> (R; $\pi$ → 3 <i>s</i> ) |                         | 92.6     | 6.14                  | CCSDT/aVTZ | 6.16  | aVQZ |
| <sup>1</sup> B <sub>1</sub> (R; $\pi$ → 3 <i>p</i> )                       |  | 0.010  | 90.1                    | 6.14     | CCSDT/aVTZ            | 6.11       | aVQZ  |      |
| <sup>1</sup> A <sub>2</sub> (R; $\pi$ → 3 <i>p</i> )                       |  |  | 91.8                    | 6.21     | CCSDT/aVTZ            | 6.18       | aVQZ  |      |
| <sup>1</sup> B <sub>1</sub> (R; $\pi$ → 3 <i>s</i> )                       |  | 0.000  | 92.8                    | 6.49     | CCSDT/aVTZ            | 6.52       | aVQZ  |      |
| <sup>1</sup> B <sub>2</sub> (R; $\pi$ → 3 <i>p</i> )                       |  | 0.082  | 92.4                    | 7.29     | CCSDT/aVTZ            | 7.18       | aVQZ  |      |
| <sup>1</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                          |  | 0.314  | 86.5                    | 7.31     | E                     | 7.29       | aVQZ  |      |
| <sup>3</sup> B <sub>2</sub> (V; $\pi$ → $\pi^*$ )                          |  |  | 98.2                    | 3.92     | D                     | 3.91       | aVQZ  |      |
| <sup>3</sup> A <sub>1</sub> (V; $\pi$ → $\pi^*$ )                          |  |  | 97.7                    | 4.76     | D                     | 4.76       | aVQZ  |      |
| <sup>3</sup> B <sub>1</sub> (R; $\pi$ → 3 <i>p</i> )                       |  |  | 96.6                    | 5.93     | D                     | 5.90       | aVQZ  |      |
| <sup>3</sup> A <sub>2</sub> (R; $\pi$ → 3 <i>s</i> )                       |  |  | 97.5                    | 6.08     | D                     | 5.98       | aVQZ  |      |
| thiopropynal   |  | <sup>1</sup> A''(V; <i>n</i> → $\pi^*$ )             | 0.000                   | 87.5     | 2.03                  | CCSDT/aVTZ | 2.04  | aVQZ |

Table 11. continued

|          | state   | <i>f</i> | % <i>T</i> <sub>1</sub> | TBE/aVTZ |                       | TBE/CBS |       |
|----------|---|----------|-------------------------|----------|-----------------------|---------|-------|
|          |   |          |                         | value    | protocol <sup>b</sup> | value   | corr. |
| triazine | <sup>3</sup> A''(V; <i>n</i> → π*)              |          | 97.2                    | 1.80     | D                     | 1.81    | aVQZ  |
|          | <sup>1</sup> A <sub>1</sub> '(V; <i>n</i> → π*) |          | 88.3                    | 4.72     | CCSDT/aVTZ            | 4.72    | aVQZ  |
|          | <sup>1</sup> A <sub>2</sub> '(V; <i>n</i> → π*) | 0.014    | 88.3                    | 4.75     | CCSDT/aVTZ            | 4.74    | aVQZ  |
|          | <sup>1</sup> E''(V; <i>n</i> → π*)              |          | 88.3                    | 4.78     | CCSDT/aVTZ            | 4.78    | aVQZ  |
|          | <sup>1</sup> A <sub>2</sub> '(V; π → π*)        |          | 85.7                    | 5.75     | CCSDT/aVTZ            | 5.75    | aVQZ  |
|          | <sup>1</sup> A <sub>1</sub> '(V; π → π*)        |          | 90.4                    | 7.24     | CCSDT/aVTZ            | 7.23    | aVQZ  |
|          | <sup>1</sup> E'(R; <i>n</i> → 3s)               | 0.016    | 90.9                    | 7.32     | CCSDT/aVTZ            | 7.36    | aVQZ  |
|          | <sup>1</sup> E'(V; <i>n</i> → π*)               |          | 82.6                    | 7.78     | CCSDT/aVTZ            | 7.76    | aVQZ  |
|          | <sup>1</sup> E'(V; π → π*)                      | 0.451    | 90.0                    | 7.94     | CCSDT/aVTZ            | 7.93    | aVQZ  |
|          | <sup>3</sup> A <sub>2</sub> '(V; <i>n</i> → π*) |          | 96.7                    | 4.33     | D                     | 4.34    | aVQZ  |
|          | <sup>3</sup> E''(V; <i>n</i> → π*)              |          | 96.6                    | 4.51     | D                     | 4.51    | aVQZ  |
|          | <sup>3</sup> A <sub>1</sub> '(V; <i>n</i> → π*) |          | 96.2                    | 4.73     | D                     | 4.74    | aVQZ  |
|          | <sup>3</sup> A <sub>1</sub> '(V; π → π*)        |          | 98.2                    | 4.85     | D                     | 4.86    | aVQZ  |
|          | <sup>3</sup> E'(V; π → π*)                      |          | 96.9                    | 5.59     | E                     | 5.59    | aVQZ  |
|          | <sup>3</sup> A <sub>2</sub> '(V; (π → π*))      |          | 97.6                    | 6.62     | D                     | 6.61    | aVQZ  |

<sup>a</sup>The composite protocol to generate these TBEs is also reported (see footnote *b*). In the right-most column, we list the TBE values obtained by including an additional correction (obtained at the CC3 level) for the basis set incompleteness error. Values displayed in italics are likely to be relatively less accurate. All values are obtained in the frozen-core approximation. <sup>b</sup>Protocol A: CCSDT/aVTZ value corrected by the difference between CCSDTQ/aVDZ and CCSDT/aVDZ. Protocol B: CCSDT/aVTZ value corrected by the difference between CCSDTQ/6-31+G(d) and CCSDT/6-31+G(d). Protocol C: CC3/aVTZ value corrected by the difference between CCSDTQ/6-31+G(d) and CC3/6-31+G(d). Protocol D: CC3/aVTZ value corrected by the difference between CCSDT/aVDZ and CC3/aVDZ. Protocol E: CC3/aVTZ value corrected by the difference between CCSDT/6-31+G(d) and CC3/6-31+G(d). Protocol F: FCI/aVDZ value (from ref 72) corrected by the difference between CCSDT/aVTZ and CCSDT/aVDZ. Protocol G: FCI/6-31+G(d) value corrected by the difference between CCSDT/aVTZ and CCSDT/6-31+G(d). Protocol H: FCI/6-31+G(d) value corrected by the difference between CC3/aVTZ and CC3/6-31+G(d).

with a nearby Rydberg transition that is separated by only 0.03 eV at the CC3/aVTZ level. This obviously makes the analysis particularly challenging for this specific transition.

The results obtained for both pyrimidine and triazine are listed in Tables 10 and S10. Because the former derivative can be viewed as the smallest model of DNA bases, previous theoretical<sup>23,25–27,30,66,69,160,183,212,213,218,249–251</sup> and experimental<sup>234,252,253</sup> studies are rather extensive. For triazine, which belongs to a nonabelian point group, theoretical studies are scarcer,<sup>23,25–27,30,66,69,212,213,244,254,255</sup> especially for the triplets,<sup>244,254,255</sup> whereas the experimental data are also limited.<sup>234,244</sup> As in pyridazine and pyridine, all the ESs listed in Table 10 show %*T*<sub>1</sub> values larger than 85% for singlets and 95% for triplets, so that CC3 and CCSDT are highly coherent, except maybe for the <sup>3</sup>A<sub>1</sub>'(π → π\*) transitions in pyrimidine. The basis set effects are also small, with no variation larger than 0.10 (0.03) eV between double- and triple-ζ (triple- and quadruple-ζ) for valence transitions and only slightly larger variations for the two Rydberg transitions (+0.04 eV between aVTZ and aVQZ). For both compounds, the current values are almost systematically larger than previously published data, with our CC3 values being typically bracketed by the published CASPT2 and the present NEVPT2 estimates. For the triplets of triazine, the three lowest ESs previously estimated by CASPT2<sup>255</sup> are too low by roughly half an electronvolt.

#### 4. THEORETICAL BEST ESTIMATES

Table 11 reports our two sets of TBEs: a set obtained with the aVTZ basis set and one set including an additional correction for the one-electron basis set incompleteness error. The details of our protocol employed to generate these TBEs are also provided in Table 11. For all states with a dominant single-excitation character (that is when %*T*<sub>1</sub> > 80%), we rely on CC results using an incremental strategy to generate these TBEs. As explained in


the footnotes of Table 11, this means that we add the basis set correction (i.e., the excitation energy difference between two calculations performed with a large and a small basis set) obtained with a “lower” level of theory, e.g., CC3, to correct the result obtained at a “higher” level of theory, e.g., CCSDTQ, but with the smaller basis set. In our previous contribution,<sup>28</sup> we have extensively tested this protocol for small compounds for which CCSDTQ/aVTZ calculations were achievable. It turned out that correcting CCSDTQ/6-31+G(d) with the CC3 or CCSDT basis set effect was very effective with a MAE of 0.01 eV as compared to the true value. There are only two exceptions for which we eschew using this CC incremental strategy: two ESs in acrolein for which nicely converged FCI values indicated non-negligible CCSDT errors. For ES with %*T*<sub>1</sub> values between 70% and 80%, our previous works indicated that CCSDT tends to overshoot the transition energies by roughly 0.05–0.10 eV and that NEVPT2 errors tend to be, on average, slightly larger.<sup>72</sup> Therefore, if CCSDTQ or FCI results are not available, it is extremely difficult to make the final call. For the other transitions, we relied either on the current or previous FCI data or the NEVPT2 values as reference. The italicized transition energies in Table 11 are believed to be (relatively) less accurate. This is the case when (i) the NEVPT2 result has to be selected; (ii) the CC calculations yield quite large changes in excitation energies while incrementing the excitation order despite large %*T*<sub>1</sub>; and (iii) there is a very strong ES mixing making hard to follow a specific transition from one method (or one basis) to another.

To determine the basis set corrections beyond augmented triple-ζ, we use the CC3/aVQZ or CC3/aVSZ results. For several compounds, we also provide in the SI CC3/d-aVQZ transition energies (i.e., with an additional set of diffuse functions). However, we do not consider such values as reference because the addition of a second set of diffuse orbitals

**Table 12.** Mean Signed Error (MSE), Mean Absolute Error (MAE), Root-Mean-Square Error (RMSE), Standard Deviation of the Errors (SDE), as Well as the Maximum Positive [Max(+)] and Negative [Max(-)] Errors with Respect to the TBE/aVTZ Reported in Table 11<sup>a</sup>

| method     | count | MSE  | MAE  | RMSE | SDE  | Max(+) | Max(-) |
|------------|-------|------|------|------|------|--------|--------|
| CIS(D)     | 221   | 0.16 | 0.23 | 0.29 | 0.24 | 0.96   | -0.69  |
| ADC(2)     | 218   | 0.01 | 0.14 | 0.20 | 0.19 | 0.64   | -0.73  |
| CC2        | 223   | 0.03 | 0.15 | 0.21 | 0.20 | 0.59   | -0.68  |
| STEOM-CCSD | 190   | 0.01 | 0.12 | 0.15 | 0.14 | 0.59   | -0.42  |
| CCSD       | 223   | 0.11 | 0.13 | 0.16 | 0.12 | 0.62   | -0.16  |
| CCSDR(3)   | 134   | 0.05 | 0.05 | 0.07 | 0.05 | 0.36   | -0.03  |
| CCSDT-3    | 127   | 0.05 | 0.05 | 0.07 | 0.04 | 0.26   | 0.00   |
| CC3        | 223   | 0.00 | 0.01 | 0.02 | 0.02 | 0.17   | -0.05  |
| NEVPT2     | 223   | 0.09 | 0.13 | 0.17 | 0.14 | 0.46   | -0.42  |

<sup>a</sup>All these statistical quantities are reported in eV and have been obtained with the aVTZ basis set. "Count" refers to the number of states.



**Figure 1.** Histograms of the error distribution obtained with various levels of theory, taking the TBE/aVTZ of Table 11 as references. Note the differences in scaling in the vertical axes.

only significantly modifies the transition energies when they also induce a stronger ES mixing. We also stick to the frozen-core approximation for two reasons: (i) the effect of correlating the core electrons is generally negligible (typically  $\pm 0.02$  eV) for the compounds under study (see the SI for examples), and (ii) it would be, in principle, necessary to add core polarization functions in such a case, which would have been technically difficult for most of the compounds considered here.

Table 11 encompasses 238 ESs, each of them obtained, at least, at the CCSDT level. This set can be decomposed as follows: 144 singlet and 94 triplet transitions, or 174 valence ( $99 \pi \rightarrow \pi^*$ ,  $71 n \rightarrow \pi^*$ , and 4 double excitations) and 64 Rydberg

transitions. Among these transition energies, 14 can be considered as "unsafe" and are reported in italics accordingly. This definitely corresponds to a very significant extension of our previous ES data sets (see Introduction). Taken all together, they offer a consistent, diverse, and accurate ensemble of transition energies for approximately 350 electronic transitions of various natures in small and medium-sized organic molecules. Table 11 also reports 90 oscillator strengths,  $f$ , which makes it, to the best of our knowledge, the largest set of CC3/aVTZ oscillator strengths reported to date, the previous effort being mostly performed at the CC3/TZVP level for Thiel's set.<sup>69</sup> It should also be pointed out that all these data are obtained on

CC3/aVTZ geometries, consistently with our previous works.<sup>28,72</sup>

## 5. BENCHMARKS

Having at hand such a large set of accurate transition energies, it seems natural to pursue previous benchmarking efforts. More specifically, we assess here the performance of eight popular wave function approaches, namely, CIS(D), ADC(2), CC2, STEOM-CCSD, CCSD, CCSDR(3), CCSDT-3, and CC3. The complete list of results can be found in Table S40 of the SI. To identify the ESs for all approaches, we have made, as for the TBE above, choices based on the usual criteria (symmetry, oscillator strength, ordering, and nature of the involved orbitals). Except for a few cases (see above), assignments are unambiguous. In addition, because all tested approaches are single-reference methods, we have removed from the reference set the “unsafe” transition energies (in italics in Table 11), as well as the four transitions with a dominant double excitation character (with %  $T_1 < 50\%$  as listed in Table 12). For the latter transitions, only CCSDT-3 and CC3 are able to detect their presence but with, of course, extremely large errors. A comprehensive list of results are collected in Table 12 which, more specifically, gathers the MSE, MAE, RMSE, SDE, Max(+), and Max(−). As benchmarks of the NEVPT2 method are quite rare, we have also considered the above-listed NEVPT2 values in our method evaluation. Of course, the results of such multiconfigurational approaches significantly depend on the active space, but our main purpose is to know what typical error one can expect with such a model when reasonable, yet “chemically meaningful” active spaces are considered. Figure 1 shows histograms of the error distributions for these nine methods. Before discussing these, let us stress two obvious biases of this molecular set: (i) it encompasses only conjugated organic molecules containing 4 to 6 non-hydrogen atoms, and (ii) we mainly used CCSDTQ (4 atoms) or CCSDT (5–6 atoms) reference values. As discussed in Section 3.1.5 and in our previous work,<sup>28</sup> the MAEs obtained with these two methods are of the order of 0.01 and 0.03 eV, respectively. This means that any statistical quantity smaller than ~0.02–0.03 eV is very likely to be irrelevant.

Let us analyze the global performance of all these methods, starting with the most accurate and computationally demanding single-reference models. The relative accuracy of CC3 and CCSDT-3 as compared to CCSDT remains an open question in the literature.<sup>56,71</sup> Indeed, to the best of our knowledge, the only two previous studies discussing this specific aspect are limited to very small compounds.<sup>28,70</sup> According to the results gathered in Table 12, it appears that CC3 has a slight edge over CCSDT-3, although CCSDT-3 is closer to CCSDT in terms of formalism. Indeed, CCSDT-3 seems to provide slightly too large transition energies (MSE of +0.05 eV). These conclusions are qualitatively consistent with the analyses performed on smaller derivatives,<sup>28,70</sup> but the amplitude of the CCSDT-3 deviations is larger in the present set. Although the performance of CC3 might be unduly inflated by the use of CCSDT and CCSDTQ reference values, it is also clear that CC3 very rarely fails (Figure 1). Consequently, CC3 transition energies can be viewed as extremely solid references for any transition with a dominant single-excitation character. This conclusion is again consistent with previous analyses performed on smaller compounds,<sup>28,70,148</sup> as well as with recent comparisons between theoretical and experimental 0-0 energies performed by some of us on medium-sized molecules.<sup>116,117,173</sup> To state it more boldly: it appears likely that CC3 is even more accurate than

previously thought. In addition, thanks to the exhaustive and detailed comparisons made in the present work, we can safely conclude that CC3 regularly outperforms CASPT2 and NEVPT2, even when these methods are combined with relatively large active spaces. This statement seems to hold as long as the considered ES does not show a strong multiple excitation character, that is, when % $T_1 > 70\%$ .

The perturbative inclusion of triples as in CCSDR(3) yields a very small MAE (0.05 eV) for a much lighter computational cost as compared to CCSDT. Nevertheless, as with CCSDT-3, the CCSDR(3) transition energies have a clear tendency of being too large, an error sign likely inherited from the parent CCSD model. The 0.05 eV MAE for CCSDR(3) is rather similar to the one obtained for smaller compounds when comparing to FCI (0.04 eV)<sup>28</sup> and is also inline with the 2009 benchmark study of Sauer et al.<sup>24</sup>

CCSD provides an interesting case study. The calculated MSE (+0.11 eV), indicating an overestimation of the transition energies, fits well with several previous reports.<sup>23,28,66,69–71,76,160,256</sup> It is, nonetheless, larger than the one determined for smaller molecules (+0.05 eV),<sup>28</sup> hinting that the performance of CCSD deteriorates for larger compounds. Moreover, the CCSD MAE of 0.13 eV is much smaller than the one reported by Thiel in his original work (0.49 eV)<sup>23</sup> but of the same order of magnitude as in the more recent study of Kánnár and Szalay performed on Thiel’s set (0.18 eV for transitions with % $T_1 > 90\%$ ).<sup>69</sup> Retrospectively, it is pretty obvious that Thiel’s much larger MAE is very likely due to the CASPT2 reference values.<sup>23</sup> Indeed, as we have shown several times in the present study, CASPT2 transition energies tend to be significantly too low, therefore exacerbating the usual CCSD overestimation.

With a single detailed benchmark study to date,<sup>66</sup> the STEOM-CCSD approach has received relatively little attention, and its overall accuracy still needs to be corroborated. It is noteworthy that STEOM-CCSD provides a smaller MSE than CCSD and comparable MAE and RMSE. The spread of the error is however slightly larger as evidenced by Figure 1 and the SDE values reported in Table 12. These trends are the same as for smaller compounds.<sup>28</sup> For Thiel’s set, Dutta and co-workers also reported a rather good performance for STEOM-CCSD with respect to the CC3/TZVP reference data, though a slightly negative MSE is obtained in their case.<sup>66</sup> This could well be due to the different basis sets considered in these two studies. It should be nevertheless stressed that we only consider “clean” STEOM-CCSD results in the present work (see Computational Details), therefore removing several difficult cases that are included in the CCSD statistics, e.g., the  $A_g$  excitation in butadiene, which can slightly bias the relative performance of STEOM-CCSD and CCSD.

For the three second-order methods, namely, CIS(D), ADC(2), and CC2, that are often used as reference to benchmark TD-DFT for “real-life” applications, the performance of the former method is clearly worse compared to the latter two which exhibit very similar statistical behaviors. These trends were also reported in previous works.<sup>18,21,23,26,28,59,70,76,148</sup> Interestingly, the CC2 MAE obtained here (0.15 eV) is significantly smaller than the one we found for smaller compounds (0.22 eV).<sup>28</sup> Therefore, in contrast to CCSD, CC2 performance seems to improve with molecular size. As above, Thiel’s original MAE for CC2 (0.29 eV) was likely too large due to the selection of CASPT2 reference values.<sup>23</sup> As already noticed by Szalay’s group,<sup>69,70</sup> although the MSE of CC2 is smaller than the one of CCSD, the

standard deviation is significantly larger with the former model, i.e., CC2 is less consistent in terms of trends than CCSD.

Finally, one obtains a reasonably tight distribution with NEVPT2, with a net overestimation trend (MSE of 0.09 eV) and a general behavior that is in fact quite comparable to (STEOM-)CCSD in terms of average and maximal deviations. Nonetheless, we wish to recall that NEVPT2 has the obvious advantage over CCSD to be able to treat accurately ES characterized by a dominant double excitation character. As mentioned above, these were not included in the present benchmark set.

In Table 13, we report a MAE decomposition for different subsets of ES. Note that, due to implementational limitations,

**Table 13. MAE (in eV) Obtained with Different Methods for Various Classes of Excited States**

| method     | singlet | triplet | valence | Rydberg | $n \rightarrow \pi^*$ | $\pi \rightarrow \pi^*$ |
|------------|---------|---------|---------|---------|-----------------------|-------------------------|
| CIS(D)     | 0.21    | 0.25    | 0.26    | 0.15    | 0.22                  | 0.28                    |
| ADC(2)     | 0.15    | 0.13    | 0.13    | 0.17    | 0.08                  | 0.17                    |
| CC2        | 0.16    | 0.14    | 0.14    | 0.18    | 0.08                  | 0.19                    |
| STEOM-CCSD | 0.11    | 0.13    | 0.11    | 0.12    | 0.08                  | 0.15                    |
| CCSD       | 0.16    | 0.09    | 0.14    | 0.09    | 0.19                  | 0.11                    |
| CCSDR(3)   | 0.05    | 0.07    | 0.07    | 0.02    | 0.08                  | 0.06                    |
| CCSDT-3    | 0.05    | 0.06    | 0.06    | 0.03    | 0.08                  | 0.04                    |
| CC3        | 0.01    | 0.01    | 0.01    | 0.01    | 0.01                  | 0.02                    |
| NEVPT2     | 0.15    | 0.12    | 0.13    | 0.15    | 0.11                  | 0.14                    |

only singlet ES could be computed with CCSDR(3) and CCSDT-3 which explains the lack of data for triplet ES. A few interesting conclusions emerge from these results. First, the errors for singlet and triplet transitions are rather similar in all the models, except for CCSD which is significantly more effective for triplets. Dutta and co-workers observed the same trend for Thiel's set with MAEs of 0.20 and 0.11 eV for the singlet and triplet ESs, respectively.<sup>66</sup> Turning to the comparison between valence and Rydberg states, we find that CC2 provides a better description of the former, whereas CCSD (and higher-order methods) yields the opposite trend. In fact, CC2 has the clear tendency to overestimate valence ES energies (MSE of +0.10 eV) and to underestimate Rydberg ES energies (MSE of -0.17 eV). CCSD is found to be much more consistent with MSEs of 0.12 and 0.09 eV, respectively (see the SI). This relatively poorer performance of CC2 as compared to CCSD for Rydberg ES is again perfectly consistent with other benchmarks,<sup>66,70</sup> although the MAE for CC2 (0.18 eV) reported in Table 13 remains relatively small as compared to the one given in ref 70. We believe that it is likely due to the distinct types of Rydberg states considered in these two studies. Indeed, we consider here (relatively) low-lying Rydberg transitions in medium-sized molecules, while Kánnár and Szalay (mostly) investigated higher-lying Rydberg states in smaller compounds. CIS(D), ADC(2), CC2, and STEOM-CCSD better describe  $n \rightarrow \pi^*$  transitions, whereas CCSD seems more suited for  $\pi \rightarrow \pi^*$  transitions. The variations between the two subsets are probably not significant for the higher-order approaches. The former observation agrees well with previous results obtained for smaller compounds<sup>28</sup> and for Thiel's set,<sup>23,69</sup> whereas the latter, less expected observation is likely dependent on the selected ES subset.<sup>23,70</sup> Finally, the average errors obtained with NEVPT2 are rather uniform for all subsets.

## 6. CONCLUDING REMARKS

We have computed highly accurate vertical transition energies for a set of 27 organic molecules containing from 4 to 6 (non-hydrogen) atoms. To this end, we employed several state-of-the-art theoretical models with increasingly large diffuse basis sets. Most of our theoretical best estimates are based on CCSDTQ (4 atoms) or CCSDT (5 and 6 atoms) excitation energies. For the vast majority of the listed excited states, the present contribution is the very first to disclose (sometimes basis-set extrapolated) CCSDT/aVTZ and (true) CC3/aVQZ transition energies as well as CC3/aVTZ oscillator strengths for each dipole-allowed transition. The present set contains a total of 238 transition energies and 90 oscillator strengths, with a reasonably good balance between singlet, triplet, valence, and Rydberg states. Among these 238 transitions, we believe that 224 are "solid" TBE; i.e., they are chemically accurate (MAE below 0.043 eV or 1 kcal·mol<sup>-1</sup>) for the considered geometry. It allowed us to establish a reasonable error bar for several popular ES models with a lower computational cost: CIS(D), ADC(2), CC2, STEOM-CCSD, CCSD, CCSDR(3), CCSDT-3, CC3, and NEVPT2. It turns out that CC3 is extremely accurate and very likely should be considered as globally more robust and trustworthy than CASPT2 or NEVPT2, except for ES with a predominant double excitation character. Other methods including corrections for the triples yield a mean absolute deviation around 0.05 eV, whereas none of the second-order approaches has been found to be chemically accurate with MAEs in the 0.12–0.23 eV range.

Paraphrasing Thiel and co-workers,<sup>23</sup> we hope that this new set of vertical transition energies, combined or not with the ones described in our previous works,<sup>28,72</sup> will be useful for the community, will stimulate further developments and analyses in the field, and will provide new grounds for appraising the *pros* and *cons* of ES models already available or currently under development. Looking into a crystal ball, we can forecast that the emergence of new SCI algorithms optimized for modern supercomputer architectures will likely lead to the revision of some of the present TBEs, allowing us to climb even higher on the accuracy ladder.<sup>257</sup>

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.jctc.9b01216>.

Basis set and frozen core effects, definition of the active spaces for the multiconfigurational calculations, additional details about the SCI calculations and their extrapolation, benchmark data and further statistical analysis, and geometries (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Authors

**Pierre-François Loos** – Laboratoire de Chimie et Physique Quantiques, CNRS et Université Toulouse III - Paul Sabatier, 31062 Toulouse, France; [orcid.org/0000-0003-0598-7425](https://orcid.org/0000-0003-0598-7425); Email: [loos@irsamc.ups-tlse.fr](mailto:loos@irsamc.ups-tlse.fr)

**Filippo Lipparini** – Dipartimento di Chimica e Chimica Industriale, University of Pisa, S6124 Pisa, Italy; [orcid.org/0000-0002-4947-3912](https://orcid.org/0000-0002-4947-3912); Email: [filippo.lipparini@unipi.it](mailto:filippo.lipparini@unipi.it)

**Denis Jacquemin** – CEISAM Lab, UMR 6230, Université de Nantes, CNRS, F-44000 Nantes, France; [orcid.org/0000-0002-4217-0708](https://orcid.org/0000-0002-4217-0708); Email: [denis.Jacquemin@univ-nantes.fr](mailto:denis.Jacquemin@univ-nantes.fr)

## Authors

Martial Boggio-Pasqua – Laboratoire de Chimie et Physique Quantiques, CNRS et Université Toulouse III - Paul Sabatier, 31062 Toulouse, France; [orcid.org/0000-0001-6684-5223](https://orcid.org/0000-0001-6684-5223)

Anthony Scemama – Laboratoire de Chimie et Physique Quantiques, CNRS et Université Toulouse III - Paul Sabatier, 31062 Toulouse, France

Complete contact information is available at:  
<https://pubs.acs.org/10.1021/acs.jctc.9b01216>

## Funding

P.-F.L. would like to thank the Centre National de la Recherche Scientifique for funding. D.J. acknowledges the Région des Pays de la Loire for financial support. This research used resources of (i) the GENCI-TGCC (Grant No. 2019-A0060801738); (ii) CCIPL (Centre de Calcul Intensif des Pays de Loire); (iii) the Troy cluster installed in Nantes; and (iv) CALMIP under allocations 2019-18005 (Toulouse). This work has been supported through the EUR Grant NanoX ANR-17-EURE-0009 in the framework of the “Programme des Investissements d’Avenir”.

## Notes

The authors declare no competing financial interest.

## REFERENCES

- (1) Casida, M. E. In *Time-Dependent Density-Functional Response Theory for Molecules*; Chong, D. P., Ed.; Recent Advances in Density Functional Methods; World Scientific: Singapore, 1995; Vol. 1; pp 155–192.
- (2) Ullrich, C. *Time-Dependent Density-Functional Theory: Concepts and Applications*; Oxford Graduate Texts; Oxford University Press: New York, 2012.
- (3) Adamo, C.; Jacquemin, D. The calculations of Excited-State Properties with Time-Dependent Density Functional Theory. *Chem. Soc. Rev.* **2013**, *42*, 845–856.
- (4) Laurent, A. D.; Jacquemin, D. TD-DFT Benchmarks: A Review. *Int. J. Quantum Chem.* **2013**, *113*, 2019–2039.
- (5) Dreuw, A.; Head-Gordon, M. Failure of Time-Dependent Density Functional Theory for Long-Range Charge-Transfer Excited States: the Zincbacteriochlorin-Bacteriochlorin and Bacteriochlorophyll-Spheroidene Complexes. *J. Am. Chem. Soc.* **2004**, *126*, 4007–4016.
- (6) Peach, M. J. G.; Benfield, P.; Helgaker, T.; Tozer, D. J. Excitation Energies in Density Functional Theory: an evaluation and a Diagnostic Test. *J. Chem. Phys.* **2008**, *128*, 044118.
- (7) Dierksen, M.; Grimme, S. A Density Functional Calculation of the Vibronic Structure of Electronic Absorption Spectra. *J. Chem. Phys.* **2004**, *120*, 3544–3554.
- (8) Dierksen, M.; Grimme, S. The Vibronic Structure of Electronic Absorption Spectra of Large Molecules: A Time-Dependent Density Functional Study on the Influence of Exact Hartree-Fock Exchange. *J. Phys. Chem. A* **2004**, *108*, 10225–10237.
- (9) Avila Ferrer, F. J.; Cerezo, J.; Stendardo, E.; Improta, R.; Santoro, F. Insights for an Accurate Comparison of Computational Data to Experimental Absorption and Emission Spectra: Beyond the Vertical Transition Approximation. *J. Chem. Theory Comput.* **2013**, *9*, 2072–2082.
- (10) Charaf-Eddin, A.; Planchat, A.; Mennucci, B.; Adamo, C.; Jacquemin, D. Choosing a Functional for Computing Absorption and Fluorescence Band Shapes with TD-DFT. *J. Chem. Theory Comput.* **2013**, *9*, 2749–2760.
- (11) Latouche, C.; Baiardi, A.; Barone, V. Virtual Eyes Designed for Quantitative Spectroscopy of Inorganic Complexes: Vibronic Signatures in the Phosphorescence Spectra of Terpyridine Derivatives. *J. Phys. Chem. B* **2015**, *119*, 7253–7257.
- (12) Muniz-Miranda, F.; Pedone, A.; Battistelli, G.; Montalti, M.; Bloino, J.; Barone, V. Benchmarking TD-DFT against Vibrationally

Resolved Absorption Spectra at Room Temperature: 7-Amino-coumarins as Test Cases. *J. Chem. Theory Comput.* **2015**, *11*, 5371–5384.

(13) Vazart, F.; Latouche, C.; Bloino, J.; Barone, V. Vibronic Coupling Investigation to Compute Phosphorescence Spectra of Pt(II) Complexes. *Inorg. Chem.* **2015**, *54*, 5588–5595.

(14) Santoro, F.; Jacquemin, D. Going Beyond the Vertical Approximation with Time-Dependent Density Functional Theory. *WIREs Comput. Mol. Sci.* **2016**, *6*, 460–486.

(15) Goerigk, L.; Grimme, S. Assessment of TD-DFT Methods and of Various Spin Scaled CIS<sub>n</sub>D and CC2 Versions for the Treatment of Low-Lying Valence Excitations of Large Organic Dyes. *J. Chem. Phys.* **2010**, *132*, 184103.

(16) Jacquemin, D.; Planchat, A.; Adamo, C.; Mennucci, B. A TD-DFT Assessment of Functionals for Optical 0–0 Transitions in Solvated Dyes. *J. Chem. Theory Comput.* **2012**, *8*, 2359–2372.

(17) Chibani, S.; Charaf-Eddin, A.; Le Guennic, B.; Jacquemin, D. Boronil and Related NBO Dyes: Insights From Theory. *J. Chem. Theory Comput.* **2013**, *9*, 3127–3135.

(18) Winter, N. O. C.; Graf, N. K.; Leutwyler, S.; Hättig, C. Benchmarks for 0–0 Transitions of Aromatic Organic Molecules: DFT/B3LYP, ADC(2), CC2, SOS-CC2 and SCS-CC2 Compared to High-resolution Gas-Phase Data. *Phys. Chem. Chem. Phys.* **2013**, *15*, 6623–6630.

(19) Fang, C.; Oruganti, B.; Durbeej, B. How Method-Dependent Are Calculated Differences Between Vertical, Adiabatic and 0–0 Excitation Energies? *J. Phys. Chem. A* **2014**, *118*, 4157–4171.

(20) Jacquemin, D.; Moore, B.; Planchat, A.; Adamo, C.; Autschbach, J. Performance of an Optimally Tuned Range-Separated Hybrid Functional for 0–0 Electronic Excitation Energies. *J. Chem. Theory Comput.* **2014**, *10*, 1677–1685.

(21) Jacquemin, D.; Duchemin, I.; Blase, X. 0–0 Energies Using Hybrid Schemes: Benchmarks of TD-DFT, CIS(D), ADC(2), CC2, and BSE/GW formalisms for 80 Real-Life Compounds. *J. Chem. Theory Comput.* **2015**, *11*, 5340–5359.

(22) Loos, P.-F.; Jacquemin, D. Evaluating 0–0 Energies with Theoretical Tools: a Short Review. *ChemPhotoChem.* **2019**, *3*, 684–696.

(23) Schreiber, M.; Silva-Junior, M. R.; Sauer, S. P. A.; Thiel, W. Benchmarks for Electronically Excited States: CASPT2, CC2, CCSD and CC3. *J. Chem. Phys.* **2008**, *128*, 134110.

(24) Sauer, S. P. A.; Schreiber, M.; Silva-Junior, M. R.; Thiel, W. Benchmarks for Electronically Excited States: A Comparison of Noniterative and Iterative Triples Corrections in Linear Response Coupled Cluster Methods: CCSDR(3) versus CC3. *J. Chem. Theory Comput.* **2009**, *5*, 555–564.

(25) Silva-Junior, M. R.; Sauer, S. P. A.; Schreiber, M.; Thiel, W. Basis Set Effects on Coupled Cluster Benchmarks of Electronically Excited States: CC3, CCSDR(3) and CC2. *Mol. Phys.* **2010**, *108*, 453–465.

(26) Silva-Junior, M. R.; Schreiber, M.; Sauer, S. P. A.; Thiel, W. Benchmarks of Electronically Excited States: Basis Set Effects Benchmarks of Electronically Excited States: Basis Set Effects on CASPT2 Results. *J. Chem. Phys.* **2010**, *133*, 174318.

(27) Schwabe, T.; Goerigk, L. Time-Dependent Double-Hybrid Density Functionals with Spin-Component and Spin-Opposite Scaling. *J. Chem. Theory Comput.* **2017**, *13*, 4307–4323.

(28) Loos, P.-F.; Scemama, A.; Blondel, A.; Garniron, Y.; Caffarel, M.; Jacquemin, D. A Mountaineering Strategy to Excited States: Highly-Accurate Reference Energies and Benchmarks. *J. Chem. Theory Comput.* **2018**, *14*, 4360–4379.

(29) Giner, E.; Scemama, A.; Toulouse, J.; Loos, P. F. Chemically Accurate Excitation Energies With Small Basis Sets. *J. Chem. Phys.* **2019**, *151*, 144118.

(30) Fulscher, M. P.; Andersson, K.; Roos, B. O. Toward an Accurate Molecular Orbital Theory for Excited States: the Azabenzene. *J. Phys. Chem.* **1992**, *96*, 9204–9212.

(31) Serrano-Andrés, L.; Merchán, M.; Nebot-Gil, I.; Lindh, R.; Roos, B. O. Towards an Accurate Molecular Orbital Theory for Excited



States: Ethene, Butadiene, and Hexatriene. *J. Chem. Phys.* **1993**, *98*, 3151–3162.

(32) Serrano-Andrés, L.; Merchán, M.; Nebot-Gil, I.; Roos, B. O.; Fulscher, M. Theoretical Study of the Electronic Spectra of Cyclopentadiene, Pyrrole, and Furan. *J. Am. Chem. Soc.* **1993**, *115*, 6184–6197.

(33) Serrano-Andrés, L.; Merchán, M.; Fulscher, M.; Roos, B. O. A Theoretical Study of the Electronic Spectrum of Thiophene. *Chem. Phys. Lett.* **1993**, *211*, 125–134.

(34) Lorentzon, J.; Malmqvist, P.-Å.; Fulscher, M.; Roos, B. O. A CASPT2 Study of the Valence and Lowest Rydberg Electronic States of Benzene and Phenol. *Theor. Chim. Acta* **1995**, *91*, 91–108.

(35) Merchán, M.; González-Luque, R.; Roos, B. O. A Theoretical Determination of the Electronic Spectrum of Methylene-cyclopropene. *Theor. Chim. Acta* **1996**, *94*, 143–154.

(36) Merchán, M.; Roos, B. O.; McDiarmid, R.; Xing, X. A Combined Theoretical and Experimental Determination of the Electronic Spectrum of Acetone. *J. Chem. Phys.* **1996**, *104*, 1791–1804.

(37) Roos, B. O.; Andersson, K.; Fulscher, M. P.; Malmqvist, P.-Å.; Serrano-Andrés, L. In *Advances in Chemical Physics*; Prigogine, I., Rice, S. A., Eds.; Wiley: New York, 1996; Vol. *XCIII*, pp 219–331.

(38) Serrano-Andrés, L.; Fulscher, M. P.; Roos, B. O.; Merchán, M. Theoretical Study of the Electronic Spectrum of Imidazole. *J. Phys. Chem.* **1996**, *100*, 6484–6491.

(39) Tozer, D. J.; Amos, R. D.; Handy, N. C.; Roos, B. O.; Serrano-Andrés, L. Does Density Functional Theory Contribute to the Understanding of Excited States of Unsaturated Organic Compounds? *Mol. Phys.* **1999**, *97*, 859–868.

(40) Burcl, R.; Amos, R. D.; Handy, N. C. Study of Excited States of Furan and Pyrrole by Time-Dependent Density Functional Theory. *Chem. Phys. Lett.* **2002**, *355*, 8–18.

(41) Zobel, J. P.; Nogueira, J. J.; Gonzalez, L. The IPEA Dilemma in CASPT2. *Chem. Sci.* **2017**, *8*, 1482–1499.

(42) Silva-Junior, M. R.; Schreiber, M.; Sauer, S. P. A.; Thiel, W. Benchmarks for Electronically Excited States: Time-Dependent Density Functional Theory and Density Functional Theory Based Multireference Configuration Interaction. *J. Chem. Phys.* **2008**, *129*, 104103.

(43) Goerigk, L.; Moellmann, J.; Grimme, S. Computation of Accurate Excitation Energies for Large Organic Molecules with Double-Hybrid Density Functionals. *Phys. Chem. Chem. Phys.* **2009**, *11*, 4611–4620.

(44) Jacquemin, D.; Wathelet, V.; Perpète, E. A.; Adamo, C. Extensive TD-DFT Benchmark: Singlet-Excited States of Organic Molecules. *J. Chem. Theory Comput.* **2009**, *5*, 2420–2435.

(45) Rohrdanz, M. A.; Martins, K. M.; Herbert, J. M. A Long-Range-Corrected Density Functional That Performs Well for Both Ground-State Properties and Time-Dependent Density Functional Theory Excitation Energies, Including Charge-Transfer Excited States. *J. Chem. Phys.* **2009**, *130*, 054112.

(46) Jacquemin, D.; Perpète, E. A.; Ciofini, I.; Adamo, C. Assessment of Functionals for TD-DFT Calculations of Singlet-Triplet Transitions. *J. Chem. Theory Comput.* **2010**, *6*, 1532–1537.

(47) Jacquemin, D.; Perpète, E. A.; Ciofini, I.; Adamo, C.; Valero, R.; Zhao, Y.; Truhlar, D. G. On the Performances of the M06 Family of Density Functionals for Electronic Excitation Energies. *J. Chem. Theory Comput.* **2010**, *6*, 2071–2085.

(48) Silva-Junior, M. R.; Thiel, W. Benchmark of Electronically Excited States for Semiempirical Methods: MNDO, AM1, PM3, OM1, OM2, OM3, INDO/S, and INDO/S2. *J. Chem. Theory Comput.* **2010**, *6*, 1546–1564.

(49) Mardirossian, N.; Parkhill, J. A.; Head-Gordon, M. Benchmark Results for Empirical Post-GGA Functionals: Difficult Exchange Problems and Independent Tests. *Phys. Chem. Chem. Phys.* **2011**, *13*, 19325–19337.

(50) Jacquemin, D.; Perpète, E. A.; Ciofini, I.; Adamo, C. Assessment of the  $\omega$ B97 Family for Excited-State Calculations. *Theor. Chem. Acc.* **2011**, *128*, 127–136.

(51) Huix-Rotllant, M.; Ipatov, A.; Rubio, A.; Casida, M. E. Assessment of Dressed Time-Dependent Density-Functional Theory for the Low-Lying Valence States of 28 Organic Chromophores. *Chem. Phys.* **2011**, *391*, 120–129.

(52) Della Sala, F.; Fabiano, E. Accurate Singlet and Triplet Excitation Energies Using the Localized Hartree–Fock Kohn–Sham Potential. *Chem. Phys.* **2011**, *391*, 19–26.

(53) Trani, F.; Scalmani, G.; Zheng, G. S.; Carnimeo, I.; Frisch, M. J.; Barone, V. Time-Dependent Density Functional Tight Binding: New Formulation and Benchmark of Excited States. *J. Chem. Theory Comput.* **2011**, *7*, 3304–3313.

(54) Peverati, R.; Truhlar, D. G. Performance of the M11 and M11-L Density Functionals for Calculations of Electronic Excitation Energies by Adiabatic Time-Dependent Density Functional Theory. *Phys. Chem. Chem. Phys.* **2012**, *14*, 11363–11370.

(55) Domínguez, A.; Aradi, B.; Frauenheim, T.; Lutsker, V.; Niehaus, T. A. Extensions of the Time-Dependent Density Functional Based Tight-Binding Approach. *J. Chem. Theory Comput.* **2013**, *9*, 4901–4914.

(56) Demel, O.; Datta, D.; Nooijen, M. Additional Global Internal Contraction in Variations of Multireference Equation of Motion Coupled Cluster Theory. *J. Chem. Phys.* **2013**, *138*, 134108.

(57) Schapiro, I.; Sivalingam, K.; Neese, F. Assessment of  $n$ -Electron Valence State Perturbation Theory for Vertical Excitation Energies. *J. Chem. Theory Comput.* **2013**, *9*, 3567–3580.

(58) Voityuk, A. A. INDO/X: A New Semiempirical Method for Excited States of Organic and Biological Molecules. *J. Chem. Theory Comput.* **2014**, *10*, 4950–4958.

(59) Harbach, P. H. P.; Wormit, M.; Dreuw, A. The Third-Order Algebraic Diagrammatic Construction Method (ADC(3)) for the Polarization Propagator for Closed-Shell Molecules: Efficient Implementation and Benchmarking. *J. Chem. Phys.* **2014**, *141*, 064113.

(60) Yang, Y.; Peng, D.; Lu, J.; Yang, W. Excitation Energies from Particle-Particle Random Phase Approximation: Davidson Algorithm and Benchmark Studies. *J. Chem. Phys.* **2014**, *141*, 124104.

(61) Sauer, S. P.; Pitzner-Frydendahl, H. F.; Buse, M.; Jensen, H. J. A.; Thiel, W. Performance of SOPPA-Based Methods in the Calculation of Vertical Excitation Energies and Oscillator Strengths. *Mol. Phys.* **2015**, *113*, 2026–2045.

(62) Piecuch, P.; Hansen, J. A.; Ajala, A. O. Benchmarking the Completely Renormalised Equation-Of-Motion Coupled-Cluster Approaches for Vertical Excitation Energies. *Mol. Phys.* **2015**, *113*, 3085–3127.

(63) Tajti, A.; Szalay, P. G. Investigation of the Impact of Different Terms in the Second Order Hamiltonian on Excitation Energies of Valence and Rydberg States. *J. Chem. Theory Comput.* **2016**, *12*, 5477–5482.

(64) Maier, T. M.; Bahmann, H.; Arbuznikov, A. V.; Kaupp, M. Validation of Local Hybrid Functionals for TDDFT Calculations of Electronic Excitation Energies. *J. Chem. Phys.* **2016**, *144*, 074106.

(65) Rishi, V.; Perera, A.; Nooijen, M.; Bartlett, R. J. Excited States from Modified Coupled Cluster Methods: are they any Better than EOM-CCSD? *J. Chem. Phys.* **2017**, *146*, 144104.

(66) Dutta, A. K.; Nooijen, M.; Neese, F.; Izsák, R. Exploring the Accuracy of a Low Scaling Similarity Transformed Equation of Motion Method for Vertical Excitation Energies. *J. Chem. Theory Comput.* **2018**, *14*, 72–91.

(67) Helmich-Paris, B. Benchmarks for Electronically Excited States with CASSCF Methods. *J. Chem. Theory Comput.* **2019**, *15*, 4170–4179.

(68) Haase, P. A. B.; Faber, R.; Provasi, P. F.; Sauer, S. P. A. Noniterative Doubles Corrections to the Random Phase and Higher Random Phase Approximations: Singlet and Triplet Excitation Energies. *J. Comput. Chem.* **2020**, *41*, 43–55.

(69) Kännár, D.; Szalay, P. G. Benchmarking Coupled Cluster Methods on Valence Singlet Excited States. *J. Chem. Theory Comput.* **2014**, *10*, 3757–3765.

- (70) Kánnár, D.; Tajti, A.; Szalay, P. G. Accuracy of Coupled Cluster Excitation Energies in Diffuse Basis Sets. *J. Chem. Theory Comput.* **2017**, *13*, 202–209.
- (71) Watson, T. J.; Lotrich, V. F.; Szalay, P. G.; Perera, A.; Bartlett, R. J. Benchmarking for Perturbative Triple-Excitations in EE-EOM-CC Methods. *J. Phys. Chem. A* **2013**, *117*, 2569–2579.
- (72) Loos, P.-F.; Boggio-Pasqua, M.; Scemama, A.; Caffarel, M.; Jacquemin, D. Reference Energies for Double Excitations. *J. Chem. Theory Comput.* **2019**, *15*, 1939–1956.
- (73) Chung, L. W.; Sameera, W. M. C.; Ramozzi, R.; Page, A. J.; Hatanaka, M.; Petrova, G. P.; Harris, T. V.; Li, X.; Ke, Z.; Liu, F.; Li, H.-B.; Ding, L.; Morokuma, K. The ONIOM Method and Its Applications. *Chem. Rev.* **2015**, *115*, S678–S796.
- (74) Papajak, E.; Zheng, J.; Xu, X.; Leverentz, H. R.; Truhlar, D. G. Perspectives on Basis Sets Beautiful: Seasonal Plantings of Diffuse Basis Functions. *J. Chem. Theory Comput.* **2011**, *7*, 3027–3034.
- (75) Budzák, S.; Scalmani, G.; Jacquemin, D. Accurate Excited-State Geometries: a CASPT2 and Coupled-Cluster Reference Database for Small Molecules. *J. Chem. Theory Comput.* **2017**, *13*, 6237–6252.
- (76) Jacquemin, D. What is the Key for Accurate Absorption and Emission Calculations? Energy or Geometry? *J. Chem. Theory Comput.* **2018**, *14*, 1534–1543.
- (77) Bremond, E.; Savarese, M.; Adamo, C.; Jacquemin, D. Accuracy of TD-DFT Geometries: a Fresh Look. *J. Chem. Theory Comput.* **2018**, *14*, 3715–3727.
- (78) Aidas, K.; Angeli, C.; Bak, K. L.; Bakken, V.; Bast, R.; Boman, L.; Christiansen, O.; Cimiraglia, R.; Coriani, S.; Dahle, P.; Dalskov, E. K.; Ekström, U.; Enevoldsen, T.; Eriksen, J. J.; Ettenhuber, P.; Fernández, B.; Ferrighi, L.; Fliegl, H.; Frediani, L.; Hald, K.; Halkier, A.; Hättig, C.; Heiberg, H.; Helgaker, T.; Hennum, A. C.; Hettema, H.; Hjertenæs, E.; Høst, S.; Høyvik, I.-M.; Iozzi, M. F.; Jansik, B.; Jensen, H. J. A.; Jonsson, D.; Jørgensen, P.; Kauczor, J.; Kirpekar, S.; Kjærgaard, T.; Klopper, W.; Knecht, S.; Kobayashi, R.; Koch, H.; Kongsted, J.; Krapp, A.; Kristensen, K.; Ligabue, A.; Lutnæs, O. B.; Melo, J. I.; Mikkelsen, K. V.; Myhre, R. H.; Neiss, C.; Nielsen, C. B.; Norman, P.; Olsen, J.; Olsen, J. M. H.; Osted, A.; Packer, M. J.; Pawłowski, F.; Pedersen, T. B.; Provasi, P. F.; Reine, S.; Rinkevicius, Z.; Ruden, T. A.; Ruud, K.; Rybkin, V. V.; Salek, P.; Samson, C. C. M.; de Merás, A. S.; Saue, T.; Sauer, S. P. A.; Schimmelpfennig, B.; Snekov, K.; Steindal, A. H.; Sylvester-Hvid, K. O.; Taylor, P. R.; Teale, A. M.; Tellgren, E. I.; Tew, D. P.; Thorvaldsen, A. J.; Thøgersen, L.; Vahtras, O.; Watson, M. A.; Wilson, D. J. D.; Ziolkowski, M.; Ågren, H. The Dalton Quantum Chemistry Program System. *WIREs Comput. Mol. Sci.* **2014**, *4*, 269–284.
- (79) CFOUR, Coupled-Cluster techniques for Computational Chemistry, a quantum-chemical program package by J. F. Stanton, J. Gauss, L. Cheng, M. E. Harding, D. A. Matthews, and P. G. Szalay with contributions from A. A. Auer, R. J. Bartlett, U. Benedikt, C. Berger, D. E. Bernholdt, Y. J. Bomble, O. Christiansen, F. Engel, R. Faber, M. Heckert, O. Heun, M. Hilgenberg, C. Huber, T.-C. Jagau, D. Jonsson, J. Jusélius, T. Kirsch, K. Klein, W. J. Lauderdale, F. Lipparini, T. Metzroth, L. A. Mück, D. P. O'Neill, D. R. Price, E. Prochnow, C. Puzzarini, K. Ruud, F. Schiffmann, W. Schwalbach, C. Simmons, S. Stopkowitz, A. Tajti, J. Vázquez, F. Wang, and J. D. Watts, and the integral packages MOLECULE (J. Almlöf and P. R. Taylor), PROPS (P. R. Taylor), ABACUS (T. Helgaker, H. J. Aa. Jensen, P. Jørgensen, and J. Olsen), and ECP routines by A. V. Mitin and C. van Wüllen. For the current version, see <http://www.cfour.de>.
- (80) Garniron, Y.; Applencourt, T.; Gasperich, K.; Benali, A.; Ferté, A.; Paquier, J.; Pradines, B.; Assaraf, R.; Reinhardt, P.; Toulouse, J.; Barbareco, P.; Renon, N.; David, G.; Malrieu, J.-P.; Véril, M.; Caffarel, M.; Loos, P.-F.; Giner, E.; Scemama, A. Quantum Package 2.0: An Open-Source Determinant-Driven Suite of Programs. *J. Chem. Theory Comput.* **2019**, *15*, 3591–3609.
- (81) Booth, G. H.; Thom, A. J. W.; Alavi, A. Fermion Monte Carlo Without Fixed Nodes: A Game of Life, Death, and Annihilation in Slater Determinant Space. *J. Chem. Phys.* **2009**, *131*, 054106.
- (82) Scemama, A.; Caffarel, M.; Benali, A.; Jacquemin, D.; Loos, P.-F. Influence of Pseudopotentials on Excitation Energies from Selected Configuration Interaction and Diffusion Monte Carlo. *Res. Chem.* **2019**, *1*, 100002.
- (83) Werner, H.-J.; Knowles, P. J.; Knizia, G.; Manby, F. R.; Schütz, M. Molpro: a general-purpose quantum chemistry program package. *Wiley Interdisciplinary Reviews: Computational Molecular Science* **2012**, *2*, 242–253.
- (84) Angeli, C.; Cimiraglia, R.; Evangelisti, S.; Leininger, T.; Malrieu, J.-P. Introduction of  $n$ -Electron Valence States for Multireference Perturbation Theory. *J. Chem. Phys.* **2001**, *114*, 10252–10264.
- (85) Angeli, C.; Cimiraglia, R.; Malrieu, J.-P.  $N$ -Electron Valence State Perturbation Theory: A Fast Implementation of the Strongly Contracted Variant. *Chem. Phys. Lett.* **2001**, *350*, 297–305.
- (86) Angeli, C.; Cimiraglia, R.; Malrieu, J.-P.  $N$ -Electron Valence State Perturbation Theory: A Spinless Formulation and an Efficient Implementation of the Strongly Contracted and of the Partially Contracted Variants. *J. Chem. Phys.* **2002**, *117*, 9138–9153.
- (87) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Petersson, G. A.; Nakatsuji, H.; Li, X.; Caricato, M.; Marenich, A. V.; Bloino, J.; Janesko, B. G.; Gomperts, R.; Mennucci, B.; Hratchian, H. P.; Ortiz, J. V.; Izmaylov, A. F.; Sonnenberg, J. L.; Williams-Young, D.; Ding, F.; Lipparini, F.; Egidi, F.; Goings, J.; Peng, B.; Petrone, A.; Henderson, T.; Ranasinghe, D.; Zakrzewski, V. G.; Gao, J.; Rega, N.; Zheng, G.; Liang, W.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Throssell, K.; Montgomery, J. A., Jr.; Peralta, J. E.; Ogliaro, F.; Bearpark, M. J.; Heyd, J. J.; Brothers, E. N.; Kudin, K. N.; Staroverov, V. N.; Keith, T. A.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A. P.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Millam, J. M.; Klene, M.; Adamo, C.; Cammi, R.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Farkas, O.; Foresman, J. B.; Fox, D. J. *Gaussian 16*, Revision A.03; Gaussian Inc.: Wallingford, CT, 2016.
- (88) Neese, F. The ORCA Program System. *WIREs Comput. Mol. Sci.* **2012**, *2*, 73–78.
- (89) Rolik, Z.; Szegedy, L.; Ladjanski, I.; Ladóczki, B.; Kállay, M. An Efficient Linear-Scaling CCSD(T) Method Based on Local Natural Orbitals. *J. Chem. Phys.* **2013**, *139*, 094105.
- (90) Kállay, M.; Rolik, Z.; Csontos, J.; Nagy, P.; Samu, G.; Mester, D.; Csóka, J.; Szabó, B.; Ladjanski, I.; Szegedy, L.; Ladóczki, B.; Petrov, K.; Farkas, M.; Mezei, P. D.; Hégyely, B. MRCC, Quantum Chemical Program; 2017. See [www.mrcc.hu](http://www.mrcc.hu).
- (91) Shao, Y.; Gan, Z.; Epifanovsky, E.; Gilbert, A. T.; Wormit, M.; Kussmann, J.; Lange, A. W.; Behn, A.; Deng, J.; Feng, X.; Ghosh, D.; Goldey, M.; Horn, P. R.; Jacobson, L. D.; Kalman, I.; Khaliullin, R. Z.; Kus, T.; Landau, A.; Liu, J.; Proynov, E. I.; Rhee, Y. M.; Richard, R. M.; Rohrdanz, M. A.; Steele, R. P.; Sundstrom, E. J.; Woodcock, H. L.; Zimmerman, P. M.; Zuev, D.; Albrecht, B.; Alguire, E.; Austin, B.; Beran, G. J. O.; Bernard, Y. A.; Berquist, E.; Brandhorst, K.; Bravaya, K. B.; Brown, S. T.; Casanova, D.; Chang, C.-M.; Chen, Y.; Chien, S. H.; Closser, K. D.; Crittenden, D. L.; Diedenhofen, M.; DiStasio, R. A.; Do, H.; Dutoi, A. D.; Edgar, R. G.; Fatehi, S.; Fusti-Molnar, L.; Ghysels, A.; Golubeva-Zadorozhnaya, A.; Gomes, J.; Hanson-Heine, M. W.; Harbach, P. H.; Hauser, A. W.; Hohenstein, E. G.; Holden, Z. C.; Jagau, T.-C.; Ji, H.; Kaduk, B.; Khistyayev, K.; Kim, J.; Kim, J.; King, R. A.; Klunzinger, P.; Kosenkov, D.; Kowalczyk, T.; Krauter, C. M.; Lao, K. U.; Laurent, A. D.; Lawler, K. V.; Levchenko, S. V.; Lin, C. Y.; Liu, F.; Livshits, E.; Lochan, R. C.; Luenser, A.; Manohar, P.; Manzer, S. F.; Mao, S.-P.; Mardirossian, N.; Marenich, A. V.; Maurer, S. A.; Mayhall, N. J.; Neuscamman, E.; Oana, C. M.; Olivares-Amaya, R.; O'Neill, D. P.; Parkhill, J. A.; Perrine, T. M.; Peverati, R.; Prociuk, A.; Rehn, D. R.; Rosta, E.; Russ, N. J.; Sharada, S. M.; Sharma, S.; Small, D. W.; Sodt, A.; Stein, T.; Stück, D.; Su, Y.-C.; Thom, A. J.; Tsuchimochi, T.; Vanovschi, V.; Vogt, L.; Vydrov, O.; Wang, T.; Watson, M. A.; Wenzel, J.; White, A.; Williams, C. F.; Yang, J.; Yeganeh, S.; Yost, S. R.; You, Z.-Q.; Zhang, I. Y.; Zhang, X.; Zhao, Y.; Brooks, B. R.; Chan, G. K.; Chipman, D. M.; Cramer, C. J.; Goddard, W. A.; Gordon, M. S.; Hehre, W. J.; Klamt, A.; Schaefer, H. F.; Schmidt, M. W.; Sherrill, C. D.; Truhlar, D. G.; Warshel, A.; Xu, X.; Aspuru-Guzik, A.; Baer, R.; Bell, A. T.; Besley, N. A.; Chai, J.-D.; Dreuw, A.; Dunietz, B. D.; Furlani, T. R.; Gwaltney, S. R.; Hsu, C.-

P.; Jung, Y.; Kong, J.; Lambrecht, D. S.; Liang, W.; Ochsenfeld, C.; Rassolov, V. A.; Slipchenko, L. V.; Subotnik, J. E.; Van Voorhis, T.; Herbert, J. M.; Krylov, A. I.; Gill, P. M.; Head-Gordon, M. Advances in Molecular Quantum Chemistry Contained in the Q-Chem 4 Program Package. *Mol. Phys.* **2015**, *113*, 184–215.

(92) Christiansen, O.; Koch, H.; Jørgensen, P. Response Functions in the CC3 Iterative Triple Excitation Model. *J. Chem. Phys.* **1995**, *103*, 7429–7441.

(93) Koch, H.; Christiansen, O.; Jørgensen, P.; Sanchez de Merás, A. M.; Helgaker, T. The CC3Model: An Iterative Coupled Cluster Approach Including Connected Triples. *J. Chem. Phys.* **1997**, *106*, 1808–1818.

(94) Watts, J. D.; Bartlett, R. J. Iterative and Non-Iterative Triple Excitation Corrections in Coupled-Cluster Methods for Excited Electronic States: the EOM-CCSDT-3 and EOM-CCSD( $\tilde{T}$ ) Methods. *Chem. Phys. Lett.* **1996**, *258*, 581–588.

(95) Prochnow, E.; Harding, M. E.; Gauss, J. Parallel Calculation of CCSDT and Mk-MRCCSDT Energies. *J. Chem. Theory Comput.* **2010**, *6*, 2339–2347.

(96) Noga, J.; Bartlett, R. J. The Full CCSDT Model for Molecular Electronic Structure. *J. Chem. Phys.* **1987**, *86*, 7041–7050.

(97) Kucharski, S. A.; Bartlett, R. J. Recursive Intermediate Factorization and Complete Computational Linearization of the Coupled-Cluster Single, Double, Triple, and Quadruple Excitation Equations. *Theor. Chim. Acta* **1991**, *80*, 387–405.

(98) Christiansen, O.; Koch, H.; Jørgensen, P. The Second-Order Approximate Coupled Cluster Singles and Doubles Model CC2. *Chem. Phys. Lett.* **1995**, *243*, 409–418.

(99) Hättig, C.; Weigend, F. CC2 Excitation Energy Calculations on Large Molecules Using the Resolution of the Identity Approximation. *J. Chem. Phys.* **2000**, *113*, 5154–5161.

(100) Purvis, G. P., III; Bartlett, R. J. A Full Coupled-Cluster Singles and Doubles Model: The Inclusion of Disconnected Triples. *J. Chem. Phys.* **1982**, *76*, 1910–1918.

(101) Christiansen, O.; Koch, H.; Jørgensen, P. Perturbative Triple Excitation Corrections to Coupled Cluster Singles and Doubles Excitation Energies. *J. Chem. Phys.* **1996**, *105*, 1451–1459.

(102) Head-Gordon, M.; Rico, R. J.; Oumi, M.; Lee, T. J. A Doubles Correction to Electronic Excited States From Configuration Interaction in the Space of Single Substitutions. *Chem. Phys. Lett.* **1994**, *219*, 21–29.

(103) Head-Gordon, M.; Maurice, D.; Oumi, M. A Perturbative Correction to Restricted Open-Shell Configuration-Interaction with Single Substitutions for Excited-States of Radicals. *Chem. Phys. Lett.* **1995**, *246*, 114–121.

(104) Nooijen, M.; Bartlett, R. J. A New Method for Excited States: Similarity Transformed Equation-Of-Motion Coupled-Cluster Theory. *J. Chem. Phys.* **1997**, *106*, 6441–6448.

(105) Dreuw, A.; Wormit, M. The Algebraic Diagrammatic Construction Scheme for the Polarization Propagator for the Calculation of Excited States. *WIREs Comput. Mol. Sci.* **2015**, *5*, 82–95.

(106) Luo, C.; Du, W. N.; Duan, X. M.; Li, Z. S. A Theoretical Study of the Photodissociation Mechanism of Cyanoacetylene in its Lowest Singlet and Triplet Excited States. *Astrophys. J.* **2008**, *687*, 726–730.

(107) Job, V.; King, G. The Electronic Spectrum of Cyanoacetylene: Part I. Analysis of the 2600-Å System. *J. Mol. Spectrosc.* **1966**, *19*, 155–177.

(108) Job, V.; King, G. The Electronic Spectrum of Cyanoacetylene: Part II. Analysis of the 2300-Å System. *J. Mol. Spectrosc.* **1966**, *19*, 178–184.

(109) Callomon, J. H.; Davey, A. B. Rotational Analysis of the 3000 Absorption System of Cyanogen, C<sub>2</sub>N<sub>2</sub>. *Proc. Phys. Soc., London* **1963**, *82*, 335–336.

(110) Bell, S.; Cartwright, G.; Fish, G.; O'Hare, D.; Ritchie, R.; Walsh, A.; Warsop, P. The Electronic Spectrum of Cyanogen. *J. Mol. Spectrosc.* **1969**, *30*, 162–163.

(111) Fish, G.; Cartwright, G.; Walsh, A.; Warsop, P. Rotational Structure in the  $^1\Sigma_u^- \leftarrow ^1\Sigma_g^+$  Transition of Cyanogen at 2200 Å. *J. Mol. Spectrosc.* **1972**, *41*, 20–32.

(112) Haink, H.-J.; Jungen, M. Excited States of the Polyacetylenes. Analysis of the Near Ultraviolet Spectra of Diacetylene and Triacetylene. *Chem. Phys. Lett.* **1979**, *61*, 319–322.

(113) Allan, M. The Excited States of 1,3-Butadiyne Determined by Electron Energy Loss Spectroscopy. *J. Chem. Phys.* **1984**, *80*, 6020–6024.

(114) Fischer, G.; Ross, I. G. Electronic Spectrum of Dicyanoacetylene. I. Calculations of the Geometries and Vibrations of Ground and Excited States of Diacetylene, Cyanoacetylene, Cyanogen, Triacetylene, Cyanodiacetylene, and Dicyanoacetylene. *J. Phys. Chem. A* **2003**, *107*, 10631–10636.

(115) Paterson, M. J.; Christiansen, O.; Pawłowski, F.; Jørgensen, P.; Hättig, C.; Helgaker, T.; Salek, P. Benchmarking Two-Photon Absorption with CC3 Quadratic Response Theory, and Comparison with Density-Functional Response Theory. *J. Chem. Phys.* **2006**, *124*, 054322.

(116) Loos, P.-F.; Galland, N.; Jacquemin, D. Theoretical 0–0 Energies with Chemical Accuracy. *J. Phys. Chem. Lett.* **2018**, *9*, 4646–4651.

(117) Loos, P.-F.; Jacquemin, D. Chemically Accurate 0–0 Energies with not-so-Accurate Excited State Geometries. *J. Chem. Theory Comput.* **2019**, *15*, 2481–2491.

(118) Hardwick, J. L.; Ramsay, D. A. The Near Ultraviolet Band System of Diacetylene. *Chem. Phys. Lett.* **1977**, *48*, 399–401.

(119) Caricato, M.; Mennucci, B.; Scalmani, G.; Trucks, G. W.; Frisch, M. J. Electronic Excitation Energies in Solution at Equation of Motion CCSD Level Within a State-Specific Polarizable Continuum Model Approach. *J. Chem. Phys.* **2010**, *132*, 084102.

(120) Leang, S. S.; Zahariev, F.; Gordon, M. S. Benchmarking the Performance of Time-Dependent Density Functional Methods. *J. Chem. Phys.* **2012**, *136*, 104101.

(121) Guareschi, R.; Filippi, C. Ground- and Excited-State Geometry Optimization of Small Organic Molecules with Quantum Monte Carlo. *J. Chem. Theory Comput.* **2013**, *9*, 5513–5525.

(122) Daday, C.; König, C.; Neugebauer, J.; Filippi, C. Wavefunction in Density Functional Theory Embedding for Excited States: Which Wavefunctions, which Densities? *ChemPhysChem* **2014**, *15*, 3205–3217.

(123) Guareschi, R.; Floris, F. M.; Amovilli, C.; Filippi, C. Solvent Effects on Excited-State Structures: A Quantum Monte Carlo and Density Functional Study. *J. Chem. Theory Comput.* **2014**, *10*, 5528–5537.

(124) Serrano-Andrés, L.; Pou-AméRigo, R.; Fülšcher, M. P.; Borin, A. C. Electronic Excited States of Conjugated Cyclic Ketones and Thioketones: A Theoretical Study. *J. Chem. Phys.* **2002**, *117*, 1649–1659.

(125) Liu, L.; Xia, S.; Fang, W.-H. Photodecarbonylation Mechanism of Cyclopropanone in the Gas Phase: Electronic Structure Calculation and AIMS Dynamics Simulation. *J. Phys. Chem. A* **2014**, *118*, 8977–8985.

(126) Harshbarger, W. R.; Kuebler, N. A.; Robin, M. B. Electronic Structure and Spectra of Small Rings. V Photoelectron and Electron Impact Spectra of Cyclopropanone. *J. Chem. Phys.* **1974**, *60*, 345–350.

(127) Breslow, R.; Oda, M. Isolation and Characterization of Pure Cyclopropanone. *J. Am. Chem. Soc.* **1972**, *94*, 4787–4788.

(128) Staley, S. W.; Norden, T. D. Synthesis and Direct Observation of Methylenecyclopropane. *J. Am. Chem. Soc.* **1984**, *106*, 3699–3700.

(129) Aquilante, F.; Barone, V.; Roos, B. O. A Theoretical Investigation of Valence and Rydberg Electronic States of Acrolein. *J. Chem. Phys.* **2003**, *119*, 12323–12334.

(130) Saha, B.; Ehara, M.; Nakatsuji, H. Singly and Doubly Excited States of Butadiene, Acrolein, and Glyoxal: Geometries and Electronic Spectra. *J. Chem. Phys.* **2006**, *125*, 014316.

(131) Mai, S.; Müller, T.; Plasser, F.; Marquetand, P.; Lischka, H.; González, L. Perturbational Treatment of Spin-Orbit Coupling for Generally Applicable High-Level Multi-Reference Methods. *J. Chem. Phys.* **2014**, *141*, 074105.

(132) Azarias, C.; Habert, C.; Budzák, Š.; Blase, X.; Duchemin, I.; Jacquemin, D. Calculations of  $n \rightarrow \pi^*$  Transition Energies:

Comparisons Between TD-DFT, ADC, CC, CASPT2, and BSE/GW Descriptions. *J. Phys. Chem. A* **2017**, *121*, 6122–6134.

(133) Bataev, V.; Panchenko, Y.; Abramov, A. A Quantum-Mechanical Analysis of trans-Acrolein Vibrational Spectra in the Ground  $S_0$  and Excited  $T_1$  and  $S_1$  Electronic States. *J. Mol. Struct.* **2017**, *1135*, 15–19.

(134) Dallos, M.; Lischka, H. A Systematic Theoretical Investigation of the Lowest Valence- and Rydberg-Excited Singlet States of Trans-Butadiene. The Character of the  ${}^1B_u$  ( $V$ ) State Revisited. *Theor. Chem. Acc.* **2004**, *112*, 16–26.

(135) Boggio-Pasqua, M.; Bearpark, M. J.; Klene, M.; Robb, M. A. A Computational Strategy for Geometry Optimization of Ionic and Covalent Excited States, Applied to Butadiene and Hexatriene. *J. Chem. Phys.* **2004**, *120*, 7849–7860.

(136) Li, X.; Paldus, J. Multi-Reference State-Universal Coupled-Cluster Approaches to Electronically Excited States. *J. Chem. Phys.* **2011**, *134*, 214118.

(137) Watson, M. A.; Chan, G. K.-L. Excited States of Butadiene to Chemical Accuracy: Reconciling Theory and Experiment. *J. Chem. Theory Comput.* **2012**, *8*, 4013–4018.

(138) Daday, C.; Smart, S.; Booth, G. H.; Alavi, A.; Filippi, C. Full Configuration Interaction Excitations of Ethene and Butadiene: Resolution of an Ancient Question. *J. Chem. Theory Comput.* **2012**, *8*, 4441–4451.

(139) Isegawa, M.; Peverati, R.; Truhlar, D. G. Performance of Recent and High-Performance Approximate Density Functionals for Time-Dependent Density Functional Theory Calculations of Valence and Rydberg Electronic Transition Energies. *J. Chem. Phys.* **2012**, *137*, 244104.

(140) Isegawa, M.; Truhlar, D. G. Valence Excitation Energies of Alkenes, Carbonyl Compounds, and Azabenzenes by Time-Dependent Density Functional Theory: Linear Response of the Ground State Compared to Collinear and Noncollinear Spin-Flip TDDFT with the Tamm-Dancoff Approximation. *J. Chem. Phys.* **2013**, *138*, 134111.

(141) Shu, Y.; Truhlar, D. G. Doubly Excited Character or Static Correlation of the Reference State in the Controversial  $2^1A_g$  State of trans-Butadiene? *J. Am. Chem. Soc.* **2017**, *139*, 13770–13778.

(142) Sokolov, A. Y.; Guo, S.; Ronca, E.; Chan, G. K.-L. Time-Dependent N-Electron Valence Perturbation Theory With Matrix Product State Reference Wavefunctions for Large Active Spaces and Basis Sets: Applications to the Chromium Dimer and all-trans Polyenes. *J. Chem. Phys.* **2017**, *146*, 244102.

(143) Chien, A. D.; Holmes, A. A.; Otten, M.; Umrigar, C. J.; Sharma, S.; Zimmerman, P. M. Excited States of Methylene, Polyenes, and Ozone from Heat-Bath Configuration Interaction. *J. Phys. Chem. A* **2018**, *122*, 2714–2722.

(144) Copan, A. V.; Sokolov, A. Y. Linear-Response Density Cumulant Theory for Excited Electronic States. *J. Chem. Theory Comput.* **2018**, *14*, 4097–4108.

(145) Tran, T.; Segarra-Martí, J.; Bearpark, M. J.; Robb, M. A. Molecular Vertical Excitation Energies Studied with First-Order RASSCF (RAS[1,1]): Balancing Covalent and Ionic Excited States. *J. Phys. Chem. A* **2019**, *123*, 5223–5230.

(146) Stanton, J. F.; Gauss, J. Theoretical Study of Electronically Excited cis- and trans-Glyoxal. *Spectrochim. Acta, Part A* **1997**, *53*, 1153–1162.

(147) Köhn, A.; Hättig, C. Analytic Gradients for Excited States in the Coupled-Cluster Model CC2 Employing the Resolution-Of-The-Identity Approximation. *J. Chem. Phys.* **2003**, *119*, 5021–5036.

(148) Hättig, C. In *Response Theory and Molecular Properties (A Tribute to Jan Lindenberg and Poul Jørgensen)*; Jensen, H. A., Ed.; Advances in Quantum Chemistry; Academic Press: 2005; Vol. 50, pp 37–60.

(149) Poorabdollah, H.; Omidyan, R.; Solimannejad, M.; Azimi, G. Hydrogen Bond Strengthening of cis–trans Glyoxal Dimers in Electronic Excited States: A Theoretical Study. *Spectrochim. Acta, Part A* **2014**, *122*, 337–342.

(150) Walsh, A. D. The Absorption Spectra of Acrolein, Crotonaldehyde and Mesityl Oxide in the Vacuum Ultra-Violet. *Trans. Faraday Soc.* **1945**, *41*, 498–505.

(151) Becker, R. S.; Inuzuka, K.; King, J. Acrolein: Spectroscopy, Photoisomerization, and Theoretical Considerations. *J. Chem. Phys.* **1970**, *52*, 5164–5170.

(152) Flicker, W. M.; Mosher, O. A.; Kuppermann, A. Electron-Impact Investigation of Excited Singlet States in 1,3-Butadiene. *Chem. Phys.* **1978**, *30*, 307–314.

(153) Doering, J. P.; McDiarmid, R. 100 eV Electron Impact Study of 1,3-Butadiene. *J. Chem. Phys.* **1981**, *75*, 2477–2478.

(154) Mosher, O. A.; Flicker, W. M.; Kuppermann, A. Electronic Spectroscopy of trans 1,3-Butadiene by Electron Impact. *J. Chem. Phys.* **1973**, *59*, 6502–6511.

(155) Verhaart, G.; Brongersma, H. Triplet  $\pi \rightarrow \pi^*$  and  $\pi \rightarrow \pi^*$  Transitions in Glyoxal and Biacetyl by Low-Energy Electron-Impact Spectroscopy. *Chem. Phys. Lett.* **1980**, *72*, 176–180.

(156) Robin, M. B. In *Higher Excited States of Polyatomic Molecules*; Robin, M. B., Ed.; Academic Press: 1985; Vol. III.

(157) Gwaltney, S. R.; Bartlett, R. J. An Application of the Equation-Of-Motion Coupled Cluster Method to the Excited States of Formaldehyde, Acetaldehyde, and Acetone. *Chem. Phys. Lett.* **1995**, *241*, 26–32.

(158) Wiberg, K. B.; Stratmann, R. E.; Frisch, M. J. A Time-Dependent Density Functional Theory Study of the Electronically Excited States of Formaldehyde, Acetaldehyde and Acetone. *Chem. Phys. Lett.* **1998**, *297*, 60–64.

(159) Wiberg, K. B.; de Oliveira, A. E.; Trucks, G. A Comparison of the Electronic Transition Energies for Ethene, Isobutene, Formaldehyde, and Acetone Calculated Using RPA, TDDFT, and EOM-CCSD. Effect of Basis Sets. *J. Phys. Chem. A* **2002**, *106*, 4192–4199.

(160) Caricato, M.; Trucks, G. W.; Frisch, M. J.; Wiberg, K. B. Electronic Transition Energies: A Study of the Performance of a Large Range of Single Reference Density Functional and Wave Function Methods on Valence and Rydberg States Compared to Experiment. *J. Chem. Theory Comput.* **2010**, *6*, 370–383.

(161) Pašteka, L. F.; Melicherčík, M.; Neogrady, P.; Urban, M. CASPT2 and CCSD(T) Calculations of Dipole Moments and Polarizabilities of Acetone in Excited States. *Mol. Phys.* **2012**, *110*, 2219–2237.

(162) Send, R.; Kühn, M.; Furche, F. Assessing Excited State Methods by Adiabatic Excitation Energies. *J. Chem. Theory Comput.* **2011**, *7*, 2376–2386.

(163) Birss, F. W.; Dong, R. Y.; Ramsay, D. A. The Band System of Propynal: Rotational Analysis of the 0–0 Band Near 4145 Å. *Can. J. Phys.* **1973**, *51*, 1810–1814.

(164) Judge, T. H.; Moule, D. C.; Bruno, A. E.; Steer, R. P. Thioketone Spectroscopy: An Analysis of the Lower Electronic Transitions in Thioacetone and Thioacetaldehyde. *Chem. Phys. Lett.* **1983**, *102*, 385–389.

(165) Brand, J.; Chan, W.; Liu, D.; Callomon, J.; Watson, J. The 3820 Å Band System of Propynal: Rotational Analysis of the 0–0 Band. *J. Mol. Spectrosc.* **1974**, *50*, 304–309.

(166) Staley, R. H.; Harding, L. B.; Goddard, W.; Beauchamp, J. Triplet States of the Amide Group. Trapped Electron Spectra of Formamide and Related Molecules. *Chem. Phys. Lett.* **1975**, *36*, 589–593.

(167) Johnson, K. E.; Johnston, D. B.; Lipsky, S. The Electron Impact Spectra of Some Monoolefinic Hydrocarbons. *J. Chem. Phys.* **1979**, *70*, 3844–3858.

(168) Judge, R.; Moule, D. A Vibronic Analysis of the Lower  $\tilde{X}^1A'' \leftarrow \tilde{X}^1A'$  Singlet-Singlet and  $\tilde{a}^3A'' \leftarrow \tilde{X}^1A'$  Triplet-Singlet Band Systems of Thiopropynal. *J. Mol. Spectrosc.* **1984**, *104*, 248–252.

(169) Robin, M. B. In *Higher Excited States of Polyatomic Molecules*; Robin, M. B., Ed.; Academic Press: 1985; Vol. III, pp 253–278.

(170) Palmer, M. H.; Beveridge, A. J.; Walker, I. C.; Abuain, T. M. Electronic and Cationic States of 2-Methylpropene (Isobutene) Studied by VUV Absorption, Scattered Electron Spectroscopy and ab

Initio Multireference Configuration Interaction Calculations. *Chem. Phys.* **1987**, *117*, 51–63.

(171) Karolczak, J.; Clouthier, D. J.; Judge, R.; Moule, D. High-Resolution Absorption and Pyrolysis Jet Spectroscopy of the  $0_0^0$  Band of the  $\tilde{A}^1A'' \leftarrow \tilde{X}^1A'' (n \rightarrow \pi^*)$  Electronic Transition of Formyl Cyanide, HCOCN. *J. Mol. Spectrosc.* **1991**, *147*, 61–70.

(172) Xing, X.; McDiarmid, R.; Philis, J. G.; Goodman, L. Vibrational Assignments in the  $3p$  Rydberg States of Acetone. *J. Chem. Phys.* **1993**, *99*, 7565–7573.

(173) Suellen, C.; Freitas, R. G.; Loos, P.-F.; Jacquemin, D. Cross Comparisons Between Experiment, TD-DFT, CC, and ADC for Transition Energies. *J. Chem. Theory Comput.* **2019**, *15*, 4581–4590.

(174) Nakano, H.; Tsuneda, T.; Hashimoto, T.; Hirao, K. Theoretical Study of the Excitation Spectra of Five-Membered Ring Compounds: Cyclopentadiene, Furan, and Pyrrole. *J. Chem. Phys.* **1996**, *104*, 2312–2320.

(175) Trofimov, A.; Schirmer, J. Polarization Propagator Study of Electronic Excitation in key Heterocyclic Molecules II. Furan. *Chem. Phys.* **1997**, *224*, 175–190.

(176) Christiansen, O.; Halkier, A.; Koch, H.; Jorgensen, P.; Helgaker, T. Integral-Direct Coupled Cluster Calculations of Frequency-Dependent Polarizabilities, Transition Probabilities and Excited-State Properties. *J. Chem. Phys.* **1998**, *108*, 2801–2816.

(177) Christiansen, O.; Jorgensen, P. The Electronic Spectrum of Furan. *J. Am. Chem. Soc.* **1998**, *120*, 3423–3430.

(178) Wan, J.; Meller, J.; Hada, M.; Ehara, M.; Nakatsuji, H. Electronic Excitation Spectra of Furan and Pyrrole: Revisited by the Symmetry Adapted Cluster–Configuration Interaction Method. *J. Chem. Phys.* **2000**, *113*, 7853–7866.

(179) Gromov, E. V.; Trofimov, A. B.; Vitkovskaya, N. M.; Schirmer, J.; Köppel, H. Theoretical Study of the Low-Lying Excited Singlet States of Furan. *J. Chem. Phys.* **2003**, *119*, 737–753.

(180) Pastore, M.; Angeli, C.; Cimiraaglia, R. An Application of Second and Third-Order  $n$ -Electron Valence State Perturbation Theory to the Calculation of the Vertical Electronic Spectrum of Furan. *Chem. Phys. Lett.* **2006**, *426*, 445–451.

(181) Shen, J.; Li, S. Block Correlated Coupled Cluster Method with the Complete Active-Space Self-Consistent-Field Reference Function: Applications for Low-Lying Electronic Excited States. *J. Chem. Phys.* **2009**, *131*, 174101.

(182) Li, X.; Paldus, J. A Multireference Coupled-Cluster Study of Electronic Excitations in Furan and Pyrrole. *J. Phys. Chem. A* **2010**, *114*, 8591–8600.

(183) Sauri, V.; Serrano-Andrés, L.; Shahi, A. R. M.; Gagliardi, L.; Vancoillie, S.; Pierloot, K. Multiconfigurational Second-Order Perturbation Theory Restricted Active Space (RASPT2) Method for Electronic Excited States: A Benchmark Study. *J. Chem. Theory Comput.* **2011**, *7*, 153–168.

(184) Holland, D.; Seddon, E.; Trofimov, A.; Gromov, E.; Wormit, M.; Dreuw, A.; Korona, T.; de Oliveira, N.; Archer, L.; Joyeux, D. A Study of the Excited Electronic States of Normal and Fully Deuterated Furan by Photoabsorption Spectroscopy and High-Level *ab initio* Calculations. *J. Mol. Spectrosc.* **2015**, *315*, 184–195.

(185) Van Veen, E. Triplet  $\pi \rightarrow \pi^*$  Transitions in Thiophene, Furan and Pyrrole by Low-Energy Electron-Impact Spectroscopy. *Chem. Phys. Lett.* **1976**, *41*, 535–539.

(186) Flicker, W. M.; Mosher, O. A.; Kuppermann, A. Electron Impact Investigation of Electronic Excitations in Furan, Thiophene, and Pyrrole. *J. Chem. Phys.* **1976**, *64*, 1315–1321.

(187) Pastore, M.; Angeli, C.; Cimiraaglia, R. The Vertical Electronic Spectrum of Pyrrole: A second and Third Order  $n$ -Electron Valence State Perturbation Theory Study. *Chem. Phys. Lett.* **2006**, *422*, 522–528.

(188) Christiansen, O.; Gauss, J.; Stanton, J. F.; Jorgensen, P. The Electronic Spectrum of Pyrrole. *J. Chem. Phys.* **1999**, *111*, 525–537.

(189) Heindl, M.; González, L. A XMS-CASPT2 non-Adiabatic Dynamics Study on Pyrrole. *Comput. Theor. Chem.* **2019**, *1155*, 38–46.

(190) Flicker, W. M.; Mosher, O. A.; Kuppermann, A. Triplet States of Furan, Thiophene, and Pyrrole. *Chem. Phys. Lett.* **1976**, *38*, 489–492.

(191) Palmer, M. H.; Wilson, P. J. The Singlet Electronic States of Pyrrole: a Theoretical Study by Both *ab initio* Multi-Reference Configuration Interaction Methods and Time-Dependent Density Functional Theory and a Reconsideration of the Experimental VUV Spectral Data. *Mol. Phys.* **2003**, *101*, 2391–2408.

(192) Horváth, G.; Kiss, Á. The Electronic Spectra of Five-membered Heterocyclic Compounds. *Spectrochim. Acta, Part A* **1967**, *23*, 921–924.

(193) Bavia, M.; Bertinelli, F.; Taliani, C.; Zauli, C. The Electronic Spectrum of Pyrrole in the Vapor and Crystal. *Mol. Phys.* **1976**, *31*, 479–489.

(194) Trofimov, A.; Schirmer, J. Polarization Propagator Study of Electronic Excitation in key Heterocyclic Molecules I. Pyrrole. *Chem. Phys.* **1997**, *214*, 153–170.

(195) Palmer, M. H.; Walker, I. C.; Guest, M. F. The Electronic States of Pyrrole Studied by Optical (VUV) Absorption, Near-Threshold Electron Energy-Loss (EEL) Spectroscopy and *ab initio* Multi-Reference Configuration Interaction Calculations. *Chem. Phys.* **1998**, *238*, 179–199.

(196) Roos, B. O.; Malmqvist, P.-Å.; Molina, V.; Serrano-Andrés, L.; Merchán, M. Theoretical Characterization of the Lowest-Energy Absorption Band of Pyrrole. *J. Chem. Phys.* **2002**, *116*, 7526–7536.

(197) Neville, S. P.; Worth, G. A. A Reinterpretation of the Electronic Spectrum of Pyrrole: A Quantum Dynamics Study. *J. Chem. Phys.* **2014**, *140*, 034317.

(198) Wan, J.; Ehara, M.; Hada, M.; Nakatsuji, H. Electronic Excitation and Ionization Spectra of Cyclopentadiene: Revisit by the Symmetry-Adapted Cluster–Configuration Interaction Method. *J. Chem. Phys.* **2000**, *113*, 5245–5252.

(199) Frueholz, R. P.; Flicker, W. M.; Mosher, O. A.; Kuppermann, A. Electronic Spectroscopy of 1,3-Cyclopentadiene, 1,3-Cyclohexadiene and 1,3-Cycloheptadiene by Electron Impact. *J. Chem. Phys.* **1979**, *70*, 2003–2013.

(200) McDiarmid, R.; Sabljic, A. Analysis of the Absorption Spectrum of the 195-nm Region of Cyclopentadiene. *J. Phys. Chem.* **1991**, *95*, 6455–6462.

(201) McDiarmid, R.; Sabljic, A.; Doering, J. P. Valence Transitions in 1,3-Cyclopentadiene, 1,3-Cyclohexadiene, and 1,3-Cycloheptadiene. *J. Chem. Phys.* **1985**, *83*, 2147–2152.

(202) Sabljic, A.; McDiarmid, R.; Gedanken, A. A Two-Photon Resonant Multiphoton Ionization Study of the  $3p$ -Rydberg  $\leftarrow \tilde{X}$  Transitions of Cyclopentadiene. *J. Phys. Chem.* **1992**, *96*, 2442–2448.

(203) Devine, A. L.; Cronin, B.; Nix, M. G. D.; Ashfold, M. N. R. High Resolution Photofragment Translational Spectroscopy Studies of the near Ultraviolet Photolysis of Imidazole. *J. Chem. Phys.* **2006**, *125*, 184302.

(204) Wan, J.; Hada, M.; Ehara, M.; Nakatsuji, H. Electronic Excitation Spectrum of Thiophene Studied by Symmetry-Adapted Cluster Configuration Interaction Method. *J. Chem. Phys.* **2001**, *114*, 842–850.

(205) Pastore, M.; Angeli, C.; Cimiraaglia, R. A Multireference Perturbation Theory Study on the Vertical Electronic Spectrum of Thiophene. *Theor. Chem. Acc.* **2007**, *118*, 35–46.

(206) Holland, D. M. P.; Trofimov, A. B.; Seddon, E. A.; Gromov, E. V.; Korona, T.; de Oliveira, N.; Archer, L. E.; Joyeux, D.; Nahon, L. Excited Electronic States of Thiophene: High Resolution Photoabsorption Fourier Transform Spectroscopy and *ab initio* Calculations. *Phys. Chem. Chem. Phys.* **2014**, *16*, 21629–21644.

(207) Di Lonardo, G.; Galloni, G.; Trombetti, A.; Zauli, C. Electronic Spectrum of Thiophen and Some Deuterated Thiophens. *J. Chem. Soc., Faraday Trans. 2* **1972**, *68*, 2009–2016.

(208) Varsányi, G.; Nyulászi, L.; Veszprémi, T.; Narisawa, T. Vibronic Analysis and Symmetry of the Lowest Energy Ultraviolet Transition of Thiophen. *J. Chem. Soc., Perkin Trans. 2* **1982**, 761–765.

(209) Haberkern, H.; Asmis, K. R.; Allan, M.; Swiderek, P. Triplet States in Oligomeric Materials: Electron Energy Loss Spectroscopy of Thiophene and Bithiophene and Extrapolation to the Polymer. *Phys. Chem. Chem. Phys.* **2003**, *5*, 827–833.

- (210) Palmer, M. H.; Walker, I. C.; Guest, M. F. The Electronic States of Thiophene Studied by Optical (VUV) Absorption, Near-Threshold Electron Energy-Loss (EEL) Spectroscopy and *ab initio* Multi-Reference Configuration Interaction Calculations. *Chem. Phys.* **1999**, *241*, 275–296.
- (211) Kleinschmidt, M.; Tatchen, J.; Marian, C. M. Spin-Orbit Coupling of DFT/MRCI Wavefunctions: Method, Test Calculations, and Application to Thiophene. *J. Comput. Chem.* **2002**, *23*, 824–833.
- (212) Del Bene, J. E.; Watts, J. D.; Bartlett, R. J. Coupled-Cluster Calculations of the Excitation Energies of Benzene and the Azabenzenes. *J. Chem. Phys.* **1997**, *106*, 6051–6060.
- (213) Nooijen, M. Similarity Transformed Equation of Motion Coupled-Cluster Study of Excited States of Selected Azabenzenes. *Spectrochim. Acta, Part A* **1999**, *55*, 539–559.
- (214) Sobolewski, A. L.; Woywod, C.; Domcke, W. *Ab Initio* Investigation of Potential-Energy Surfaces Involved in the Photophysics of Benzene and Pyrazine. *J. Chem. Phys.* **1993**, *98*, 5627–5641.
- (215) Christiansen, O.; Koch, H.; Halkier, A.; Jorgensen, P.; Helgaker, T.; Sánchez de Merás, A. Large-Scale Calculations of Excitation Energies in Coupled Cluster Theory: The Singlet Excited States of Benzene. *J. Chem. Phys.* **1996**, *105*, 6921–6939.
- (216) Packer, M. K.; Dalskov, E. K.; Enevoldsen, T.; Jensen, H. J.; Oddershede, J. A New Implementation of the Second-Order Polarization Propagator Approximation (SOPPA): The Excitation Spectra of Benzene and Naphthalene. *J. Chem. Phys.* **1996**, *105*, 5886–5900.
- (217) Hald, K.; Jørgensen, P.; Christiansen, O.; Koch, H. Implementation of Electronic Ground States and Singlet and Triplet Excitation Energies in Coupled Cluster Theory with Approximate Triples Corrections. *J. Chem. Phys.* **2002**, *116*, 5963–5970.
- (218) Li, Y.; Wan, J.; Xu, X. Theoretical study of the vertical excited states of benzene, pyrimidine, and pyrazine by the symmetry adapted cluster—Configuration interaction method. *J. Comput. Chem.* **2007**, *28*, 1658–1667.
- (219) Devarajan, A.; Gaenko, A. V.; Khait, Y. G.; Hoffmann, M. R. Generalized van Vleck Perturbation Theory (GVVPT2) Study of the Excited States of Benzene and the Azabenzenes. *J. Phys. Chem. A* **2008**, *112*, 2677–2682.
- (220) Sharma, P.; Bernales, V.; Truhlar, D. G.; Gagliardi, L. Valence  $\pi\pi^*$  Excitations in Benzene Studied by Multiconfiguration Pair-Density Functional Theory. *J. Phys. Chem. Lett.* **2019**, *10*, 75–81.
- (221) Doering, J. P. Low-Energy Electron Impact Study of the First, Second, and Third Triplet States of Benzene. *J. Chem. Phys.* **1969**, *51*, 2866–2870.
- (222) Nakashima, N.; Inoue, H.; Sumitani, M.; Yoshihara, K. Laser Flash Photolysis of Benzene. iii.  $S_n \leftarrow S_1$  Absorption of Gaseous Benzene. *J. Chem. Phys.* **1980**, *73*, 5976–5980.
- (223) Johnson, P. M. The Multiphoton Ionization Spectrum of Benzene. *J. Chem. Phys.* **1976**, *64*, 4143–4148.
- (224) Johnson, P.; Korenowski, G. The Discovery of a  $3p$  Rydberg State in Benzene by Three-Photon Resonant Multiphoton Ionization Spectroscopy. *Chem. Phys. Lett.* **1983**, *97*, 53–56.
- (225) Hiraya, A.; Shobatake, K. Direct Absorption Spectra of Jet-Cooled Benzene in 130–260 Nm. *J. Chem. Phys.* **1991**, *94*, 7700–7706.
- (226) Weber, P.; Reimers, J. R. *Ab Initio* and Density Functional Calculations of the Energies of the Singlet and Triplet Valence Excited States of Pyrazine. *J. Phys. Chem. A* **1999**, *103*, 9821–9829.
- (227) Woywod, C.; Papp, A.; Halász, G. J.; Vibók, Á. Theoretical Investigation of the Electronic Spectrum of Pyrazine. *Theor. Chem. Acc.* **2010**, *125*, 521–533.
- (228) Stanton, J. F.; Gauss, J. The First Excited Singlet State of *s*-Tetrazine: A Theoretical Analysis of Some Outstanding Questions. *J. Chem. Phys.* **1996**, *104*, 9859–9869.
- (229) Rubio, M.; Roos, B. O. A Theoretical Study of the Electronic Spectrum of *s*-Tetrazine. *Mol. Phys.* **1999**, *96*, 603–615.
- (230) Adamo, C.; Barone, V. A TDDFT Study of the Electronic Spectrum of *s*-Tetrazine in the Gas-Phase and in Aqueous Solution. *Chem. Phys. Lett.* **2000**, *330*, 152–160.
- (231) Nooijen, M. Electronic Excitation Spectrum of *s*-Tetrazine: An Extended-STEOM-CCSD Study. *J. Phys. Chem. A* **2000**, *104*, 4553–4561.
- (232) Angeli, C.; Cimraglia, R.; Cestari, M. A Multireference *n*-electron Valence State Perturbation Theory Study of the Electronic Spectrum of *s*-tetrazine. *Theor. Chem. Acc.* **2009**, *123*, 287–298.
- (233) Pastorczak, E.; Pernal, K. Electronic Excited States from the Adiabatic-Connection Formalism with Complete Active Space Wave Functions. *J. Phys. Chem. Lett.* **2018**, *9*, 5534–5538.
- (234) Bolovinos, A.; Tsekeris, P.; Philis, J.; Pantos, E.; Andritsopoulos, G. Absolute Vacuum Ultraviolet Absorption Spectra of Some Gaseous Azabenzenes. *J. Mol. Spectrosc.* **1984**, *103*, 240–256.
- (235) Okuzawa, Y.; Fujii, M.; Ito, M. Direct Observation of Second Excited  $^1, 3 (n, \pi^*)$  States of Pyrazine by UV-IR Double Resonance dip Spectroscopy. *Chem. Phys. Lett.* **1990**, *171*, 341–346.
- (236) Walker, I. C.; Palmer, M. H. The Electronic States of the Azines. IV. Pyrazine, Studied by VUV Absorption, Near-Threshold Electron Energy-Loss Spectroscopy and *Ab Initio* Multi-Reference Configuration Interaction Calculations. *Chem. Phys.* **1991**, *153*, 169–187.
- (237) Stener, M.; Decleva, P.; Holland, D. M. P.; Shaw, D. A. A Study of the Valence Shell Electronic States of Pyrimidine and Pyrazine by Photoabsorption Spectroscopy and Time-Dependent Density Functional Theory Calculations. *J. Phys. B: At., Mol. Opt. Phys.* **2011**, *44*, 075203.
- (238) Palmer, M. H.; McNab, H.; Reed, D.; Pollacchi, A.; Walker, I. C.; Guest, M. F.; Siggel, M. R. The Molecular and Electronic States of 1,2,4,5-tetrazine Studied by VUV absorption, Near-Threshold Electron Energy-Loss Spectroscopy and *ab initio* Multi-Reference Configuration Interaction Studies. *Chem. Phys.* **1997**, *214*, 191–211.
- (239) Palmer, M. H.; Walker, I. C. The Electronic States of the Azines. v. Pyridazine, Studied by VUV Absorption, Near Threshold Electron Energy-Loss Spectroscopy and *Ab Initio* Multi-Reference Configuration Interaction Calculations. *Chem. Phys.* **1991**, *157*, 187–200.
- (240) Linert, I.; Zubek, M. Electron Energy-loss Spectroscopy of Excited States of the Diazine Molecules: Pyridazine. *J. Electron Spectrosc. Relat. Phenom.* **2019**, *233*, 69–76.
- (241) Lorentzon, J.; Fülischer, M. P.; Roos, B. O. A Theoretical Study of the Electronic Spectra of Pyridine and Phosphabenzene. *Theor. Chim. Acta* **1995**, *92*, 67–81.
- (242) Walker, I. C.; Palmer, M. H.; Hopkirk, A. The Electronic States of the Azines. II. Pyridine, Studied by VUV Absorption, Near-Threshold Electron Energy Loss Spectroscopy and *Ab Initio* Multi-Reference Configuration Interaction Calculations. *Chem. Phys.* **1990**, *141*, 365–378.
- (243) Linert, I.; Zubek, M. Electron Energy-Loss Spectroscopy of Excited States of the Pyridine Molecules. *Eur. Phys. J. D* **2016**, *70*, 74.
- (244) Walker, I. C.; Palmer, M. H.; Ballard, C. C. The Electronic States of the Azines. VI. 1,3,5-Triazine, Studied by VUV Absorption, Near-Threshold Electron Energy-Loss Spectroscopy and *ab initio* Multi-Reference Configuration Interaction Calculations. *Chem. Phys.* **1992**, *167*, 61–75.
- (245) Fischer, G.; Wormell, P. Vibronic Analyses of the Lowest Singlet–Singlet and Singlet–Triplet Band Systems of Pyridazine. *Chem. Phys.* **2000**, *257*, 1–20.
- (246) Cai, Z.-L.; Reimers, J. R. The Low-Lying Excited States of Pyridine. *J. Phys. Chem. A* **2000**, *104*, 8389–8408.
- (247) Wan, J.; Hada, M.; Ehara, M.; Nakatsuji, H. Electronic Excitation and Ionization Spectra of Azabenzenes: Pyridine Revisited by the Symmetry-Adapted Cluster Configuration Interaction Method. *J. Chem. Phys.* **2001**, *114*, 5117–5123.
- (248) Innes, K.; Ross, I.; Moomaw, W. R. Electronic States of Azabenzenes and Azanaphthalenes: A Revised and Extended Critical Review. *J. Mol. Spectrosc.* **1988**, *132*, 492–544.
- (249) Serrano-Andrés, L.; Fülischer, M. P.; Karlström, G. Solvent Effects on Electronic Spectra Studied by Multiconfigurational Perturbation Theory. *Int. J. Quantum Chem.* **1997**, *65*, 167–181.
- (250) Öhrn, A.; Christiansen, O. Electronic Excitation Energies of Pyrimidine Studied using Coupled Cluster Response Theory. *Phys. Chem. Chem. Phys.* **2001**, *3*, 730–740.

(251) Fischer, G.; Cai, Z.-L.; Reimers, J. R.; Wormell, P. Singlet and Triplet Valence Excited States of Pyrimidine. *J. Phys. Chem. A* **2003**, *107*, 3093–3106.

(252) Palmer, M. H.; Walker, I. C.; Guest, M. F.; Hopkirk, A. The Electronic States of the Azines. iii. Pyrimidine, Studied by VUV Absorption, Near-Threshold Electron Energy-Loss Spectroscopy and ab initio Multi-Reference Configuration Calculations. *Chem. Phys.* **1990**, *147*, 19–33.

(253) Linert, I.; Zubek, M. A Study of the Electronic States of Pyrimidine by Electron Energy Loss Spectroscopy. *Chem. Phys. Lett.* **2015**, *624*, 1–5.

(254) Palmer, M. H.; Walker, I. C.; Guest, M. F.; Siggel, M. R. The Electronic States of the Azines. VII. 1,2,4-Triazine, Studied by Photon Absorption, Near-Threshold Electron Energy Loss Spectroscopy and ab initio Multi-Reference Configuration Interaction Calculations. *Chem. Phys.* **1995**, *201*, 381–391.

(255) Oliva, J. M.; Azenha, M. E. D. G.; Burrows, H. D.; Coimbra, R.; Seixas de Melo, J. S.; Canle L., M.; Fernández, M. I.; Santaballa, J. A.; Serrano-Andrés, L. On the Low-Lying Excited States of sym-Triazine-Based Herbicides. *ChemPhysChem* **2005**, *6*, 306–314.

(256) Jacquemin, D.; Duchemin, I.; Blase, X. Is the Bethe–Salpeter Formalism Accurate for Excitation Energies? Comparisons with TD-DFT, CASPT2, and EOM-CCSD. *J. Phys. Chem. Lett.* **2017**, *8*, 1524–1529.

(257) Eriksen, J. J.; Lipparini, F.; Gauss, J. Virtual Orbital Many-Body Expansions: A Possible Route towards the Full Configuration Interaction Limit. *J. Phys. Chem. Lett.* **2017**, *8*, 4633–4639.